
(a) exact/series ratio of sector-averaged coefficients, $b_0 = 0^\circ$
exact / series coefficient
D/r
τ_0/q_0 (monopole)
$\tau_1/(\varepsilon o_1)$ (dipole)
τ_2/q_2 (quadrupole)
$\tau_3/(\varepsilon o_3)$ (octupole)
(b) exact/series ratio of sector-averaged coefficients, $b_0 = 19.1^\circ$
exact / series coefficient
D/r
τ_0/q_0 (monopole)
$\tau_1/(\varepsilon o_1)$ (dipole)
τ_2/q_2 (quadrupole)
$\tau_3/(\varepsilon o_3)$ (octupole)
(c) predicted octupole once m_0, m_1, m_2 are matched: exact vs series
predicted A_3
D/r (held fixed)
data $A_3 \pm 1\sigma$ (sector-correlated)
data $A_3 \pm 1\sigma$ (naive)
exact: match m_0-m_2, on-axis
exact: match m_0-m_2, flipped
series: match m_0-m_2, on-axis
series: match m_0-m_2, flipped
(d) exact point-mass curves with rotation + 2ℓ transforms
$A/A_{\rm all}$
Galactic longitude ℓ [deg]
data Fourier fit ($m \le 3$)
exact free, on-axis: R^2=0.857
exact D/r=2: R^2=0.887
exact D/r=3: R^2=0.863
exact D/r=4: R^2=0.848
Migkas Fig. 6