Centre-of-Expansion shell cosmology: from Migkas anisotropy to a flat CMB

Chris Alameddin

29 September 2026

Directional Hubble measurements from Migkas et al.[1] fix an off-centre Sedov–Taylor[2, 3] / Lemaitre–Tolman–Bondi (LTB)[4, 5, 6] shell geometry. The same geometry is then tested against the cosmic microwave background (CMB). The CMB is treated as an independent consistency check, not as a fit target[7].

1 Problem and stance

Migkas et al. report a sky-dependent cluster Hubble rate. Standard cosmology assumes large-scale isotropy, and the CMB is isotropic to $\sim 10^{-5}$[7]. This letter takes both seriously.

2 Best-fit geometry

Local Hubble velocity is free-fall toward the CoE minus our motion, $\vec{V}_H = \vec{V}_g - \vec{V}_O$. The observer ("US") sits inside a thick shell; vision radius $R \approx 14.3\,\mathrm{Gpc}$. Preferred relativistic fit: $D \approx 32.9\,\mathrm{Gpc}$, $R_{\mathrm{in}} \approx 18.1\,\mathrm{Gpc}$, $R_{\mathrm{out}} \approx 49.7\,\mathrm{Gpc}$.




Figure 1: Geometry: CoE, observer (US) inside a thick shell, and the vision sphere of radius R.

3 Approach

1. Migkas A/A_{all} vs. Galactic ℓ (and Fig. 8); slice fits by redshift.
2. Fit CoE + thick shell (Newtonian and relativistic LTB)[4, 5, 6].
3. Orient on the sky: untilted $\rightarrow$ $+15°$ tilt $\rightarrow$ gradient $\rightarrow$ $+80.8°$ roll; match Migkas Fig. 8.
4. Donut-universe evolution: shells $> c$, visible at c, fixed recombination surface.
5. CMB stress test: gravitational multipole $\rightarrow$ kill function $\rightarrow$ two-emitter construction.

Table 1: Newtonian and relativistic shell+core fits to Migkas $A(\ell)$.

	Newt. Ω_s free	Newt. $\Omega_s = 0.049$	Rel. LTB $\Omega_s = 0.049$
D (Gpc)	93.7	49.2	32.9
Ω_s	1.01	0.049	0.049
l_0 (deg)	12.8	13.0	13.5
R^2 vs Migkas	0.909	0.893	0.885
R^2 vs A_F	0.987	0.977	0.963




Figure 2: Migkas Fig. 6: $A/A_{\rm all}$ versus Galactic longitude, with Newtonian and relativistic shell+core model curves.




Figure 3: Migkas $A(\ell)$ in three redshift slices (0.03–0.07, $z > 0.07$, and all-z), with the CoE model recomputed at each slice's median distance, plus the D–core-mass degeneracy.

4 Einstein and Friedmann equations for the CoE+shell

The LTB metric is written about the CoE, not about us. The observer (US) sits at comoving distance D from that centre. Sky anisotropy is an off-centre sampling of radial structure, not a direction-dependent scale factor $a(t, \theta)$. Einstein's equation is

$$G_{\mu\nu} = 8\pi G\, T_{\mu\nu}. \tag{1}$$

For concentric dust shells about the CoE the Lemaitre–Tolman–Bondi line element is[4, 5, 6]

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + \frac{(R')^2}{1 + 2E(r)}\,\mathrm{d}r^2 + R(t, r)^2\,\mathrm{d}\Omega^2, \tag{2}$$

with areal radius $R(t, r)$, $R' \equiv \partial_r R$, and $\dot{R} \equiv \partial_t R$. Each shell obeys the Friedmann-like constraint

$$\dot{R}^2 = 2E(r) + \frac{2GM(r)}{R}. \tag{3}$$

The interior mass uses a background density scale ν and a radial mass multiplier $\beta(r)$ (not a scale factor),

$$M(r) = \frac{4\pi}{3} R^3 \nu \beta(r), \tag{4}$$

so $\beta = 1$ recovers critical-density normalisation. In the CoE+shell geometry $\beta(r)$ is small in the cavity, rises through the thick shell, and returns toward the exterior background beyond $R_{\rm out}$.

Including radiation on the same radial scaffolding,

$$H_a^2 \propto \frac{\nu_m \beta_m(r)}{a^3} + \frac{\nu_r \beta_r(r)}{a^4} - \frac{k(r)}{a^2}, \tag{5}$$

with $H_a = \dot{a}/a$ and $a(t, r)$ the local expansion factor of each shell. Early times are radiation-dominated ($a \sim t^{1/2}$); late times are dust-dominated ($a \sim t^{2/3}$). Curvature $k(r)$ follows from $E(r)$ and introduces no angular dependence into the metric coefficients.

For the Newtonian $\Omega_s = 0.049$ CoE+shell fit in Table 1, the solved geometry matching Migkas $A(\ell)$ is

$$D \approx 49.17\,\text{Gpc}, \quad R_{\rm in} \approx 34.48\,\text{Gpc}, \quad R_{\rm out} \approx 70.27\,\text{Gpc}, \quad \Omega_s = 0.049, \quad \ell_0 = 13^\circ.$$

The shell rule with vision radius $R \approx 14.3\,\text{Gpc}$ requires $D \gtrsim 14.7\,\text{Gpc}$; the fitted D lies above that bound with the observer mid-shell between $R_{\rm in}$ and $R_{\rm out}$.




Figure 4: Full Migkas $A/A_{\rm all}$ profile and residual panel for the Newtonian and relativistic fits.

5 Sky orientation

The model must be placed on the Galactic sky. Starting from an untilted, no-roll configuration, we apply a +15° tilt, then a radial gradient, then an +80.8° roll about the CoE axis. The predicted MAX/MIN loci are compared with Migkas Fig. 8 in the following section.




Figure 5: Sky orientation sequence: untilted; +15° tilt; tilt with gradient; tilt, gradient, and +80.8° roll. Each row shows the model frame, the Mollweide sky map, and a 3D view.




Figure 6: Model MAX/MIN bands overlaid on Migkas Fig. 8 for +15° tilt; tilt with gradient; and tilt, gradient, and +80.8° roll.

6 Time evolution and the CMB

In the donut-universe picture the CoE is fixed while thick shells expand outward faster than light in coordinate sense; the sphere of events visible to us grows at c. Recombination is a spherical surface of fixed physical radius around the CoE. The observer remains mid-shell between the inner and outer edges (S1 and S2). Frames A–D show shells crossing that recombination surface while the visible sphere grows more slowly.




Figure 7: Donut-universe time sequence (A–D): fixed recombination surface; shells expand through it; S2 is filled in every frame; the observer sits mid-shell.




Figure 8: Two-emitter baseline. If opposite plates (or shells) share the observer's bulk velocity, the kinematic sky is flat before gravitational multipoles are included.

The same off-centre geometry produces a gravitational multipole on the last-scattering sky. The temperature pattern required to cancel that multipole—the *kill function*—is minus the Doppler residual as a function of angle from the CoE. Exploratory two-emitter and multipole-partner constructions follow the kill-function morphology at large D and remove most of the demeaned power through $\ell \leq 4$, but they do not yet reach the CMB bound of order 10^{-5}.




Figure 9: Unit-normalized kill function compared with exploratory plate, shell, and shear residuals at three distances D.




Figure 10: CoE Doppler term plus a multipole partner: residuals after a dipole+P_2 cancellation and after an $\ell \leq 4$ partner.

7 What is interesting

The Migkas anisotropy is a gravitational effect: free-fall relative to an off-centre CoE. On the last-scattering sky that same gravity produces a temperature multipole. Separately, a pure electromagnetic Doppler pattern—from radial motion of emitting surfaces—has nearly the same angular shape. The *kill function* is the gravitational residual written with the opposite sign. Today's result is that this gravitational kill tracks the electromagnetic Doppler morphology across the distances we scanned. That is unexpected if gravity and the photon bath are

unrelated, and it suggests a geometric route to CMB flatness that does not throw away the Migkas fit.

8 What we do next

1. Evolve $D(t)$, $R_{\rm in}(t)$, and $R_{\rm out}(t)$ from the present back through recombination, with the observer mid-shell.
2. Integrate null geodesics on that history to obtain $T(\theta)$ on our sky, including gravitational and kinematic contributions together.
3. Ask whether a shared bulk velocity of the emitting surfaces, plus the gravitational multipole, can keep $\ell \geq 2$ below $\sim 10^{-5}$ while the low-redshift $A(\ell)$ pattern remains Migkas-compatible.
4. If a residual remains, identify the smallest additional degree of freedom (timing of recombination, shell thickness history, or a mild partner multipole) that removes it without reintroducing a fixed-curve conflict across redshift slices.

9 Bottom line

Migkas fixes an off-centre CoE+shell that fits directional Hubble across slices. Gravity on that geometry predicts a last-scattering multipole whose kill function matches the shape of an electromagnetic Doppler pattern from the same radii. The open problem is no longer whether the shapes talk to each other—they do—but whether a full time-dependent light-cone calculation can make them cancel at the 10^{-5} level without spoiling the Migkas anisotropy.

References

[1] K. Migkas, G. Schellenberger, T. H. Reiprich, F. Pacaud, M. E. Ramos-Ceja, and L. Lovisari, Astron. Astrophys. **636**, A15 (2020); arXiv:2004.03305.

[2] L. I. Sedov, J. Appl. Math. Mech. **10**, 241 (1946).

[3] G. I. Taylor, Proc. R. Soc. Lond. A **201**, 159 (1950).

[4] G. Lemaître, Ann. Soc. Sci. Bruxelles **A53**, 51 (1933).

[5] R. C. Tolman, Proc. Natl. Acad. Sci. USA **20**, 169 (1934).

[6] H. Bondi, Mon. Not. R. Astron. Soc. **107**, 410 (1947).

[7] Planck Collaboration, Astron. Astrophys. **641**, A7 (2020).

Acknowledgments

I thank Luke for the important question that sharpened this work, and Dr. Rendler for his support and guidance. Computational assistance is gratefully acknowledged.

Respectfully,

Chris Alameddin

Appendix

Shell expansion and pointing at the CoE

The LTB shell equation integrates to the usual early radiation and late matter asymptotics, with the sign of $E(r)$ fixing whether a dust shell expands forever or recollapses. Figure A1 collects those solutions for a flat matter+radiation track, dust with $E = 0$, $E > 0$, and $E < 0$.




Figure A1: LTB shell evolution. The flat matter+radiation track is shown with the $E = 0$ dust solution and the $E > 0$ (expand-forever) and $E < 0$ (recollapsing) branches.

Because the CoE is a fixed direction on the sky in our geometry, it is a place you can point to. From Central Texas (where this work was done), that direction rises in the northeast and sets in the northwest on a sidereal schedule fixed by the fitted CoE coordinates. Figure A2 compares CoE rise, culmination, and set with the Sun through 2026. Table A1 lists the nine Sun–CoE event crossings for 2027; the highlighted diagonal is rise with rise, highest with highest, and set with set.




Figure A2: CoE and Sun timing, Plano 2026.

Sun / CoE Event Crossings — Central Texas (2027)

Sun ↓ / CoE →	CoE Rise	CoE Highest	CoE Set
Sunrise	**Jun 26 6:40 AM NE · 0°**	Oct 9 7:38 AM N · 83°	Jan 21 7:38 AM NW · 0°
Sun Highest	Mar 9 12:48 PM NE · 0°	**Jul 8 1:43 PM N · 84°**	Nov 10 12:21 PM NW · 0°
Sunset	Dec 26 5:37 PM NE · 0°	Apr 5 7:54 PM N · 83°	**Jul 23 8:34 PM NW · 0°**

Highlighted diagonal: rise–rise · highest–highest · set–set

Table A1: Sun/CoE crossings, Central Texas 2027.

Figure A3 gives the annual track of the Centre of Expansion (CoE) on the local sky as seen from Plano at 12:00 local mean time on each day of 2026. The month labels make the calendar progression explicit: as the Earth moves through its orbit, one fixed celestial direction traces a large oval in local horizon coordinates. This is a locus on the sky, not the Sun's familiar figure-eight (analemma).




Figure A3: Annual local-sky path of the Centre of Expansion (CoE) from Plano, Texas, evaluated at 12:00 local mean time on each day of 2026. The colour gradient and labelled dates mark the progression through the year; the large oval is the locus of the fixed celestial direction in local horizon coordinates, not a solar figure-eight.

