Unexpectedly low marine heatwave forecast skill during the 2023–2024 El Niño

Supplementary Methods and data coverage

Scope of the supplementary analyses

These analyses address three questions: whether MHW forecast performance differs from historical expectations; how tropical and remote atmospheric forecasts differ; and how regional SST associations enter the diagnostic allocation of the regional map difference. Figures 1–4 and 16–17 assess MHW intensity, occurrence and sampling coverage; Figs. 5–11 examine tropical and remote atmospheric forecasts; Figs. 12–15 evaluate regional SST associations, their temporal transfer, a single-model process comparison and the signed map accounting. Supplementary Table 1 identifies the data and sampling support of each analysis.

Events, repeated samples and uncertainty

The earlier strong events are 1997–1998 and 2015–2016; the focal event is 2023–2024. The nine-event amplitude regression additionally uses 1991–1992, 1994–1995, 2002–2003, 2004–2005, 2006–2007, 2009–2010 and 2018–2019. Month, lead, model and spatial-grid replication do not create additional independent ENSO events. In Fig. 2, the same 2,000 resampled calendar-year blocks are used jointly for both phases, yielding percentile intervals for correlations and their slope contrast. Prediction intervals in Fig. 1 describe a new event under the linear regression assumptions. Lead and historical-reference ranges elsewhere are descriptive ranges, not confidence intervals. No new multiple-comparison significance claims are made.

AUC and conditional intensity errors

Global AUC in main Fig. 1a,b remains the original monthly spatial discrimination metric. The spatial sensitivity analyses in Supplementary Figs. 1c and 16–17 pool raw cell-month samples within an 11 × 11 array of 1° ocean grid cells (centre ±5 grid centres), restricted to the same major ocean basin and 60°S–60°N. Longitude wraps periodically. The primary window is April of the event year through September of the next year (18 months). Forecast records from the original July–June window are reused without modification; additional months use the same archived reader. Neighbour cells must have complete paired records for all events and leads. For each event and lead, weighted ROC AUC ranks forecast SST minus its local calendar-month Q90 against observed SST above Q90, with cosine-latitude sample weights and half credit for ties. A pool with only one observed class is undefined. Canonical-minus-target differences use equal weights for the two historical events and nine leads; only centres with all 27 finite AUCs enter the map. No interpolation or smoothing of the computed AUC is applied. Overlapping neighbourhoods reduce effective spatial independence despite the unchanged 1° output grid. Figure 17 separately checks 12 versus 18 months and single-cell versus 11° support. Forecast/observed intensity is max(SST − Q90,0); squared errors are pooled on observed-MHW samples only, using the same neighbourhood and time support. Roots follow MSE aggregation, including equal-event canonical averaging and region averaging. Regional neighbourhood metrics cannot be summed to recover the original global AUC gap. No claim of increased independent sample size is made.

Forecast extraction shared with the main SST figures

Figures 1–4 and 15–17 retain the archived SST extraction used in main Figs. 1 and 4. That extraction averages stored initialization coordinates in some selected forecast files. A file-level audit has identified cases where filename-based and exact initialization-coordinate selections differ, including GFDL files for September and December 2023 with conflicting source-coordinate attribution. Consequently these supplementary SST panels describe the retained analysis, not an independently initialization-controlled validation.
Exact-coordinate extraction changes the regression-based global deficit in the separately available sensitivity analysis; it is not silently substituted here. The magnitude of the SST-derived skill differences is therefore conditional on this extraction rule.

Tropical correlation comparisons

a,c,e, Cosine-latitude-weighted spatial correlations between forecast and observed DJF anomalies for SST, zonal wind stress and convection, represented by precipitation. All variables use NASA GEOS-S2S, the same four ensemble members (1–4), the same target months and leads, and the same tropical Pacific ocean mask (20°S–20°N, 140°E–100°W; 4,833 grid cells). The two historical strong El Niño events are shown separately. b,d,f, Relative correlation decrease: 100 × (r_reference − r_2023–2024)/r_reference, using each historical event and their equal-event mean. Positive values indicate deterioration; negative values indicate improvement. All right-hand panels share the same vertical scale. Lead 9 is unavailable and remains blank/NA; no nearest-lead substitution is used. Forecast initialization equals the verifying month minus the labelled lead; the internal S coordinate must match that initialization and L must equal lead + 0.5. Monthly anomalies are calculated against 1992–2020, with separate lead-dependent model and observed climatologies using identical available calendar months. DJF is averaged before spatial correlation. The same ocean mask is used for all 72 valid event–variable–lead correlations. Eight leads within each of three events are repeated evaluations, not independent events; reference curves are sensitivity comparisons, not confidence intervals. This is a controlled single-model comparison, not a multimodel attribution of MHW forecast-skill loss. Across eight leads, mean correlations change from 0.933 to 0.668 for SST, from 0.764 to 0.273 for zonal wind stress and from 0.810 to 0.612 for convection. Thus this matched comparison supports particularly strong wind-stress deterioration, but does not show greater deterioration in both atmospheric variables than in SST. The earlier mixed-protocol figure is retained in the preceding release, not used for this strict comparison.

Exact-coordinate DJF circulation comparisons

Figures 6–11 use DJF targets and exact available future leads 1–8 for COLA–RSMAS–CCSM4 (members 1–10), NASA–GEOS-S2S (1–4) and GFDL–SPEAR (1–15). Equal weights are applied to members within a model, then to model fields. All required fields must be finite. NCEP–CFSv2 is not in this fixed source set because its 1997–1998 Z200 product is unavailable there; NASA limits the common exact lead range to 1–8. Each calendar-month climatology spans 1992–2020, excluding February 2018 uniformly after the corresponding NASA field failed the availability check, leaving 29 December, 29 January and 28 February samples. No target-event DJF month is removed. The spatial correlation centers each field before cosine-latitude-weighted covariance; RMSE retains amplitude and sign-independent error magnitude. Pooled tropical–remote associations include between-event differences; their within-event counterparts cannot be substituted for them.

Region definitions

For main Figs. 3–4 and Supplementary Figs. 7 and 12–15, NOAA WOA major-basin identity is intersected with latitude bands: TP/TA, 23.5°S–23.5°N; NP/NA, 23.5°N–60°N; SP/SA, 60°S–23.5°S. IO spans the Indian Ocean from 60°S to its northern boundary. Grid-centre membership assigns −23.5° to the tropical band and +23.5° to the northern band, with no overlap. NP, SP, TA, NA, SA and IO are the six atmospheric recipient ocean masks. Global circulation comparisons additionally include land. The spatial sensitivity analyses in Supplementary Figs. 1c and 16–17 use the retained ±20° split; they are shown explicitly as a different support, not silently relabelled as the process regions.

Historical SST association and reconstruction

Figures 12–13 use regional ERSST SST and ORAS5 mixed-layer temperature, surface heating and horizontal advection. The fitting record is January 1985–December 2020 (432 months). ENSO is a separate group; IOD, PDO, NAO and SAM form the other-mode group. Local variables are previous-month SST and mixed-layer temperature, contemporaneous net surface heating divided by mixed-layer heat capacity, and horizontal advection. Full-sample month effects and linear trend are removed for the in-sample variance allocation. Group Shapley values average marginal explained variance over group orders; nested Owen values split the local parent while preserving group membership. The unresolved fraction is 1 − full-model R^2. All shares sum to one. For forward reconstruction, the seasonal cycle, trend, predictor scaling and regression are trained only before each test block. Test-time contemporaneous observed predictors make this diagnostic reconstruction rather than an actionable forecast.

CFSv2-only process check

Figure 14 uses a separately retrieved CFSv2 process archive, four members (00, 06, 12 and 18), DJF and leads 1, 3 and 6 months. Verification uses ERA5 upper-level circulation and ORAS5 net ocean surface heat flux. The circulation quantity is the departure from contemporaneous zonal mean, not the calendar-month anomaly used in main Fig. 3. Net surface flux is positive downward into the ocean. Squared errors are averaged across the three months and cosine-latitude-weighted common ocean cells before taking the root. Each RMSE is divided by the mean of the two earlier-event RMSEs at the same region and lead. The regional masks have been recomputed at ±23.5°. CFSv2 is also one component of the main SST ensemble; this is therefore a separately processed single-model consistency check, not a model-independent validation of the mechanism.

Exact relationship between Supplementary Fig. 15 and main Fig. 4

Let $d(g) = 0.7050185255 - \mathrm{AUC}(g)$, where AUC(g) is the nine-lead mean of the local 18-month, 11°-neighbourhood AUC, and let a(g) be cosine-latitude area weights normalized over all evaluable map cells. For region b, $L(b) = \sum a(g) \max[d(g),0]$ and $H(b) = \sum a(g) \max[-d(g),0]$. The common denominator is $N = \sum_b [L(b) - H(b)] = 0.1065143704$. The positive and negative percentages are $100\, L(b)/N$ and $-100\, H(b)/N$. They sum to +120.790563% and −20.790563%, respectively, and hence 100% net. Historical SST-association weights w(b,k) sum to one within each region; component amounts are $\sum_b L(b)w(b,k)$ and $-\sum_b H(b)w(b,k)$. Local subcomponents replace, rather than duplicate, their parent allocation. The global reference is not an estimate of each grid cell's expected AUC. The map aggregate differs from main Fig. 1's globally pooled monthly spatial-AUC deficit (0.1090497372); these are distinct estimands, with different ranking samples and aggregation order. Exact closure verifies accounting, not causal attribution. Source tables provide regional and component crosswalks to main Fig. 4.


Supplementary Table 1 | Data coverage

Figures	Variables, sources and periods	Spatial support, samples and leads
S1, S16–S17	SST; conditional intensity RMSE; pooled temporal AUC ERSST + retained four-model SST extraction Q90:1985–2014; maps:April–September next year; S17 also July–June	Ocean ≤60°; same-basin 11° neighbourhood; inset masks ±20° S1a:10 events; S1b:404 months; maps:3 events ×18 months ×9 leads Lead labels 1–9; single-class/invalid pools missing; original extraction limits retained
S2–S4	Monthly spatial AUC; Niño3.4 Same archived four-model SST ensemble as main Fig. 1 Monthly record; nine historical events + focal event	Global evaluation mask S2: phase-eligible months; S3: 9 events; S4: 1,080 records Leads 1–9; 12 verifying months per event; no independent-month inference
S5	SST, zonal wind stress, convection correlations NASA GEOS-S2S members 1–4; ERSST and ERA5 DJF 1997–98, 2015–16, 2023–24; climatology 1992–2020	20°S–20°N, 140°E–100°W; 4,833 common ocean cells 3 events × 3 variables × 8 leads = 72 correlations Exact starts and leads 1–8; lead 9 unavailable for all variables
S6–S9	200-hPa height; tropical precipitation for S9 ERA5; CCSM4, GEOS-S2S, SPEAR DJF; 1992–2020 monthly means, February 2018 excluded	S6/S8: all longitudes ≤60° incl. land; S7/S9: six ±23.5° recipient masks 3 events; 8 leads; S9: 144 repeated pairs Exact leads 1–8; NASA lacks 9; CFSv2 historical Z200 absent from this source set

Supplementary Table 1 | Data coverage (continued)

Figures	Variables, sources and periods	Spatial support, samples and leads
S10–S11	Height anomaly patterns, amplitude, MSE Same exact-coordinate three-model/ERA5 products DJF and common anomaly baseline	S10a global; S10b/c: 30–60°N, 40–60°S, 120°E east to20°E; S11:45°N,50°S 3 strong events; S10a additionally 6 other El Niño winters Means of exact leads 1–8; six reference winters:1994,2002,2004,2006,2009,2018
S12–S13	SST; indices; lagged SST/MLT; heating; advection ERSST, climate-index series and ORAS5 1985–2020; month effects and linear trend	Seven disjoint ±23.5° analysis regions 432 months/region; three forward test blocks No forecast lead axis; contemporaneous observed predictors explicitly diagnostic
S14	CFSv2 height and net surface heat-flux errors Separate CFSv2 process download; ERA5/ORAS5 verification DJF; normalization by two earlier event RMSEs	Seven ±23.5° regions; fixed common finite ocean cells 3 events × 7 regions × 3 leads; 54 remote pairs Leads 1,3,6; four members; not an independent model from main SST ensemble
S15	Local AUC differences and historical SST-association weights Same spatial AUC map and association weights as main Fig. 4 April 2023–September 2024; weights:1985–2020	Seven ±23.5° regions + Other seas; 11° same-basin pools 27,508 valid map centres; nine leads; weights:432 months/region Undefined AUC excluded; net map denominator 0.1065143704; distinct from Fig. 1 global AUC gap


Supplementary Fig. 1 | MHW-intensity forecast errors in time and space



a, Event-mean MHW-intensity root-mean-square error (RMSE) versus peak Niño3.4 anomaly. The regression and grey 95% prediction interval use nine earlier El Niño events; the red star marks 2023–2024, excluded from fitting. b, Monthly global intensity RMSE (404 months); shading marks the three strong events. c, Intensity RMSE in 2023–2024 minus the reference from 1997–1998 and 2015–2016, using 18-month windows and 11° × 11° same-basin neighbourhoods. Red indicates larger errors and blue smaller errors; grey ocean cells lack a valid comparison. The inset summarizes all seven analysis regions, retaining all nine leads and their range. TP, tropical Pacific; NP, North Pacific; SP, South Pacific; TA, tropical Atlantic; NA, North Atlantic; SA, South Atlantic; IO, Indian Ocean. Intensity is positive SST excess above the local 1985–2014 calendar-month 90th percentile. Verification includes only observed-MHW samples: smaller RMSE indicates better intensity prediction but does not assess false alarms outside observed MHWs.

Spatial windows run from April of the event year to September of the next year. The reference RMSE is the root of the equal-event mean squared error; regional summaries also aggregate squared errors before taking roots. Lead ranges are not confidence intervals. Panels a,b retain monthly spatial aggregation. The 1° display grid does not imply independent 1° information. Coverage and pooling sensitivity are shown in Supplementary Figs. 16–17.


Supplementary Fig. 2 | ENSO-phase associations with MHW forecast skill



a,b, Global MHW area under the ROC curve (AUC) against absolute Niño3.4 anomaly during El Niño and La Niña months, respectively. The thresholds are +0.5 °C and −0.5 °C; neutral months are outside this comparison. Lines are ordinary-least-squares fits and r is the monthly Pearson correlation. Both panels use the same AUC axis. The 95% correlation intervals are percentile intervals from 2,000 whole-calendar-year resamples; the same sampled years are used for both phases when estimating the slope difference. Months within years and successive ENSO months are dependent; they are not independent events. Year-block resampling addresses within-year dependence but may not capture the full duration of multiyear ENSO episodes. The 95% year-block intervals for the correlations are 0.31 to 0.78 for El Niño and −0.13 to 0.39 for La Niña. The El Niño minus La Niña slope difference is 0.062 AUC °C□¹ (95% year-block interval -0.011 to 0.128); the interval includes zero.

All phase-eligible months are retained; no independent-month P values or multiple-comparison significance symbols are used.


Supplementary Fig. 3 | Sensitivity of the ENSO–skill relationship to event selection



a, Mean AUC versus midpoint year for the nine earlier El Niño events. b, Niño3.4 amplitude and AUC after removing their respective linear relationships with event year. c, The Niño3.4 coefficient in AUC = intercept + amplitude coefficient × Niño3.4 + year coefficient × event year, refitted after omitting each named event. Each point uses eight events; the red dashed line is the coefficient from all nine events. The 2023–2024 event is excluded throughout. These nine omission estimates describe sensitivity to event selection; their spread is not a confidence interval and they are not nine independent experiments. Adjustment for event year assesses linear temporal confounding only.

Nine historical events; deterministic ordinary-least-squares fits. Event amplitudes and AUCs are identical to main Fig. 1a.


Supplementary Fig. 4 | MHW forecast skill across lead months and event development



a, AUC expected from earlier events (dark line) and realized in 2023–2024 (red line), evaluated separately at each lead from 1 to 9 months. Connecting vertical segments show the expected-minus-realized difference. At each lead, AUC is averaged over July–June and regressed on El Niño amplitude using the nine earlier events; 2023–2024 is excluded. b, AUC averaged over leads 1–9 for each verifying month in the three strong events. The light-grey, dark-grey and red lines denote 1997–1998, 2015–2016 and 2023–2024. All 1,080 event–month–lead records are used: ten events × twelve months × nine leads. Leads and months are repeated evaluations, not independent events.

No lead-specific significance test is claimed. These panels retain the forecast extraction and monthly spatial-AUC definition of main Fig. 1a,b.


Supplementary Fig. 5 | Tropical forecast skill under matched model, season and lead conditions



a,c,e, Cosine-latitude-weighted spatial correlations between forecast and observed DJF anomalies for SST, zonal wind stress and convection, represented by precipitation. All variables use NASA GEOS-S2S, the same four ensemble members (1–4), the same target months and leads, and the same tropical Pacific ocean mask (20°S–20°N, 140°E–100°W; 4,833 grid cells). The two historical strong El Niño events are shown separately. b,d,f, Relative correlation decrease: 100 × (r_reference − r_2023–2024)/r_reference, using each historical event and their equal-event mean. Positive values indicate deterioration; negative values indicate improvement. All right-hand panels share the same vertical scale. Lead 9 is unavailable and remains blank/NA; no nearest-lead substitution is used.

Forecast initialization equals the verifying month minus the labelled lead; the internal S coordinate must match that initialization and L must equal lead + 0.5. Monthly anomalies are calculated against 1992–2020, with separate lead-dependent model and observed climatologies using identical available calendar months. DJF is averaged before spatial correlation. The same ocean mask is used for all 72 valid event–variable–lead correlations. Eight leads within each of three events are repeated evaluations, not independent events; reference curves are sensitivity comparisons, not confidence intervals. This is a controlled single-model comparison, not a multimodel attribution of MHW forecast-skill loss.


Supplementary Fig. 6 | Observed and forecast upper-level circulation in three El Niño events



Rows show December–February (DJF) of 1997–1998, 2015–2016 and 2023–2024; columns show ERA5 200-hPa geopotential-height anomalies, multimodel forecast anomalies and forecast minus observed anomalies. Forecasts first average members within each model, then equally weight the three available models and the eight common future leads. Observed and forecast fields share the ±160 m colour scale; differences share ±100 m, with extensions marking values beyond the limits. Red denotes positive height anomalies or positive forecast error, blue negative values. Anomalies use calendar-month means from 1992–2020; February 2018 is omitted consistently from all model and observation climatologies because the corresponding GEOS-S2S field is missing. No event DJF month is excluded. Lead 9 is absent from the fixed three-model cohort. Maps retain their native 1° resolution without added smoothing.

Models are COLA–RSMAS–CCSM4, NASA–GEOS-S2S and GFDL–SPEAR. These are seasonal composites, not isolated causal ENSO responses. Signed averaging can cancel opposing errors; Fig. 8 also evaluates RMSE.


Supplementary Fig. 7 | Regional circulation forecast skill across lead months



a–c, Forecast–observation spatial correlations of DJF 200-hPa height anomalies in 1997–1998, 2015–2016 and 2023–2024. d, Correlation in 2023–2024 minus the equal-event mean of the two earlier events. Panels a–c share the −1 to 1 scale; in d, red denotes lower and blue higher correlation. Grey cells labelled NA denote unavailable lead 9, not zero skill; 48 region–lead combinations are valid per event. NP, North Pacific; SP, South Pacific; TA, tropical Atlantic; NA, North Atlantic; SA, South Atlantic; IO, Indian Ocean. Here NA on a row label is North Atlantic, whereas NA inside a cell means not available. Pacific and Atlantic masks use ±23.5° and ±60° boundaries as in main Figs. 3–4; tropical Pacific is the source and is omitted from this six-recipient comparison. The full boundaries are in the coverage table.

These six ocean-mask diagnostics are not four idealized PNA/PSA/Indian/Atlantic waveguide boxes. No significance is inferred from the number of leads or grid cells.


Supplementary Fig. 8 | Circulation forecast correlations and RMSE across models



a, Global spatial correlation of forecast and observed DJF 200-hPa height anomalies. b, Corresponding area-weighted RMSE. Each row is one model or the equal-weight multimodel mean. Colours denote 1997–1998 (light grey), 2015–2016 (dark grey) and 2023–2024 (red). Large points show the mean over eight leads; small points retain all lead values; whiskers span their minimum and maximum. The multimodel mean is formed from fields before verification. The domain is all longitudes, 60°S–60°N, including land, weighted by cosine latitude. NCEP–CFSv2 is absent because 1997–1998 200-hPa data were unavailable in the common source set used for main Fig. 3; it is assessed separately in Fig. 14.

The plot includes 72 single-model and 24 multimodel event–lead evaluations from three events. Lead ranges are not confidence intervals. RMSE squares spatial errors before averaging and is therefore not reduced by cancellation of positive and negative errors.


Supplementary Fig. 9 | Links between tropical and remote forecast errors depend on the El Niño event



a–c, Tropical Pacific precipitation forecast–observation correlation versus remote 200-hPa height forecast–observation correlation, shown separately for the three events. Symbols identify the six recipient regions defined in Fig. 7. Each panel contains all 48 region–lead pairs; r summarizes their within-event association. d, The pooled association for all events and after omitting each event. The pooled relationship compares both differences between events and differences within events, whereas a–c assess the latter alone. Each tropical lead value is repeated for six recipients. Thus 144 plotted pairs represent three event clusters, not 144 independent events; the weak within-event associations cannot establish a stable predictive relationship or one-way transmission of forecast error.

All values, including negative circulation correlations, are retained. Event omission is a sensitivity analysis, not out-of-event prediction validation. No independent-sample P value is reported.


Supplementary Fig. 10 | Circulation pattern similarity, amplitude and error relative to climatology



a, Global anomaly correlations of the 2023–2024 forecast (red) and the forecast composite of the two earlier strong events (dark grey) against each of six other El Niño winters, identified by start year. These are events in the main Fig. 1 list within the historical-reference window 1992–1993 to 2021–2022, excluding 1997–1998 and 2015–2016; 1991–1992 is outside that window. b, Forecast-to-observed spatial standard-deviation ratio of height anomalies. c, Skill against a zero-anomaly forecast, defined as 1 − MSE_forecast/MSE_zero-anomaly. Values above zero indicate lower error than climatology. Panels b,c use 30°–60°N and 40°–60°S along the sector from 120°E eastward to 20°E. The dashed line in b marks equal forecast and observed amplitude.

Individual-event correlations are not equivalent to correlation with the averaged observed field. A smaller forecast anomaly amplitude does not by itself establish better or worse forecast skill. No confidence intervals are estimated for these deterministic comparisons.


Supplementary Fig. 11 | Longitudinal differences in observed and forecast height anomalies



a,c, Observed (solid) and forecast (dashed) DJF 200-hPa height anomalies along 45°N and 50°S. Dark grey denotes the equal-event composite of 1997–1998 and 2015–2016; red denotes 2023–2024. b,d, Forecast minus observed height along the same sections. All 360 longitude nodes per section are retained, starting at 120°E and continuing eastward around the globe. The two latitudes sample the northern and southern midlatitude circulation to illustrate differences in the longitude and strength of anomaly centres; they do not identify fixed Rossby-wave trajectories or the seasonally most active waveguide. Positive errors indicate heights predicted too high and negative errors heights predicted too low, relative to the observed field, irrespective of the sign of the observed anomaly.

Forecasts average the eight common leads. Anomalies and source fields are identical to Fig. 6. The focus is longitudinal error structure, not a new estimate of global MHW skill.


Supplementary Fig. 12 | Historical SST variance associated with climate modes and local processes



a, Allocation of historical regional SST variance among ENSO, other climate modes, prior ocean thermal state, surface heating, horizontal advection and an unresolved fraction. The denominator is total SST variance. b, The three local components divided by their combined share, so the denominator is the local-group allocation, not total variance. Prior thermal state comprises SST and mixed-layer temperature one month earlier. Other modes comprise IOD, PDO, NAO and SAM indices. Both panels use 432 monthly records per region during 1985–2020, with the ±23.5° masks of main Fig. 4. Seasonal means and a linear trend are removed before group-respecting variance allocation. Shared explanatory information is averaged over predictor-group orders, with nested ordering within the local group.

These are statistical SST associations, not heat-budget fractions or measured contributions of process-specific forecast errors in 2023–2024. Their connection to main Fig. 4b is the explicitly defined weighting rule in Supplementary Methods.


Supplementary Fig. 13 | Testing regional SST estimates in years excluded from fitting



This analysis uses contemporaneous observed climate indices, surface heating and horizontal advection, together with prior SST and mixed-layer temperature; it is diagnostic reconstruction, not an operational forecast. a, Later-period regional SST reconstruction using climate indices alone or all measured predictors. b, Skill decrease after removing each predictor group from the full model; negative values indicate improved reconstruction after removal. Three forward test blocks are 2003–2008, 2009–2014 and 2015–2020. Each model is trained only on earlier months beginning in 1985; seasonal terms, linear trend, scaling and coefficients are fitted within training data. Reconstruction skill is 1 − (model SSE / baseline SSE), where SSE is the sum of squared errors and the baseline is the training-fitted seasonal cycle plus linear trend. Squared errors are summed across test blocks before taking the ratio.

All seven regions and seven predictor configurations are retained. The three successive test blocks are not independent ENSO events. This score is not MHW AUC and does not validate the causal interpretation of the Sankey diagram.


Supplementary Fig. 14 | CFSv2-only circulation and surface heat-flux error comparison



a, CFSv2 circulation RMSE versus net surface heat-flux RMSE, each divided by its own 1997–1998/2015–2016 mean at the same region and lead. Colours and symbols distinguish the three events. The dashed line is the pooled fit. b, Percentage RMSE increases in 2023–2024 relative to those earlier-event means, averaged over leads 1, 3 and 6 months. Red denotes greater error and blue smaller error. c, Correlations between the two normalized errors after omitting each event; the red dashed line denotes the pooled estimate. All masks now use the ±23.5° divisions of main Figs. 3–4. The 54 remote pairs comprise three events × six recipient regions × three leads; nine additional tropical-Pacific pairs enter a.

This is a separately downloaded and processed CFSv2-only check, not an independent model: CFSv2 also contributes to the main SST ensemble. Circulation is departure from contemporaneous zonal mean, verified against ERA5; net heat flux is verified against ORAS5 and is positive into the ocean. Four members and three DJF months are equally weighted. The circulation metric differs from the anomaly correlation in main Fig. 3; no causal or independent-event significance is inferred.


Supplementary Fig. 15 | Regional and component accounting of lower and higher forecast skill



a, Area-weighted regional contributions to the net difference between the globally expected AUC (0.7050185) and the local neighbourhood AUC in 2023–2024, matching main Fig. 4. Red bars sum cells below that reference; blue hatched bars subtract cells above it; black diamonds show the net contribution. b, The same signed amounts allocated using historical SST-association weights. The three local components are shown separately and their parent group is not added again. Both panels divide by the same net map difference, 0.1065144 AUC. Positive shares sum to 120.79% and negative shares to −20.79%, giving 100% net. Regions use ±23.5° boundaries; Other seas comprises evaluated cells outside the seven masks.

The map uses April 2023–September 2024, nine leads and 11° same-basin neighbourhoods. Its area-weighted local-AUC difference is not the monthly spatial-AUC deficit in main Fig. 1 (0.1090497). Component weights describe historical SST associations, not causal contributions of process forecast errors. Unresolved is the unallocated association weight. Supplementary Methods gives the signed-sum identity and distinguishes the two AUC metrics.


Supplementary Fig. 16 | Regional skill changes and valid ocean-area coverage



a, Mean 1997–1998/2015–2016 neighbourhood AUC minus 2023–2024 AUC. c, 2023–2024 conditional intensity RMSE minus the two-event reference. Bars show region summaries; small points and whiskers retain all nine lead values and their minimum–maximum range. Positive differences indicate poorer forecasts in 2023–2024; negative differences indicate improvement. b,d, Cosine-latitude-weighted area coverage where all three events and all nine leads are evaluable for the respective metric. Denominators are the full original analysis-region ocean areas, not the already valid cells. TP, tropical Pacific; NP, North Pacific; SP, South Pacific; TA, tropical Atlantic; NA, North Atlantic; SA, South Atlantic; IO, Indian Ocean. These ±20° sensitivity-analysis boundaries differ from the ±23.5° masks in main Figs. 3–4 and Supplementary Fig. 15.

All spatial comparisons use April–September of the following year and 11° × 11° same-basin ocean neighbourhoods. AUC requires both observed classes; missing comparisons stay undefined. Neighbourhoods overlap and months/leads are dependent. Regional AUC differences are not additive contributions to the global monthly spatial-AUC gap. Lead ranges are descriptive, not confidence intervals.

Supplementary Fig. 17 | Sensitivity to verification duration and spatial pooling



Four combinations compare the original July–June window (12 months) and its symmetric three-month extension (April–September of the following year; 18 months), evaluated at individual grid cells or within fixed 11° × 11° neighbourhoods. a,c, Regional AUC decreases and conditional intensity RMSE increases. b,d, Their common valid ocean-area coverage. Red diamonds identify the 18-month, 11° configuration used in Supplementary Figs. 1c and 16. All three events use the same window, lead range and neighbourhood definition. The 12-month, single-cell calculation provides the unpooled reference. All seven fixed regions are retained; no region is removed because its sign differs. Individual-cell estimates use each grid cell separately; pooled estimates combine samples within an 11° × 11° same-basin neighbourhood.

Each point is a region-level summary, not an independent replicate or significance estimate. The longer window changes the seasonal/event-life-cycle support. The larger neighbourhood changes spatial support; higher coverage is not evidence of higher independent spatial resolution or a more accurate forecast. This sensitivity analysis retains the main archived input profile and its documented source-selection limitations.