

**1. D'Angelo 1.40.** Assume that  $f : \mathcal{S}^1 \rightarrow \mathbb{C}$  is  $k$  times continuously differentiable. Show that there is a constant  $C$  such that for  $n > 0$

$$|\widehat{f}(n)| \leq \frac{C}{n^k}.$$

**Solution:**

First, observe

$$\begin{aligned} \mathcal{F}(f^{(j)}) &= \int_0^{2\pi} f^{(j)} e^{-inx} dx \\ &= f^{(k-1)} e^{-inx} \Big|_{x=0}^{2\pi} + \frac{in}{2\pi} \int_0^{2\pi} f^{(j-1)} e^{-inx} dx \\ &= 0 + in \mathcal{F}(f^{(j-1)}). \end{aligned}$$

Proceeding inductively from  $k$ , we have

$$\mathcal{F}(f^{(j)}) = (in)^k \mathcal{F}(f).$$

Hence,

$$|\mathcal{F}(f)| = \frac{|\mathcal{F}(f^{(j)})|}{n^k} \leq \frac{\|f^{(j)}\|_{L_1}}{n^k}$$

which is finite since  $f^{(j)}$  is continuous on the compact set  $\mathcal{S}^1$ .

□

**2. D'Angelo 1.41.** Assume that  $f(x) = -1$  for  $-\pi < x < 0$  and  $f(x) = 1$  for  $0 < x < \pi$ . Compute the Fourier series for  $f$ .

**Solution:**

We calculate

$$\begin{aligned} \int_{-\pi}^{\pi} f(x) e^{-inx} dx &= \int_{-\pi}^{\pi} f(x) \cos(nx) dx - i \int_{-\pi}^{\pi} f(x) \sin(nx) dx \\ &= 0 - 2i \int_0^{\pi} \sin(nx) dx \\ &= \frac{-2i(\cos(n\pi) - 1)}{n} \\ &= \begin{cases} \frac{4i}{n} & \text{if } n \text{ is even} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

□

**3. D'Angelo 1.44.** Put  $D_k = \sum_{-k}^k e^{inx}$ . Define  $F_N$  by

$$F_N = \frac{D_0(x) + D_1(x) + \cdots + D_{N-1}(x)}{N},$$

and show that

$$F_N = \frac{1}{N} \frac{\sin^2\left(\frac{Nx}{2}\right)}{\sin^2\left(\frac{x}{2}\right)}.$$

**Solution:**

Denote  $\omega = e^{inx}$ , and note that  $\omega^{-1} = \bar{\omega}$ . Expanding  $F_N$ , we have

$$\begin{aligned} F_N(x) &= \frac{1}{N} \sum_{k=0}^{N-1} \sum_{n=-k}^k \omega^n = \frac{1}{N} \sum_{k=0}^{N-1} \omega^{-k} \sum_{n=-k}^k \omega^{k+n} \\ &= \frac{1}{N} \sum_{k=0}^{N-1} \omega^{-k} \sum_{n=0}^{2k} \omega^n \\ &= \frac{1}{N} \sum_{k=0}^{N-1} \omega^{-k} \frac{1 - \omega^{2k+1}}{1 - \omega} \\ &= \frac{1}{N} \frac{1}{1 - \omega} \sum_{k=0}^{N-1} \bar{\omega}^k - \omega^{k+1} \\ &= \frac{1}{N} \frac{1}{1 - \omega} \left( \frac{1 - \bar{\omega}^N}{1 - \bar{\omega}} - \frac{\omega - \omega^{N+1}}{1 - \omega} \right) \\ &= \frac{1}{N} \frac{1}{|1 - \omega|^2} \left( 1 - \bar{\omega}^N - (1 - \bar{\omega}) \frac{\omega - \omega^{N+1}}{1 - \omega} \right) \\ &= \frac{1}{N} \frac{1}{|1 - \omega|^2} \left( 1 - \bar{\omega}^N - \frac{-(1 - \omega) + \omega^N(1 - \omega)}{1 - \omega} \right) \\ &= \frac{1}{N} \frac{1}{|1 - \omega|^2} (2 - \bar{\omega}^N - \omega^N) \\ &= \frac{1}{N} \frac{2 - 2\cos(Nx)}{(1 - \cos x)^2 + \sin^2 x} \\ &= \frac{1}{N} \frac{2 - 2\cos(Nx)}{2 - 2\cos x}. \end{aligned}$$

The result follows upon using the following identity,

$$\frac{1 - \cos(\alpha)}{2} = \sin^2\left(\frac{\alpha}{2}\right).$$

□

**4. D'Angelo 1.49.** Derive the Weierstrass approximation theorem from Corollary 1.8.

**Solution:**

The Weierstrass approximation theorem states that for any continuous  $f : [a, b] \rightarrow \mathbb{C}$ , for any  $\varepsilon > 0$ , there exists a polynomial  $p$  so that

$$\|p - f\|_{\sup} < \varepsilon.$$

Without loss of generality, we can consider  $f$  on the circle by applying the continuous transformation

□

**5. D'Angelo 1.50.** If  $\{s_n\}$  is a monotone sequence of real numbers, show that the averages  $\sigma_N = \frac{1}{N} \sum_{j=1}^N s_j$  also define a monotone sequence. Give an example where the converse is false.

**Solution:**

Without loss of generality, assume that  $s_n \leq s_{n+1}$ . We must show that  $\sigma_N \leq \sigma_{N+1}$ . Observe

$$\begin{aligned} \sigma_{N+1} - \sigma_N &= \frac{s_1 + \cdots + s_{N+1}}{N+1} - \frac{s_1 + \cdots + s_N}{N} \\ &= \frac{N(s_1 + \cdots + s_{N+1}) - (N+1)(s_1 + \cdots + s_N)}{N(N+1)} \\ &= \frac{Ns_{N+1} - (s_1 + \cdots + s_N)}{N(N+1)} \\ &= \frac{(s_{N+1} - s_1) + (s_{N+1} - s_2) + \cdots + (s_{N+1} - s_N)}{N(N+1)} \\ &\geq 0 \end{aligned}$$

by monotonicity.

For the converse assertion, consider the sequence  $\{s_n\}$  given by  $s_n = \frac{(-1)^{n+1}}{n}$ . This sequence is clearly non-monotonic, yet

$$\sigma_n =$$

□