

1. Construct the Neumann series for the Fredholm equation of the second kind

$$y(x) = \lambda \int_0^1 y(s) ds + f(x)$$

for “small”  $\lambda$  and  $f(x) = 1$  and find the solution.

**Solution:**

Let  $A$  be the integral operator  $\int_0^1 \cdot ds$ . For small enough  $\lambda$ , the map  $y \rightarrow \lambda Ay + f$  is a contraction whose fixed point is the solution and is given by repeated composition to an arbitrary function  $y_0$ . Successive applications of this map to  $y_0 = 0$  yields the Neumann series

$$y_n = f + \lambda A f + \lambda^2 A^2 f + \cdots + \lambda^n A^n f. \quad (1)$$

Note that

$$A^n f = \int_0^1 \cdots \int_0^1 1 ds = 1. \quad (2)$$

So

$$y = \lim_{n \rightarrow \infty} y_n = \sum_{k=0}^{\infty} \lambda^k = \frac{1}{1 - \lambda}$$

for  $|\lambda| < 1$ .

We can easily verify that  $y$  is a solution,

$$y = \frac{1}{1 - \lambda} = \frac{\lambda + 1 - \lambda}{1 - \lambda} = \lambda \frac{1}{1 - \lambda} + 1 = \lambda Ay + f.$$

□

2. Construct the resolvent kernel for the above equation using various approaches:

(a) Assume that  $\lambda$  is “small”. For what  $\lambda$  does the series for the resolvent kernel converge?

**Solution:**

We take  $f \in C[0, 1]$ . Observe that  $\|Ay\| \leq \|y\|(1 - 0) = \|y\|$  and  $\|A1\| = 1$ , hence  $\|A\| = 1$ . Therefore, absolute and uniform convergence in (1) is guaranteed when  $\|A\||\lambda| = |\lambda| < 1$ . Taking the limit in (1), using the result that  $A^n$  is an integral operator with a given kernel  $k_n$ , and interchanging  $\int$  with  $\sum$ , the resolvent operator is given by

$$y = f + \lambda \int_0^1 \sum_{n=0}^{\infty} \lambda^n k_{n+1}(x, s) f(s) ds \quad (3)$$

Using similar reasoning as in (2), terms in the series of the resolvent kernel are given by

$$\lambda^n k_{n+1}(x, s) = \lambda^n \int_0^1 k_n(x, s) k_n(s, x) = \lambda^n \cdot 1,$$

Continuing from (3), the resolvent is

$$y = f + \frac{\lambda}{1 - \lambda} \int_0^1 f(s) ds. \quad (4)$$

□

(b) Use the fact that the kernel is symmetric and express the resolvent through the characteristic values and eigenfunctions.

**Solution:**

The characteristic values and eigenfunctions of  $A$  satisfy

$$\lambda \int_0^1 \phi_k(s) ds = \phi_k(x).$$

Hence all the eigenfunctions for  $A$  are constant functions, and the one orthonormal eigenfunction is  $\phi_1(x) = 1$  with characteristic value  $\lambda_1 = 1$ .

Following the procedure outlined in Lecture 8, for  $\lambda \neq \lambda_1 = 1$ , the unique solution to

$$y = \lambda Ay + f$$

is given by

$$y(x) = f(x) + \lambda \int_0^1 \sum_{k=1}^{\infty} \frac{\phi_k(s)\phi_k(x)}{\lambda_k - \lambda} f(s) ds = f(x) + \frac{\lambda}{1 - \lambda} \int_0^1 f(s) ds.$$

Note that this agrees with (3) and extends it to  $\lambda > 1$ .

If  $\lambda = 1$ , then there are two cases to consider. First, if  $f$  is orthogonal to 1, i.e.  $\int_0^1 f(s) ds = 0$ , then  $y = f$  since  $Af = 0$ . Note that these are not unique. If  $f$  is not orthogonal to 1, then there is no solution.

□

(c) Use the fact that the kernel is degenerate.

**Solution:**

Since the kernel is degenerate, we can reduce the problem by setting  $c = \int_0^1 y(s) ds$ , then integrating both sides of the problem. We get

$$c = \int_0^1 \lambda c + f(s) ds = \lambda c + f_1.$$

where  $f_1 = \int_0^1 f(s) ds$ . Solving this simple linear equation for  $c$  when  $\lambda \neq 1$ , we have  $c = (1 - \lambda)^{-1} f_1$  so

$$y = \lambda c + f = \frac{\lambda}{1 - \lambda} \int_0^1 f(s) ds + f. \quad (5)$$

Note that this agrees with the previous results.

When  $\lambda = 1$ , we have the condition that  $f_1 = 0$ . So there is no solution when  $f_1 \neq 0$ , and non-unique solutions  $y = c + f$  otherwise. This coincides with the symmetric kernel analysis.

□

**3. Analyze the equation**

$$y(x) = \lambda \int_{-1}^1 (2xs^3 + 5x^2s^2)y(s) ds + 7x^4 + 3$$

and solve it for any  $\lambda$ .

**Solution:**

Let  $A$  denote the integral operator above and  $f(x) = 7x^4 + 3$ . Note that the kernel is degenerate; i.e. it is of the form  $\sum_{i=1}^2 a_i(x)b_i(s)$  where  $a_1(x) = 2x$ ,  $b_1(s) = s^3$ ,  $a_2(x) = 5x^2$ , and  $b_2(s) = s^2$ . Hence, solutions are of the form

$$y(x) = \lambda a_1(x)(y, b_1) + \lambda a_2(x)(y, b_2) + f(x), \quad (6)$$

where  $(\cdot, \cdot)$  denotes the integral inner product.

Taking integral inner products on both sides of (6) by  $b_1$  and  $b_2$ , we have,

$$\begin{cases} (y, b_1) = \lambda(a_1, b_1)(y, b_1) + \lambda(a_2, b_1)(y, b_2) + (f, b_1) \\ (y, b_2) = \lambda(a_1, b_2)(y, b_1) + \lambda(a_2, b_2)(y, b_2) + (f, b_2) \end{cases} \quad (7)$$

if and only if

$$Y = \lambda KY + F,$$

where  $Y, K$  and  $F$  are the obvious vectors/matrices of inner products. The entries of  $K$  and  $F$  are

$$K = \begin{bmatrix} (a_1, b_1) & (a_2, b_1) \\ (a_1, b_2) & (a_2, b_2) \end{bmatrix} = \begin{bmatrix} \int_{-1}^1 2x^4 dx & \int_{-1}^1 5x^5 dx \\ \int_{-1}^1 2x^3 dx & \int_{-1}^1 5x^4 dx \end{bmatrix} = \begin{bmatrix} \frac{4}{5} & 0 \\ 0 & 2 \end{bmatrix},$$

and

$$F = \begin{bmatrix} (f, b_1) \\ (f, b_2) \end{bmatrix} = \begin{bmatrix} \int_{-1}^1 7s^7 + 3s^3 ds \\ \int_{-1}^1 7s^6 + 3s^2 ds \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \end{bmatrix}.$$

Thus, (7) reduces to

$$\begin{cases} (y, b_1) = \frac{4}{5}\lambda(y, b_1) \\ (y, b_2) = 2\lambda(y, b_2) + 4 \end{cases}. \quad (8)$$

When  $\lambda = \frac{5}{4}$ , the value of  $(y, b_1) = c$  is arbitrary and  $(y, b_2) = -\frac{8}{3}$ , and from (6) we can write the non-unique solutions as

$$y(x) = c\frac{5}{2}x - \frac{50}{3}x^2 + 7x^4 + 3.$$

When  $\lambda = \frac{1}{2}$ , then (8) gives rise the contradiction  $0 = 4$ , and there is no solution.

When  $\lambda \neq \frac{5}{4}$  and  $\lambda \neq \frac{1}{2}$ , the coefficients in  $Y$  are given by

$$Y = (I - \lambda K)^{-1}F = \begin{bmatrix} \frac{1}{1 - \frac{4}{5}\lambda} & 0 \\ 0 & \frac{1}{1 - 2\lambda} \end{bmatrix} \begin{bmatrix} 0 \\ 4 \end{bmatrix} = \begin{bmatrix} 0 \\ \frac{4}{1 - 2\lambda} \end{bmatrix}$$

and finally substituting into (6) gives

$$y(x) = \frac{4\lambda}{1 - 2\lambda} 5x^2 + 7x^4 + 3.$$

□