

Real Time Signal Processing with Symmetric and Asymmetric Support Intervals

Keven Joyce and Lia Harrington
University of Montana

May 14, 2015

Outline

Introduction and Motivation

Problem Outline

Problem Approach and Steps

Code Comments

Results and Conclusions

Importance of Real Time Signal Processing

What is real time signal processing?

- ▶ Applications

- ▶ Speech recognition
- ▶ Audio signal processing
- ▶ Video compression
- ▶ Weather forecasting
- ▶ Economic forecasting
- ▶ Medical imaging (e.g., CAT, MRI)
- ▶ And more...

Outline

Introduction and Motivation

Problem Outline

Problem Approach and Steps

Code Comments

Results and Conclusions

What is the problem?

Goal: We wish to reconstruct some generated signal \hat{x} that has been distorted by some error and convolution processes.

Solution: Take the convolution inverse of \hat{x} to reconstruct the signal.

Outline

Introduction and Motivation

Problem Outline

Problem Approach and Steps

Code Comments

Results and Conclusions

Problem Approach Overview

1. Specify problem settings

- ▶ $y = a * x + \nu, \quad \nu \in (0, \sigma^2 \delta).$
- ▶ $\hat{x} = r * y$
 - ▶ $r = P^{-1}q$
- ▶ Choose some $\Delta = [-d, d] \rightarrow$ initial estimate of x
- ▶ Choose some $\Delta = [T, \tau] \rightarrow$ initial estimate of x

2. Compute optimal Δ for r

- ▶ $H(\Delta) = E(\hat{x}_i - x_i)^2 = f_0 - \langle q, P^{-1}1q \rangle \Delta$
- ▶ Plot $H(\Delta)$ vs $d \rightarrow$ optimal $d \rightarrow$ optimal r

3. Simulation of measurement and processing

4. Illustrate the result of estimation

- ▶ $\sqrt{E(\hat{x}_i - x_i)^2} = \sqrt{H}$

Problem Strategy: Step 1

Specify the main ingredients of simulated measurement system:

- ▶ Specify point spread function (influence function) a ,
 - ▶ Symmetric
 - ▶ Asymmetric
- ▶ Covariance function ϕ for the signal x :





$$\phi = Cov(x) = b * b^* \quad (1)$$

- ▶ Variance σ^2 of the independent components of the additive random noise ν

$$y = a * x + \nu, \quad \nu \sim (0, \sigma^2 \delta). \quad (2)$$

- ▶ Choose support interval $\Delta = [-d, d]$ or $\Delta = [T, \tau]$

Signal and Covariance Setup



Outline

Introduction and Motivation

Problem Outline

Problem Approach and Steps

Code Comments

Results and Conclusions

Outline

Introduction and Motivation

Problem Outline

Problem Approach and Steps

Code Comments

Results and Conclusions

References

- [1] Golubtsov, P. (2015). Theoretical Big Data Analytics course notes.