

Force on a Spring = $k \cdot \text{Displacement}$

Work = Force $\cdot$ Distance

Hydrostatic Force = Density $\cdot$ Depth $\cdot$ Area

Mass = Density $\cdot$ Area

Moment = Distance $\cdot$ Mass

Center of Mass = $(\bar{x}, \bar{y}) = \left(\frac{M_y}{m}, \frac{M_x}{m}\right)$

1. A spring has a natural length of 5 ft. A force of 4 lb stretches this spring $1/4$ ft.
 - (a) Find the work done in stretching this spring from its natural length to 7 ft.
 - (b) find the work done in stretching this spring from 8 ft to 9 ft.
2. How much work is done in emptying a cylindrical tank of radius 5 m and height 10 m by pumping the the water through a pipe extending 4 m above the tank. (Hint: Water weighs 9800 N/m^3 , or 62.4 lb/ft^3 .)
3. A rectangular gate built into a dam is 4 ft high and 8 ft wide. The top of the gate is 2 ft below the surface of the water. What is the force on the gate?
4. What would be the force if the gate were the upper half of disc of radius 4 (and the top of the gate is 2 ft below the surface of the water)?
5. Find the mass, moments, and center of mass of a the triangular lamina bounded by the x - and y - axes and $y = 4 - 2x$. (As always, assume the density of this lamina to be 1.)

Copyright 2006 Dr. Brock Williams and Texas Tech University. Unauthorized reproduction prohibited.

1. If the ammount of mayhem is constant in an episode of the Muppet Show, there is a simple formula that gives the amount of mess created:

$$\text{Mess} = \text{Mayhem} * \text{Time}$$

That is, the mess is the product of the mayhem and the length of time it lasts. Typically, however, the mayhem varies over time. In a certain 30 minute episode the amount of mayhem was given by the function $M(t) = 4 \ln(t + 1)$, where $0 \leq t \leq 30$. Describe in complete detail how integration can be used to find the total mess in the episode.

2. It is a well-known fact that the anger generated by a fan during a football game is caused by both the frustration they experience and the length of time it occurs. The anger produced by a constant ammount of frustration is given by the formula

$$\text{Anger} = \frac{1}{3} (\text{Frustration})^2 * \text{Time}.$$

In the first quarter of a game, the Frustration was given by $F(t) = \sin\left(\frac{\pi}{30}t\right)$, where $0 \leq t \leq 15$.

Describe in complete detail how integration can be used to find the total Anger in the first quarter. Discuss the role played by Riemann sums, and set up an integral to find this Anger.

3. (a) What is the fundamental idea of Chapter 6?

(b) Suppose an object moves along the x -axis from $x = 2$ to $x = 5$ and is acted on by a force $F(x) = x^2$. Set up an integral to determine the amount of work done on the particle.

(c) Use the fundamental idea of Chapter 6 to explain in detail why the work on the particle is given by your integral.