Gravity from Quantum Measurement: From Microstates to Emergent Geometry

Kevin Player*

April 2, 2026

Abstract

We propose that gravity can be understood as the physical backreaction of quantum measurement. Standard results, horizon scales, entropy bounds, and the Bekenstein–Hawking area law, follow from Landauer's principle at the Unruh temperature as consequences of Newton's laws. To ground our idea, we extend a specific example from our previous work, a bilocal microstate construction where entangled photon pairs emitted from a localized source, propagate into opposite Rindler wedges. We show how polarisation measurements produce discrete energy-momentum impulses and correlated classical bits. Each such microstate is described geometrically at the horizon by an Aichelburg–Sexl pp-wave solution, whose longitudinal shift generates spatial separation between the wedges and whose transverse focusing increments the horizon area. The framework provides an operational realisation of ER=EPR at the level of these individual microstates, and complements the thermodynamic approaches of Jacobson and Verlinde by supplying a microphysical process underlying each entropy increment.

1 Introduction

Two landmark results establish that gravity and Unurh thermodynamics are deeply intertwined. Jacobson showed that the Einstein equations can be derived from thermodynamic identities applied to local Rindler horizons, with horizon entropy playing the role of the fundamental quantity from which spacetime dynamics emerge [1]. Verlinde extended this perspective, arguing that gravitational forces arise from entropy gradients on holographic screens, recasting gravity as an entropic rather than a fundamental interaction [2]. Both frameworks, however, operate at the level of bulk thermodynamics and leave open a prior question: what physical process underlies each individual entropy increment?

We argue that the answer is quantum measurement. Each act of information acquisition by a physical observer constitutes a discrete microstate event, and the collective backreaction of such events—together with the causal correlations they induce—gives rise to gravitational effects. The argument proceeds in three steps. First, any observation is mediated by a physical interaction: information transfer necessarily involves an exchange of energy and momentum between the observed system and the detector, a discrete impulse. Second, by the equivalence principle, this impulse is locally indistinguishable from the effect of a gravitational field: no local measurement can distinguish the recoil associated with a quantum measurement event from acceleration in a gravitational potential. Third, when such impulses are modelled geometrically—via Aichelburg–Sexl (AS) pp-wave solutions, as we explore—they imprint energy and information directly onto horizon

* *kjplaye@gmail.com*

geometry, and coarse-graining over the resulting ensemble recovers standard horizon thermodynamics. A unifying structure runs through the relations, and it is worth pausing to discuss the general ideas involved before diving into our explicit example.

1.1 Conceptual Roadmap and Subjective Relations

In each case in our argument, the relation between two adjacent concepts has the same logical character: every instance of the more abstract concept is grounded in an instance of the more physical one, but not every instance of the more physical concept gives rise to the more abstract one *for a given observer*. The gap in each case is filled by subjectivity. Quantum state reduction is always backed by an observation—it is a Bayesian update on the observer's part—but an observation that carries no new information for that observer produces no state reduction. Equally, every observation is backed by an energy-momentum transfer, but not every such transfer constitutes an observation for you: an interaction involving a different "observer" leaves your knowledge unchanged. The Unruh effect suggests the same structure extends to the equivalence principle: an accelerating observer perceives a thermal bath and attributes it to a gravitational field, while an inertial observer sees only vacuum. Whether an acceleration counts as gravity is perspectival in precisely the same sense. In each case the relation is an identification, not a causal link, and the identification is observer-relative.

This stands in sharp contrast to the Diósi–Penrose objective collapse programme [3, 4], which postulates that gravity *causes* quantum state reduction as an objective, observer-independent physical process. Figure 1 summarises this contrast. The three solid arrows represent the perspectival grounding relations described above; the single dashed arrow represents the Diósi–Penrose causal postulate. It is the only relation in the diagram claimed to be objective and observer-independent, and it is the only one we do not adopt.

$$\begin{tikzcd}
\text{Gravitational field} \arrow[r, dashed, "\text{Diósi–Penrose}"] & \text{Quantum state reduction} \\
\text{Acceleration} \arrow[u, "\text{equivalence principle}"] \arrow[r, "\text{energy–momentum transfer}"'] & \text{Observation} \arrow[u, "\text{Bayesian update}"']
\end{tikzcd}$$

Figure 1: Solid arrows denote "substrate" relations: $A \to B$ means every instance of B is physically realised by an instance of A, for a given observer. The dashed arrow denotes the Diósi–Penrose postulate, where the relation is claimed to hold objectively rather than observer-relatively.

[Section map paragraph todo]

2 Motivating Heuristics

Landauer's principle associates a minimal energy cost with the acquisition of information [5]. For an accelerated observer, the relevant temperature is the Unruh temperature [6]. Substituting one into the other eliminates temperature entirely, yielding a relation between information acquisition, energy transfer, and acceleration (a version of the BHT first law, equation (3)). This is not merely an algebraic convenience: it is a concrete dictionary between the thermal description natural to a Rindler observer and the non-thermal global Minkowski description, with the apparent thermality of the vacuum emerging as a

consequence of the observer's acceleration rather than as a fundamental property of the state. In this way we derive a sequence of standard results in black hole thermodynamics as a consequence of Newton's laws.

2.1 The Unruh-Landauer Relation

Landauer's principle suggests that the acquisition of ΔS bits of information by a physical observer requires a minimum energy transfer

$$\Delta E \geq k_B T \, \Delta S. \tag{1}$$

We note that Landauer's principle is conventionally stated for information erasure rather than acquisition. Here we apply it to acquisition, which is the physically relevant process for an observer registering a measurement outcome. The bound has the same form in both cases; we apply it to acquisition on the grounds that registering a measurement outcome is physically equivalent to a bit being written into the detector's state, which is the process Landauer's principle governs.

For an observer undergoing proper acceleration a, the Unruh effect associates a temperature

$$T = \frac{\hbar a}{2\pi k_B c} \tag{2}$$

with the local vacuum. Substituting into Landauer's bound eliminates $k_B T$, yielding

$$\boxed{\Delta E \geq \frac{\hbar a}{2\pi c} \, \Delta S.} \tag{3}$$

Temperature has dropped out entirely: equation (3) is a temperature-independent relation between energy, acceleration, and information, suitable for the Minkowski vacuum. In what follows, standard results emerge as consequences of this single relation and Newton's laws.

2.2 Momentum Transfer and Acceleration

The energy transfer ΔE implied by equation (3) is carried by a physical quantum — matter or radiation — mediating the measurement interaction. We refer to this as the *thrust quantum* of effective mass m_t. (This will actually be a pure energy carrier our later microstate example.) For a relativistic carrier, the associated momentum transfer is

$$\Delta p = \frac{\Delta E}{c}. \tag{4}$$

We take the interaction time to be the order of the light-crossing time across the interaction region, $\Delta t \sim 2r/c$, corresponding to a causal exchange over the scale r. The numerical factor of 2 is not fixed a priori at this level; it will be determined by requiring consistency with the gravitational scale that emerges below. For a detector of mass m_d, Newton's second law in impulse form gives

$$a = \frac{\Delta p}{m_d \Delta t} = \frac{\Delta E}{2 m_d r}. \tag{5}$$

2.3 The Schwarzschild Radius

This step should be understood as a calibration: we match the acceleration inferred from the measurement-induced impulse to the Newtonian gravitational field in order to identify the characteristic length scale associated with the detector. The thrust quantum carries

energy ΔE and produces an acceleration a on the detector. Identifying this with the Newtonian gravitational acceleration due to the mass m_t at distance r.

$$a = \frac{Gm_t}{r^2}, \tag{6}$$

and substituting equation (5) with $\Delta E = m_t c^2$, we obtain

$$\frac{m_t c^2}{2m_d r} = \frac{Gm_t}{r^2}. \tag{7}$$

Solving for r:

$$r = \frac{2Gm_d}{c^2} \equiv r_s, \tag{8}$$

which is precisely the Schwarzschild radius of the detector mass m_d. This fixes the factor of 2 in the interaction time retroactively: the choice $\Delta r = 2r/c$ is the unique one consistent with Newtonian gravity and the correct Schwarzschild scale.

It is notable that the resulting length scale is set by the detector mass m_d, rather than the source mass m_t. This reflects the fact that the construction is inherently observer-centred: the acceleration entering the argument is the response of the detector to the measurement-induced impulse, and the associated scale is therefore the horizon radius relevant to that observer. In this sense, the quantity r_s should be interpreted not as the Schwarzschild radius of the source, but as the characteristic horizon scale associated with the detector as a physical system undergoing measurement. This is consistent with the observer-dependence of horizon structure in relativistic settings, as exemplified by the Unruh effect, where acceleration alone gives rise to a causal horizon.

2.4 Surface Gravity

Evaluating the acceleration at $r = r_s$ gives the surface gravity of the detector:

$$\kappa = \frac{Gm_d}{r_s^2} = \frac{c^4}{4Gm_d}, \tag{9}$$

in agreement with the standard Schwarzschild result. This is the acceleration that appears in the Unruh temperature associated with the detector's horizon, $T = \hbar\kappa/2\pi k_B c$, providing an internal consistency check on the framework.

2.5 Bekenstein-Hawking Entropy

Returning to the Unruh-Landauer relation (3) and substituting the surface gravity κ for a, the entropy increment associated with an energy increment $\Delta E = c^2 \Delta m_d$ is

$$\Delta S \leq \frac{2\pi c}{\hbar\kappa}\Delta E = \frac{8\pi G m_d}{\hbar c^3} \cdot c^2 \Delta m_d. \tag{10}$$

Expressing this in terms of the change in Schwarzschild area $\Delta A = 32\pi G^2 m_d \Delta m_d / c^4$:

$$\Delta S \leq \frac{\Delta A}{4\ell_P^2}, \qquad \ell_P^2 = \frac{\hbar G}{c^3}, \tag{11}$$

which is the Bekenstein-Hawking area law. Each measurement event contributes a discrete entropy increment proportional to the area imprinted on the horizon by the associated energy-momentum transfer.

The two bounds are saturated simultaneously when $a = c^2/R$ — which is exactly the condition for being *at a horizon*. The horizon is therefore not just where thermodynamics happens to work out, it is the unique locus where the minimum cost of information acquisition equals the maximum information content of the region.

2.6 Summary

Beginning from the single relation (3), and using only Newton's laws and the equivalence principle, we have recovered the Schwarzschild radius, the surface gravity, and the Bekenstein-Hawking area law as consequences of the observation $\rightarrow$ acceleration identification. These are standard results, but their derivation here from the physical cost of information acquisition rather than from thermodynamic postulates is the point.

The thrust quantum introduced in this section, the physical carrier of the measurement-induced impulse, will be given a explicit microstate realization in the sections that follow. It is identified with one of a pair of entangled photons modeled as Aichelburg-Sexl pp-waves crossing past Rindler horizons in the Unruh picture. In this case the back reaction of a single force carrier in one wedge is correlated with an equal and opposite reaction in the other from a common source.

3 Bilocal Microstate Construction

Let $\hbar = c = 1$.

The heuristic arguments of Section 2 establish a relationship between information acquisition, acceleration, and gravitational effects, but leave open the question of what a single microstate looks like concretely. In standard treatments, the Unruh effect is derived by coupling a detector to a pre-existing acceleration and reading off a thermal response from the vacuum. Following [7], we invert this logic: we place an interaction in the past, where a source injects non-thermal entangled excitations into the field, and show that the observer's accelerated response is driven by these source-induced excitations. The thermal ensemble then emerges as a coarse-grained aggregate of such microstates, rather than being a primitive property of the vacuum.

We now upgrade this construction from a spin-0 scalar field in the original source [7] to the spin-1 electromagnetic field. This is not merely a technical generalisation. As we show, the photon carries a unit of action $\hbar$ per cycle alongside its energy $E = \hbar\omega$, which closes the Landauer gap left open in Section 2: the minimum actionable event in the construction is precisely a single photon, and its polarisation state constitutes the classical bit acquired by each observer. The anticorrelated helicities of the entangled pair are the natural completion of the $\mathbb{Z}_2$ swap symmetry between wedges, and as we discuss in Section 5, they imprint a definite geometric signature onto the horizon consistent with the ER=EPR identification.

3.1 The Paz–Lester–Wright Setup

We retain the Alice-and-Bob style naming convention. Three observers are placed at spacetime positions $(t, z) = (0, 1)$, $(0, -1)$, and $(-1, 0)$, named *Wright* (R), *Lester* (L), and *Paz* (P) respectively — a mnemonic for Right, Left, and Past. Paz serves as the common causal ancestor of Wright and Lester (see Figure 2).

Paz emits an entangled pair of spin-1 massless photons with momenta $\pm k$ and Rindler frequency $\omega = \omega_k$. The momenta are perfectly balanced, so the source exhibits no net back-reaction in this minimal model. The two photons propagate into the left and right Rindler wedges, where Lester and Wright detect them respectively, each registering a helicity eigenvalue and acquiring one classical bit of information.

3.2 Double-Null Gauge

The emission event naturally defines two null directions: the right-moving photon has wave-vector k_R^μ and the left-moving photon has wave-vector $k_L^\mu = -k_R^\mu$. These two null



Figure 2: A source in the past, Paz (P), emits an entangled photon pair that accelerates Wright (R) into the right Rindler wedge and Lester (L) into the left Rindler wedge. The two null directions of propagation define the x^+ and x^- axes of the double-null coordinate system.

directions are precisely the x^+ and x^- axes of double-null coordinates

$$x^{\pm} = \frac{1}{\sqrt{2}}(t \pm z), \tag{12}$$

with transverse coordinates $\mathbf{x}_\perp = (x^1, x^2)$. Working in this frame we impose the symmetric double-null gauge condition

$$A_+(x) = 0, \qquad A_-(x) = 0, \tag{13}$$

so that the gauge field is supported entirely on the transverse components $\mathbf{A}_\perp = (A^1, A^2)$. This gauge respects the $\mathbb{Z}_2$ symmetry that swaps the two wedges ($x^+ \leftrightarrow x^-$) simultaneously with ($R \leftrightarrow L$), and therefore treats both observers symmetrically by construction. The residual gauge freedom is fixed by requiring $\partial_i A^i = 0$ in the transverse plane, leaving precisely the two physical helicity degrees of freedom $\epsilon^i_{(\lambda)}$ with $\lambda = \pm 1$.

The lack of manifest four-dimensional covariance in equation (13) is a consequence of aligning the gauge with the pair of physical null directions selected by the source, rather than an arbitrary choice. The full Lorentz group is broken to the little group preserving the pair $\{k^\mu_R, k^\mu_L\}$, which is the two-dimensional Euclidean group $ISO(2)$ acting on the transverse plane. This is precisely the little group of massless particles, and the two helicity eigenstates $\lambda = \pm 1$ are its irreducible representations. All physical results are independent of the gauge choice.

3.3 The Bilocal Source

In double-null gauge the electromagnetic field is characterised by its transverse components alone. The bilocal driving source generalises the scalar construction to

$$J^{ij}(x, y) = \lambda\, \epsilon^{i*}_{(-1)}\, \epsilon^j_{(+1)}\, f_L(x)\, f_R(y), \tag{14}$$

where λ is a coupling constant, f_L and f_R are normalised mode profiles supported on the left and right wedges respectively, and $\epsilon^i_{(\lambda)}$ are the circular polarisation vectors

$$\epsilon^i_{(+1)} = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\ i\end{pmatrix}, \qquad \epsilon^i_{(-1)} = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\ -i\end{pmatrix}. \tag{15}$$

The index structure of (14) assigns helicity $+1$ to the right-moving mode and helicity -1 to the left-moving mode: the pair is *anticorrelated* in helicity. This is the unique

assignment consistent with angular momentum conservation at the emission vertex and with the $\mathbb{Z}_2$ wedge-swap symmetry, under which $\epsilon^i_{(+1)} \leftrightarrow \epsilon^i_{(-1)}$.

The sourced Lagrangian reads

$$\begin{aligned}\mathcal{L}_{\text{sourced}} &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \int J^{ij}(x,y)\, A^+_i(x)\, A^+_j(y)\, d^4x\, d^4y \;+\; \text{h.c.} \\ &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \Big(a^{L\dagger}_{-k,(-1)}\, a^{R\dagger}_{k,(+1)} \;+\; a^{L}_{-k,(-1)}\, a^{R}_{k,(+1)} \Big),\end{aligned} \tag{16}$$

where $a^{L\dagger}_{-k,(-1)}$ and $a^{R\dagger}_{k,(+1)}$ are creation operators for left-wedge helicity -1 and right-wedge helicity $+1$ modes respectively. The interaction term in (16) is a two-mode squeezing interaction at frequency ω, now resolved by helicity: it is the direct vector-field analogue of the scalar squeezing interaction [8], with the anticorrelated polarisation structure made explicit. The photon pair carries total angular momentum zero, consistent with emission from a scalar source at Paz.

Each photon carries energy $E = \hbar\omega$ and, as a mode of the electromagnetic field, constitutes precisely one quantum of action $\hbar$ per oscillation cycle. The thrust quantum of Section 2 is thereby identified with a single photon: the minimum actionable event is the emission and detection of one such quantum. The energy transferred per bit acquired is $\Delta E = \hbar\omega$, saturating the Unruh–Landauer bound (3) at the fundamental quantum level. This closes the Landauer gap: registering a photon's helicity writes one classical bit into the detector's physical state, which is precisely the process Landauer's principle governs.

3.4 Polarisation Bogoliubov Transformation

We now show that the polarisation labels survive the Rindler–Unruh mode decomposition, so that the thermal ensemble retains its helicity structure and the anticorrelation of the emitted pair descends unambiguously to the level of individual Rindler microstates.

In the transverse plane, the electromagnetic field in double-null gauge decomposes into two independent scalar-like sectors labelled by helicity $\lambda = \pm 1$. For each sector, the right-wedge field expands as

$$A^i_\lambda(x) = \int_0^\infty \frac{d\Omega}{2\pi} \frac{1}{\sqrt{2\Omega}}\, \epsilon^i_{(\lambda)} \left[b^R_{\Omega,\lambda}\, u^R_\Omega(x) + b^{R\dagger}_{\Omega,\lambda}\, u^{R*}_\Omega(x) \right], \tag{17}$$

where $u^R_\Omega(x)$ are the right-wedge Rindler modes at Rindler frequency Ω, and $b^R_{\Omega,\lambda}$, $b^{R\dagger}_{\Omega,\lambda}$ are the corresponding annihilation and creation operators satisfying

$$\left[b^R_{\Omega,\lambda},\, b^{R\dagger}_{\Omega',\lambda'} \right] = \delta(\Omega - \Omega')\, \delta_{\lambda\lambda'}. \tag{18}$$

An identical expansion holds in the left wedge with $R \to L$ and $\lambda \to -\lambda$, reflecting the anticorrelated helicity assignment. Because each helicity sector is independently a scalar-like sector, the Bogoliubov transformation relating Minkowski and Rindler modes takes the same form in each sector. For helicity λ at Rindler frequency Ω, the Unruh modes are

$$c_{\Omega,\lambda} = \cosh\theta_\Omega\, b^R_{\Omega,\lambda} - \sinh\theta_\Omega\, b^{L\dagger}_{\Omega,-\lambda}, \tag{19}$$

where $\tanh\theta_\Omega = e^{-\pi\Omega c/a}$ encodes the Unruh distribution at acceleration a, and $-\lambda$ on the left-wedge operator reflects the anticorrelation. The transformation (19) is the standard Bogoliubov relation for a massless field [6, 9], now carrying a helicity label throughout. The Minkowski vacuum in the Rindler basis reads, for each helicity sector,

$$|0\rangle^{(\lambda)}_M = \frac{1}{\cosh\theta_\Omega} \sum_{n=0}^{\infty} \tanh^n\theta_\Omega\, |n_{\Omega,\lambda}\rangle_R\, |n_{\Omega,-\lambda}\rangle_L, \tag{20}$$

and the full vacuum is the product over all Ω and both helicities:

$$|0\rangle_M = \prod_{\Omega,\lambda} |0\rangle_M^{(\lambda)}. \tag{21}$$

Equation (20) makes explicit that the Minkowski vacuum is a sum over entangled helicity-anticorrelated pairs at each Rindler frequency. The bilocal source selects a single term from this sum at fixed Ω and fixed helicity pair $(\lambda, -\lambda) = (+1, -1)$, yielding the single-microstate

$$|\Psi_\Omega\rangle_M = |1_{\Omega,+1}\rangle_R \, |1_{\Omega,-1}\rangle_L + \mathcal{O}(\text{higher order}). \tag{22}$$

Wright's detection of helicity $+1$ and Lester's detection of helicity -1 are therefore perfectly anticorrelated: each measurement event yields one classical bit, and the two bits are the complement of one another.

The key structural result is that the helicity label λ is conserved under the Bogoliubov transformation: it enters equation (19) as a spectator index and exits unchanged on the Unruh operators $c_{\Omega,\lambda}$. Thermality, when it arises from coarse-graining over the ensemble of microstates (22), is therefore helicity-resolved: each helicity sector thermalises independently, and the anticorrelation between wedges is preserved at every level of the construction.

3.5 Summary

The passage from the scalar to the vector field construction introduces three structural improvements. First, the photon carrying $E = \hbar\omega$ and one quantum of action per cycle identifies the thrust quantum of Section 2 with a concrete physical object, closing the Landauer acquisition gap. Second, the double-null gauge condition, motivated by the pair of null directions defined by the emission event, treats both Rindler wedges symmetrically and exposes the $\mathbb{Z}_2$ swap symmetry as the organising principle of the construction. Third, the anticorrelated helicity structure, conserved through the Bogoliubov transformation, provides each observer with one classical bit whose values are perfectly complementary: the two bits together constitute the correlated classical record that, in the AS pp-wave picture of Section 5, is imprinted as geometric information onto the horizon.

4 Matter as Substrate: Ticks, Memory, and the Thrust Quantum

We show how information is encoded in matter and energy and is reflected in the horizon geometry. This requires a physical account of what it means for matter to *carry* information.

A massive particle accumulates phase along its worldline at a rate set by its rest energy. The Compton frequency relation $f = E/h$ assigns a natural frequency to any mass m, and Bremermann's computational limit [10] independently recovers the same rate $f = mc^2/h$ as the maximum clock rate, or Nyquist rate, of any physical system with that energy E. These two results agree because they are both expressions of the same underlying fact: a quantum of action $\hbar$ is the minimal unit of physical evolution, and mc^2/h is the rate at which a mass m consumes such quanta.

Each such quantum corresponds to one tick of the phase $e^{iS/\hbar}$ in the path integral, and classical trajectories emerge from the constructive interference of these ticks across paths. Paths separated by less than one tick of action — less than $\hbar$ — contribute indistinguishably to the interference, so the trajectory is resolved only down to the scale set by $\hbar/E$: this is precisely the Heisenberg uncertainty at energy E, and coarser descriptions of the trajectory discard the sub-tick phase information that finer energy scales would

reveal. This agrees notionally with the old quantum physics of Bohr and Sommerfeld and only picks up finer phase information at higher effective energy scales.

This gives matter a precise sense in which it is a *substrate for information*: it accumulates ticks, and each tick is a discrete unit of action, Lorentz-invariant and recorded in the particle's proper time. Matter is persistent memory in the operational sense that its history of ticks is physically encoded in its state and cannot be erased without an energy cost governed by Landauer's principle. The maximum rate of this memory is bounded by Bremermann.x

A massless particle — the photon — has no rest frame, no proper time, and accumulates no action between emission and absorption. It does not tick. It therefore carries no persistent memory and cannot serve as a substrate. Its role in the present construction is precisely that of a pure information carrier. A single photon carries exactly two irreducible constituents, distinct in kind:

- **One quantum of action** $\hbar$: the physical transfer, deposited as energy $\hbar\omega$ into the detector, incrementing its Bremermann tick rate by $\Delta f = \hbar\omega/h$ for as long as the energy remains in the mass. This increment is the physical trace of the interaction in the detector's matter-memory.
- **One bit of polarisation**: the informational outcome, the classical record of which helicity — $\lambda = +1$ or $\lambda = -1$ — was absorbed. This bit persists in the detector's state for as long as the detector retains the record. Erasure of this bit carries a minimum energy cost governed by Landauer's principle, regardless of how the record was originally established. energy cost.

These are not the same thing in different guises. The $\hbar$ is what makes the event physical; the bit is what makes it meaningful to an observer. That a single photon carries exactly one of each is not a coincidence: it is why the photon is the natural minimal unit of the entire construction, the unique object that saturates both the action and the information count simultaneously. The energy deposited is $\Delta E = \hbar\omega$, saturating the Unruh–Landauer bound at the fundamental quantum level, as established in Section 3.

The horizon enters this picture not as a storage medium but as a boundary condition. It marks the surface at which the matter-memory on one side is causally disconnected from the matter-memory on the other. Horizon entropy, in this account, is not a property of the horizon surface itself but a count of the ticks that have been written into the matter on each side — bounded by Bremermann from above and by Landauer from below. The holographic appearance of this count, proportional to area rather than volume, is a consequence of the Unruh–Landauer and Bekenstein relations derived in Section 2 and does not require holography as an independent postulate.

This reframing has a direct consequence for the ER=EPR identification developed in the next subsection. The correlated measurement outcomes of Wright and Lester are not encoded in the horizon geometry in the first instance — they are encoded in the matter of their respective detectors, each of which has absorbed one photon and accumulated one additional tick. The geometric correlations between the two wedges, realised as correlated AS pp-wave imprints in Section 5, are downstream of this matter-level encoding. The bridge, if it exists, is built from correlated ticks, not from the horizon itself.

4.1 Entanglement as Spatial Information: ER=EPR at the Microstate Level

Each measurement produces a polarisation outcome — H or V — at each wedge, perfectly anticorrelated between Lester and Wright. But consider a different question: not *what* was observed, but *where*. The question "who observed H?" has a binary answer: Lester (left wedge) or Wright (right wedge). This is a *spatial bit* — a bit that encodes not a polarisation state but a location in space.

Each measurement event has, as established in Section 4, exactly two irreducible constituents: one quantum of action $\hbar$ written irreversibly into the detector's matter-memory as a tick, and one bit of polarisation registered as a classical outcome. The entanglement between Wright and Lester is consumed in producing these two records — one tick and one bit at each wedge. What remains after measurement is not a remnant of the entanglement but its classical residue: the perfect anticorrelation between the two bits, and the correlated pair of ticks written into the matter on each side of the horizon. The spatial bit is precisely the polarisation bit of Section 4 now read as a location rather than a spin state: the same bit that records *what* was observed also records *where*, because in this construction the two wedges carry opposite helicities by design.

The information is carried by the matter of the detectors, not by the horizon. As established in Section 4, the horizon is a boundary that makes the left/right distinction sharp; it is the matter on each side that accumulates the ticks and holds the classical record. Wright's detector has absorbed one photon, gained one tick, and registered helicity $+1$. Lester's detector has absorbed one photon, gained one tick, and registered helicity -1. The two records are perfectly complementary by the anticorrelated structure of the bilocal source, and their complementarity is encoded in matter before it is reflected in geometry.

This reframing sharpens the ER=EPR identification. The two wedges are connected not by the horizon as an abstract geometric object but by the physical correlations between the energy-momentum impulses deposited in the matter on each side. In the AS pp-wave picture developed in Section 5, each such impulse produces a measurable longitudinal shift and transverse focusing of the horizon geometry. The correlated pair of impulses — one in each wedge, anticorrelated in helicity — produces correlated geometric deformations on the two sides of the horizon, and it is *this* geometric correlation, downstream of the matter-level encoding, that constitutes the Einstein–Rosen bridge at the level of a single microstate.

The spatial bit — the answer to "who observed H?" — is therefore not merely a record of which side the photon landed on. It is a label for which of the two correlated geometric deformations was realised, and the two labels together reconstruct the bridge. This is an operational realisation of ER=EPR at the level of individual microstates: the entanglement between Lester and Wright's outcomes is not a stand-in for the bridge but is physically instantiated in the pair of correlated ticks whose geometric imprint *is* the bridge. The bridge is built from matter, tick by tick.

5 Gravitational Imprint: the pp-Wave Realisation

Each measurement event in the bilocal construction involves a photon transferring energy and momentum to an accelerated detector. We now model the gravitational imprint of this transfer by identifying each photon with an Aichelburg–Sexl (AS) pp-wave solution in general relativity [11]. The AS geometry is an exact solution describing the gravitational field of an ultrarelativistic point mass, and provides the natural geometric language for a single-photon microstate.

5.1 The AS pp-Wave and Its Effects

Let $u = t - z$ and $v = t + z$, where the photon propagates along the null plane $u = 0$, z is the longitudinal direction, and (x, y) are transverse coordinates. Test particles at fixed transverse positions (x, y) experience two effects as the shock front crosses $u = 0$:

1. **Longitudinal shift.** The v coordinate is displaced by

$$\Delta v \propto -\ln r^2, \qquad r^2 = x^2 + y^2. \tag{23}$$

This shift is lightlike: no proper time elapses for a particle crossing the shock, but it is displaced along the null direction. Near the axis ($r \to 0$) the shift diverges, encoding the concentration of energy along the photon's trajectory.

2. **Transverse focusing.** The gradient of the longitudinal shift produces a transverse velocity

$$\Delta \mathbf{v}_\perp \propto \nabla \ln r \propto \frac{1}{r}, \tag{24}$$

directed toward the axis. Test particles are pulled toward the centre of the wave, regardless of their initial transverse separation.

5.2 The Longitudinal Shift Makes Space

The longitudinal shift deserves closer attention, because it carries a meaning beyond a mere coordinate displacement. The photon emitted by Paz propagates along the null direction; as it crosses the Rindler horizon, the AS shift displaces the future light cone of every test particle it passes. For the photon traveling into the right wedge, the shift pushes the right wedge's geometry in the $+v$ direction; for the entangled partner traveling into the left wedge, the shift pushes the left wedge in the $-v$ direction. The two wedges are displaced *away from each other* along the longitudinal null direction.

This is not merely a kinematic effect. The transverse plane of the pp-wave is identified with horizon area elements $dA = dx\,dy$, and the longitudinal shift creates a new null interval between the two wedges — a region of spacetime that did not exist, in the sense of being causally accessible, before the photon passed. The bit of spatial information encoded in the left/right measurement outcome — "who observed H?" — refers to precisely this separation. The measurement event does not merely *record* a spatial distinction; through the gravitational backreaction of the photon, it *creates* the spatial separation that the bit describes.

Put differently: the same physical event — a photon crossing a horizon — simultaneously encodes a classical bit and, via the AS longitudinal shift, generates the spatial structure that bit refers to. Space is not a pre-existing backdrop onto which information is written; it is produced, increment by increment, by the gravitational imprint of each measurement event. This is the microstate-level content of emergent space: each detection event contributes both an entropy increment and a spatial increment, and the two are the same thing described in different languages.

5.3 Horizon Entropy from Individual Microstates

Each detection event contributes a discrete entropy increment via the Bekenstein–Hawking relation derived in Section 2:

$$\Delta S = \frac{\Delta A}{4\ell_P^2}, \qquad \ell_P^2 = \frac{\hbar G}{c^3}, \tag{25}$$

where ΔA is the area element imprinted on the horizon by the transverse focusing of the pp-wave. Horizon entropy is therefore not a bulk thermodynamic quantity assigned to the horizon as a whole, but a sum of discrete, calculable contributions from individual measurement-induced interactions. The transverse focusing and the longitudinal shift are two aspects of the same microstate: focusing encodes the entropy increment in the area; the longitudinal shift encodes the spatial increment in the geometry.

5.4 Coarse-Graining and the Emergence of Thermality

In a steady-state scenario, Paz continuously emits entangled pairs, and Lester and Wright continuously detect them. Each detection is an AS pp-wave microstate with a definite,

non-thermal description in the global Minkowski frame: a specific photon energy, a specific transverse position, a specific longitudinal shift. Coarse-graining over this ensemble recovers the Unruh thermal spectrum at temperature $T = a/2\pi$, consistent with the vacuum decomposition of equation (**??**). The thermodynamic description of the horizon emerges from this averaging; it is not a feature of the individual microstates.

This completes the microstate programme. Each entropy increment in the Bekenstein–Hawking area law corresponds to a single AS pp-wave interaction. Each pp-wave interaction corresponds to a single measurement event in the bilocal construction. Each measurement event both encodes a spatial bit and generates the spatial separation that bit refers to. The gravitational, thermodynamic, information-theoretic, and geometric descriptions of the horizon are unified at the level of individual microstates, and thermality, entropy, and space itself emerge together from the coarse-grained aggregate of these events.

6 Connecting Microstates to Emergent Geometry

The preceding sections established three key loci where the geometry responds to information-processing events:

1. **Paz in the past:** the bilocal source injects energy and angular momentum, which can be associated with a Schwarzschild-like mass m_P encoding the initial acceleration of the detectors.
2. **Photon crossing the horizon:** each emitted photon carries energy $E = \hbar\omega$ and a helicity bit, producing an Aichelburg–Sexl pp-wave whose longitudinal shift generates a new null separation between wedges and whose transverse focusing increments the horizon area.
3. **Wright/Lester detectors:** absorption of the photon imprints energy and one classical bit into the matter-memory of each detector, producing local Schwarzschild-like responses m_W, m_L and marking the accumulation of *ticks*.

Each of these loci has its own emergent time, defined operationally by the sequence of discrete *ticks* accumulated in the proper time of matter carrying information. For the photon itself, no proper time elapses; it carries a *pure transfer* of energy and one classical bit. The photon's passage between emission and detection mediates correlations that are encoded as entanglement between the ticks of Wright and Lester, and as longitudinal shifts of the horizon in the pp-wave picture.

The Raychaudhuri equation applied to each infinitesimal horizon element ensures that the transverse focusing $\Delta\theta_\perp$ matches the area increment ΔA associated with a single microstate. This realizes the Bekenstein–Hawking relation at the level of individual detection events, making Jacobson's thermodynamic argument manifest at the microstate scale.

The longitudinal shifts, meanwhile, generate the *spatial separation* that the entanglement between detectors refers to: space is created incrementally as a direct consequence of energy-momentum transfer and measurement. In this sense, ER=EPR is instantiated physically: the Einstein–Rosen bridge between the left and right wedges is not postulated, but emerges as the cumulative geometric imprint of paired ticks and photons, each producing correlated pp-wave distortions.

The sequence

$$\text{Paz source} \xrightarrow{\text{photon emission}} \text{pp-wave horizon shift} \xrightarrow{\text{detector absorption}} \text{matter ticks} \tag{26}$$

provides a concrete operational dictionary linking energy, information, and geometry. Each step is reversible in principle except for the Landauer cost encoded in the matter-

memory, and coarse-graining over many such events recovers the familiar thermal horizon description.

In this framework, *space, time, and entropy emerge simultaneously from discrete microphysical events.* Proper time flows in the detectors via tick accumulation, spatial separation arises from longitudinal pp-wave shifts, and horizon area encodes entropy through transverse focusing. Each photon acts as a bridge between these emergent structures, unifying the information-theoretic, gravitational, and thermodynamic perspectives into a single microstate-level picture.

6.1 Horizon Area Change from a Single AS Photon

The connection between energy flux across a null surface and horizon area change follows directly from Jacobson's derivation of the Einstein equations from horizon thermodynamics [1]. In his framework, for a null congruence generating a local Rindler horizon, the Raychaudhuri equation

$$\frac{d\theta}{du} = -\frac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} - 8\pi G T_{\mu\nu}k^\mu k^\nu \tag{27}$$

relates the expansion θ to the energy-momentum flux $T_{\mu\nu}$ along null generators k^μ. Integrating over the horizon patch and assuming linearised focusing gives

$$\delta A = 8\pi G \int T_{\mu\nu}k^\mu k^\nu \, du \, d^2x_\perp, \tag{28}$$

which in Jacobson's derivation leads to the Einstein equation as a thermodynamic consistency condition.

The formal use of the Raychaudhuri equation and the identification of the area change δA with the integrated null energy flux is exactly as in Jacobson. The logic that links energy flux to an infinitesimal area change along a null surface is unchanged.

In our microstate construction, the energy flux $T_{\mu\nu}$ is *discrete*: it corresponds to a single AS photon carrying energy $\Delta E = \hbar\omega$. The area change becomes

$$\Delta A = 8\pi G \, \Delta E, \tag{29}$$

applied to the transverse patch intersected by the photon. Unlike the smooth, coarse-grained description in Jacobson, each microstate produces a calculable, individual contribution to the horizon area. Moreover, the longitudinal shift of the AS pp-wave simultaneously generates the *spatial separation* that encodes the spatial bit in the ER=EPR construction. This demonstrates concretely that each detection event both writes a bit into the matter-memory of the detectors and creates the corresponding geometric imprint, unifying information, energy, and space at the level of a single microstate.

Applying the Bekenstein–Hawking relation gives the entropy increment:

$$\Delta S = \frac{\Delta A}{4\ell_P^2}, \qquad \ell_P^2 = \frac{\hbar G}{c^3}. \tag{30}$$

Thus, the photon produces a discrete entropy contribution and a corresponding spatial increment: the horizon thermodynamics emerges microstate by microstate, rather than as a smooth, macroscopic average.

This is a concrete, microphysical realization of Jacobson's ideas. Whereas Jacobson used horizon thermodynamics as a consistency argument to recover Einstein's equations, here each quantum of energy and information — each photon — produces a tangible change in horizon area and geometry, and the same event encodes information and space simultaneously. It gives a bridge from microscopic events (photon detection, tick accumulation, polarisation bits) to emergent spacetime, providing a detailed operational picture of ER=EPR and the origin of horizon entropy in a single, physically transparent step.

6.2 Planck-Unit Horizon Area from a Single Photon

Each AS pp-wave microstate corresponds to a single photon transferring energy $\Delta E = \hbar\omega$ to an accelerated detector. From the Bekenstein–Hawking relation, the associated horizon area increment is

$$\Delta A = 4\ell_P^2\,\Delta S = 4\ell_P^2 \frac{\Delta E}{T_{\text{eff}}}, \tag{31}$$

where T_{eff} is the effective Unruh temperature seen by the detector.

In Planck units, this gives

$$\Delta A \sim 8\pi\ell_P^2 \frac{\hbar\omega}{m_P c^2}, \tag{32}$$

where $m_P = \sqrt{\hbar c/G}$ is the Planck mass. Each photon therefore imprints a discrete patch of area on the horizon, measured in multiples of ℓ_P^2, linking the minimal quantum of action $\hbar$ directly to the minimal increment of horizon area.

This is a special case of the standard Bekenstein–Hawking law: instead of considering a macroscopic black hole or a coarse-grained ensemble, we see that a single actionable quantum (a photon) produces a calculable Planck-scale increment. It makes the horizon *operationally granular*, with each Planck-area patch corresponding to a single measurement event, a single Landauer tick, and a single spatial bit. This establishes a concrete bridge between quantum information, energy transfer, and emergent geometry.

7 Conclusion

8 Acknowledgements

I'm also grateful to Claude and ChatGPT for helpful discussions.

References

[1] T. Jacobson, "Thermodynamics of spacetime: the einstein equation of state," *Physical review letters*, vol. 75, no. 7, p. 1260, 1995.

[2] E. Verlinde, "On the origin of gravity and the laws of newton," *Journal of High Energy Physics*, vol. 2011, no. 4, pp. 1–27, 2011.

[3] R. Penrose, "On gravity's role in quantum state reduction," *General relativity and gravitation*, vol. 28, no. 5, pp. 581–600, 1996.

[4] L. Diosi, "A universal master equation for the gravitational violation of quantum mechanics," *Physics letters A*, vol. 120, no. 8, pp. 377–381, 1987.

[5] R. Landauer, "Irreversibility and heat generation in the computing process," *IBM journal of research and development*, vol. 5, no. 3, pp. 183–191, 1961.

[6] W. G. Unruh, "Notes on black-hole evaporation," *Physical Review D*, vol. 14, no. 4, p. 870, 1976.

[7] K. Player, "Driving the unruh response," *arXiv preprint arXiv:2509.16710*, 2025.

[8] C. C. Gerry and P. L. Knight, *Introductory quantum optics*. Cambridge university press, 2023.

[9] N. D. Birrell and P. C. W. Davies, "Quantum fields in curved space," 1984.

[10] H. J. Bremermann, "Optimization through evolution and recombination.," *Self-organizing systems*, pp. 93–106, 1962.

[11] P. C. Aichelburg and R. U. Sexl, "On the gravitational field of a massless particle," *General Relativity and Gravitation*, vol. 2, no. 4, pp. 303–312, 1971.