A Role for Quantum Measurement in Gravity

Kevin Player*

April 5, 2026

Abstract

Thermodynamic approaches to gravity, from Jacobson's derivation of the Einstein equations to Verlinde's entropic force, operate at the level of bulk averages and leave open the question of what physical process underlies each entropy increment. We propose that the answer is quantum measurement: each act of information acquisition by a physical observer constitutes a discrete microstate event whose energy-momentum backreaction gives rise to gravitational effects. Standard results, horizon scales, the Bekenstein–Hawking area law, and entropy bounds, follow from Landauer's principle at the Unruh temperature as consequences of Newton's laws. We ground this in a concrete bilocal microstate construction, extending our previous work, in which each measurement event deposits one quantum of action $\hbar$ and one classical bit into the matter substrate, with matter serving as the persistent carrier of information and the horizon as a boundary rather than a storage medium. This provides an operational realisation of ER=EPR at the level of individual microstates, converting entanglement into a spatial bit that encodes which side of the horizon each measurement occurred on.

1 Introduction

Two landmark results establish that gravity and Unurh thermodynamics are deeply intertwined. Jacobson showed that the Einstein equations can be derived from thermodynamic identities applied to local Rindler horizons, with horizon entropy playing the role of the fundamental quantity from which spacetime dynamics emerge [1]. Verlinde extended this perspective, arguing that gravitational forces arise from entropy gradients on holographic screens, recasting gravity as an entropic rather than a fundamental interaction [2]. Both frameworks, however, operate at the level of bulk thermodynamics and leave open a prior question: what physical process underlies each individual entropy increment?

We argue that the answer is quantum measurement. Each act of information acquisition by a physical observer constitutes a discrete microstate event, and the collective backreaction of such events, together with the causal correlations they induce, gives rise to gravitational effects. The argument proceeds in three steps. First, any observation is mediated by a physical interaction: information transfer necessarily involves an exchange of energy and momentum between the observed system and the detector, a discrete impulse. Second, by the equivalence principle, this impulse is locally indistinguishable in that from the effect of a gravitational field: no local measurement can distinguish the recoil associated with a quantum measurement event from acceleration in a gravitational potential. Third, when such impulses are modelled geometrically, via Aichelburg–Sexl (AS) pp-wave solutions, as we explore, they imprint energy and information directly onto horizon geometry, and coarse-graining over the resulting ensemble recovers standard horizon thermodynamics. A unifying structure runs through the relations, and it is worth pausing to discuss the general ideas involved before diving into our explicit example.

* *kjplaye@gmail.com*

1.1 Relation to recent existing Literature

The idea that spacetime geometry is connected to quantum information has been developed along several distinct lines, none of which supplies the microstate-level operational picture we develop here. Jacobson's derivation of the Einstein equations from horizon thermodynamics [1] and Verlinde's entropic gravity programme [2] both operate at the level of bulk thermodynamic averages, leaving open the question of what physical process underlies each individual entropy increment. Van Raamsdonk [3] argued that entanglement between boundary degrees of freedom is what holds bulk geometry together, and that reducing entanglement tears spacetime apart, a result that motivates our identification of the ER=EPR bridge with correlated measurement events, though his framework operates with thermal boundary states rather than individual non-thermal microstates. Susskind and Maldacena's ER=EPR correspondence [4] identifies entangled pairs with Einstein–Rosen bridges at the level of bulk thermodynamics; we realize this correspondence operationally at the level of a single microstate, with a calculable geometric imprint sourced by a specific bilocal emission event.

The closest antecedent to our geometric construction is the gravitational shockwave programme of Dray and 't Hooft [5] and 't Hooft's subsequent S-matrix approach to black hole scattering [**?**]. solutions sourced by infalling classical or semiclassical particles to define scattering amplitudes at the horizon, and the longitudinal position shift of the AS wave is the fundamental horizon degree of freedom in their picture, in precise agreement with our geometric construction. The key difference is that 't Hooft's sources are classical infalling particles rather than quantum entangled pairs emitted from a common ancestor in the past, and the measurement and information acquisition framing is absent from his construction. Our bilocal source selects individual non-thermal microstates from the Unruh ensemble, giving each AS imprint a definite quantum origin and connecting the geometric shift directly to a classical bit acquired by an observer.

Microstate counting for black holes has been approached from several directions. Strominger and Vafa [6] achieved exact entropy counting for extremal black holes using D-brane configurations in string theory, establishing that black hole entropy has a microscopic origin, but the microstates are defined algebraically in a high-energy string framework far removed from the operational measurement picture developed here. Carlip [**?**] derived black hole entropy from the Virasoro algebra of diffeomorphisms at the horizon boundary, identifying microstates with boundary excitations; our AS pp-wave imprints may provide a physical realization of these boundary excitations in terms of measurement events a connection we make precise in Section 5: the AS longitudinal shift is a horizon supertranslation in the sense of Hawking, Perry, and Strominger [7], and the bilocal construction supplies the physical process — quantum measurement — that excites each mode of the algebra.

Bianchi's loop quantum gravity microstate counting [**?**] assigns discrete area increments to individual spin network punctures at the horizon, structurally analogous to our result that each measurement event contributes a discrete AS-wave area increment; the difference is that our punctures have a definite dynamical origin in a quantum measurement event rather than a kinematical definition in the LQG Hilbert space. Interior reconstruction via the Papadodimas–Raju programme [**?**] addresses how degrees of freedom behind the horizon are encoded in exterior operators, a question closely related to our spatial bit argument in Section 6. Their framework requires thermality of the boundary state, whereas our construction works microstate by microstate with a non-thermal source; it would be interesting to determine whether our classical spatial record, the anticorrelated helicity bits registered by right and left wedges, realizes the interior reconstruction map at the single-microstate level, but we leave this comparison for future work.

What distinguishes the present construction from all of these approaches is the combination of three features that do not appear together elsewhere in the literature: a non-thermal bilocal source that selects individual microstates from the Unruh ensemble

rather than assuming thermality as a primitive; the identification of each microstate with a specific quantum measurement event carrying a calculable energy-momentum transfer; and an operational realization of entanglement-induced spatial separation at the single-microstate level with a concrete geometric imprint. Each of these features appears in isolation in various forms across the literature, but their combination, and in particular the grounding of horizon geometry in individual measurement events rather than bulk thermodynamic averages, appears to be new.

1.2 Conceptual Roadmap

Our conceptual argument is a relation between adjacent concepts that have the same logical character: every instance of the more abstract concept is grounded in an instance of the more physical one, but not every instance of the more physical concept gives rise to the more abstract one *for a given observer*. The gap in each case is filled by subjectivity. Quantum state reduction is always backed by an observation, it is a Bayesian update on the observer's part, but an observation that carries no new information for that observer produces no state reduction. Equally, every observation is backed by an energy-momentum transfer, but not every such transfer constitutes an observation for you: an interaction involving a different "observer" leaves your knowledge unchanged. The Unruh effect suggests the same structure extends to the equivalence principle: an accelerating observer perceives a thermal bath and attributes it to a gravitational field, while an inertial observer sees only vacuum. Whether an acceleration counts as gravity is perspectival in an analogous but distinct sense: not epistemic, as with Bayesian update, nor operational, as with energy–momentum transfer, but *geometric*, underdetermined by any local measurement rather than by the observer's knowledge or participation. For a Rindler observer, the perceived gravitational field just is the Unruh thermal bath, which has no observer-independent meaning. In each case the relation is an identification rather than a causal link, and the identification is local and observer-relative, though the sense of observer-relativity differs across the three arrows.

This stands in sharp contrast to the Diósi–Penrose objective collapse programme [8, 9], which postulates that gravity *causes* quantum state reduction as an objective, observer-independent physical process. Figure 1 summarises this contrast. The three solid arrows represent the perspectival grounding relations described above; the single dashed arrow represents the Diósi–Penrose causal postulate. It is the only relation in the diagram claimed to be objective and observer-independent, and it is the only one we do not adopt.



Figure 1: Solid arrows denote "substrate" relations: $A \to B$ means every instance of B is physically realized by an instance of A, for a given observer. The dashed arrow denotes the Diósi–Penrose postulate, where the relation is claimed to hold objectively rather than observer-relatively.

To make this concrete, we build on the microstate construction of [10], in which bilocal driving sources generate entangled Rindler excitations that realize individual terms of

the Unruh thermal ensemble. The present paper applies and extends that construction. Entangled pairs emitted from a localised source propagate into left and right Rindler wedges; polarisation measurement events produce discrete energy-momentum impulses and correlated classical bits. We show that each such microstate carries a calculable gravitational imprint, modelled by an AS pp-wave solution, and that coarse-graining over the ensemble recovers standard horizon thermodynamics. This construction also provides an operational realization of entanglement-induced spatial separation at the level of individual microstates: entanglement is converted into geometric information, and each measurement event contributes a discrete, calculable increment to horizon entropy.

1.3 Subjectivity and Existing Ontologies

A closely related perspective is provided by relational and information-theoretic interpretations of quantum theory. Carlo Rovelli's relational quantum mechanics [11] and the QBist programme associated with Christopher Fuchs [12] both treat the quantum state as observer-dependent, encoding information rather than objective physical reality. In these frameworks, measurement is understood as an update of an observer's knowledge, not a universal physical collapse. The Unruh effect [13] provides a direct physical analogue of this observer-relativity: particle content and thermality themselves become observer-dependent notions, tied to acceleration. Our construction extends this shared principle across domains, identifying quantum state reduction, thermality, and gravitational effects as different aspects of observer-relative descriptions grounded in underlying physical interactions. This viewpoint also sharpens the conceptual tension highlighted by Eugene Wigner [14], where different observers may consistently assign different physical descriptions to the same process, suggesting that the resulting geometric imprints in our framework may likewise be observer-dependent. Finally, the present approach may be viewed as a concrete dynamical realization of Wheeler's "it from bit" programme [15]. Where Wheeler's proposal remained largely conceptual, discrete acts of information acquisition giving rise to physical law, we identify an explicit dynamical mechanism: each bit acquired by a physical observer is backed by a measurement-induced energy-momentum transfer, in our example, modelled geometrically as an AS pp-wave, whose gravitational imprint incrementally builds spacetime structure. The bit does not merely describe the geometry; it produces it.

2 Motivating Heuristics

Landauer's principle associates a minimal energy cost with the acquisition of information [16]. For an accelerated observer, the relevant temperature is the Unruh temperature [13]. Substituting one into the other eliminates temperature entirely, yielding a relation between information acquisition, energy transfer, and acceleration (a version of the BHT first law, equation (3)). This is not merely an algebraic convenience: it is a concrete dictionary between the thermal description natural to a Rindler observer and the non-thermal global Minkowski description, with the apparent thermality of the vacuum emerging as a consequence of the observer's acceleration rather than as a fundamental property of the state. In this way we derive a sequence of standard results in black hole thermodynamics as a consequence of Newton's laws.

2.1 The Unruh-Landauer Relation

Landauer's principle suggests that the acquisition of ΔH bits of information by a physical observer requires a minimum energy transfer

$$\Delta E \geq k_B T \, \Delta H. \tag{1}$$

We note that Landauer's principle is conventionally stated for information erasure rather than acquisition. Here we apply it to acquisition, which is the physically relevant process for an observer registering a measurement outcome. The bound has the same form in both cases; we apply it to acquisition on the grounds that registering a measurement outcome is physically equivalent to a bit being written into the detector's state, which is the process Landauer's principle governs.

For an observer undergoing proper acceleration a, the Unruh effect associates a temperature

$$T = \frac{\hbar a}{2\pi k_B c} \tag{2}$$

with the local vacuum. Substituting into Landauer's bound eliminates $k_B T$, yielding

$$\boxed{\Delta E \geq \frac{\hbar a}{2\pi c}\,\Delta H.} \tag{3}$$

Temperature has dropped out entirely: equation (3) is a temperature-independent relation between energy, acceleration, and information, suitable for the Minkowski vacuum. Acceleration sets the energy scale of information. In what follows, standard results emerge as consequences of this single relation and Newton's laws.

2.2 Momentum Transfer and Acceleration

The energy transfer ΔE implied by equation (3) is carried by a physical quantum — matter or radiation — mediating the measurement interaction. We refer to this as the *thrust quantum* of effective mass m_t. (This will actually be a pure energy carrier in our later microstate example.) For a relativistic carrier, the associated momentum transfer is

$$\Delta p = \frac{\Delta E}{c}. \tag{4}$$

A measurement event localised to a spatial scale r is bounded by a causal diamond: the intersection of the future light cone of its initiation and the past light cone of its completion. The temporal extent of this diamond is $\Delta t = 2r/c$, fixed entirely by the causal geometry of the event rather than by any dynamical assumption. We identify the interaction time with this natural causal duration, so that

$$a = \frac{\Delta p}{m_d \Delta t} = \frac{\Delta E}{2 m_d r}. \tag{5}$$

This perspective is closely related in spirit to Erik Verlinde's entropic gravity [2], where forces arise from entropy gradients, but differs in that the present construction grounds each entropy increment in a discrete measurement event with an associated energy-momentum transfer, rather than in a coarse-grained thermodynamic gradient.

2.3 The Schwarzschild Radius

We match the acceleration inferred from the measurement-induced impulse to the Newtonian gravitational field in order to identify the characteristic length scale associated with the detector. The thrust quantum carries energy ΔE and produces an acceleration a on the detector. Identifying this with the Newtonian gravitational acceleration due to the mass m_t at distance r.

$$a = \frac{G m_t}{r^2}, \tag{6}$$

and substituting equation (5) with $\Delta E = m_t c^2$, we obtain

$$\frac{m_t c^2}{2 m_d r} = \frac{G m_t}{r^2}. \tag{7}$$

Solving for r:

$$r = \frac{2Gm_d}{c^2} \equiv r_s, \tag{8}$$

which is precisely the Schwarzschild radius of the detector mass m_d.

It is notable that the resulting length scale is set by the detector mass m_d, rather than the source mass m_t. This reflects the fact that the construction is inherently observer-centred: the acceleration entering the argument is the response of the detector to the measurement-induced impulse, and the associated scale is therefore the horizon radius relevant to that observer. In this sense, the quantity r_s should be interpreted not as the Schwarzschild radius of the source, but as the characteristic horizon scale associated with the detector as a physical system undergoing measurement. This is consistent with the observer-dependence of horizon structure in relativistic settings, as exemplified by the Unruh effect, where acceleration alone gives rise to a causal horizon.

2.4 Surface Gravity

Evaluating the acceleration at $r = r_s$ gives the surface gravity of the detector:

$$\kappa = \frac{Gm_d}{r_s^2} = \frac{c^4}{4Gm_d}, \tag{9}$$

in agreement with the standard Schwarzschild result. This is the acceleration that appears in the Unruh temperature associated with the detector's horizon, $T = \hbar\kappa/2\pi k_B c$, providing an internal consistency check on the framework.

2.5 Bekenstein-Hawking Entropy

Returning to the Unruh-Landauer relation (3) and substituting the surface gravity κ for a, the entropy increment associated with an energy increment $\Delta E = c^2 \Delta m_d$ is

$$\Delta H \leq \frac{2\pi c}{\hbar\kappa}\Delta E = \frac{8\pi G m_d}{\hbar c^3} \cdot c^2 \Delta m_d. \tag{10}$$

Expressing this in terms of the change in Schwarzschild area $\Delta A = 32\pi G^2 m_d \Delta m_d / c^4$:

$$\Delta H \leq \frac{\Delta A}{4\ell_P^2}, \qquad \ell_P^2 = \frac{\hbar G}{c^3}, \tag{11}$$

which is the Bekenstein-Hawking area law. Each measurement event contributes a discrete entropy increment proportional to the area imprinted on the horizon by the associated energy-momentum transfer.

This derivation parallels Jacob Bekenstein's original argument [17], in which the entropy increase of a black hole is inferred from the minimal energy required to localize and drop a system across the horizon; here, the same structure emerges from the minimal energy cost of registering information at the horizon scale.

The two bounds are saturated simultaneously when $a = c^2/R$ — which is exactly the condition for being *at a horizon*. The horizon is therefore not just where thermodynamics happens to work out, it is the unique locus where the minimum cost of information acquisition equals the maximum information content of the region.

2.6 Summary

Beginning from the single relation (3), and using only Newton's laws and the equivalence principle, we have recovered the Schwarzschild radius, the surface gravity, and the Bekenstein-Hawking area law as consequences of the observation $\rightarrow$ acceleration identification. These are standard results, but their derivation here from the physical cost of information acquisition rather than from thermodynamic postulates is the point.

The thrust quantum introduced in this section, the physical carrier of the measurement-induced impulse, will be given a explicit microstate realization in the sections that follow. It is identified with one of a pair of entangled photons modeled as Aichelburg-Sexl pp-waves crossing past Rindler horizons in the Unruh picture. In this case the back reaction of a single force carrier in one wedge is correlated with an equal and opposite reaction in the other from a common source.

3 Bilocal Microstate Construction

Let $\hbar = c = 1$.

The heuristic arguments of Section 2 establish a relationship between information acquisition, acceleration, and gravitational effects, but leave open the question of what a single microstate looks like concretely. In standard treatments, the Unruh effect is derived by coupling a detector to a pre-existing acceleration and reading off a thermal response from the vacuum. Following [10], we invert this logic: we place an interaction in the past, where a source injects non-thermal entangled excitations into the field, and show that the observer's accelerated response can be driven by these source-induced excitations. The thermal ensemble then emerges as a coarse-grained aggregate of such microstates, rather than being a primitive property of the vacuum.

Each microstate, equation ((22)) upcomming, is individually non-thermal: it carries a definite helicity pair and a definite Rindler frequency Ω, neither of which is fixed by the Unruh temperature. Thermality is not a property of any single microstate but of the ensemble; it emerges when one asks only which wedge a detection occurred in, discarding the specific microstate identity, and is a consequence of coarse-graining rather than a premise of the construction.

We now upgrade this construction from a spin-0 scalar field in the original source [10] to the spin-1 electromagnetic field. This is not merely a technical generalisation. As we show, the photon carries a unit of action $\hbar$ per cycle alongside its energy $E = \hbar\omega$, which closes the Landauer gap left open in Section 2: in each wedge, the minimum actionable event in the construction is precisely a single photon, and its polarisation state constitutes the classical bit acquired by each observer from detection. The anticorrelated helicities of the entangled pair are the natural completion of the $\mathbb{Z}_2$ swap symmetry between wedges, and as we discuss in Section 5, they imprint a definite geometric signature onto the horizon consistent with the emergent bridge geometry.

3.1 A Source and Three Observers

Three observers are placed at spacetime positions $(t, z) = (0, 1)$, $(0, -1)$, and $(-1, 0)$, named $\mathcal{R}$ - *Right*, $\mathcal{L}$ - *Left*, and $\mathcal{P}$ - *Past* respectively. $\mathcal{P}$ serves as the common causal ancestor of $\mathcal{R}$ and $\mathcal{L}$ (see Figure 2).

$\mathcal{P}$ emits an entangled pair of spin-1 massless photons with momenta $\pm k$ and Rindler frequency $\omega = \omega_k$. The momenta are perfectly balanced, so the source exhibits no net back-reaction in this minimal model. The two photons propagate into the left and right Rindler wedges, where $\mathcal{L}$ and $\mathcal{R}$ detect them respectively, each registering a helicity eigenvalue and acquiring one classical bit of information.

The three-party structure $\mathcal{P}$–$\mathcal{R}$–$\mathcal{L}$ is a tripartite entanglement: neither $\mathcal{R}$ nor $\mathcal{L}$ alone holds sufficient information to reconstruct the emission event at $\mathcal{P}$, but their two complementary bits together do, making $\mathcal{P}$ the logical AND of the pair and providing a natural microstate-level realization of interior reconstruction.



Figure 2: A source in the past, $\mathcal{P}$, emits an entangled photon pair that accelerates $\mathcal{R}$ into the right Rindler wedge and $\mathcal{L}$ into the left Rindler wedge. The two null directions of propagation define the z^+ and z^- axes of the double-null coordinate system.

3.2 Double-Null Gauge

The emission event naturally defines two null directions: the right-moving photon has wave-vector k_R^μ and the left-moving photon has wave-vector $k_L^\mu = -k_R^\mu$. These two null directions are precisely the z^+ and z^- axes of double-null coordinates

$$z^\pm = \frac{1}{\sqrt{2}}(t \pm z), \tag{12}$$

with transverse coordinates $\mathbf{z}_\perp = (z^1, z^2)$. Working in this frame we impose the symmetric double-null gauge condition

$$A_+(z) = 0, \qquad A_-(z) = 0, \tag{13}$$

so that the gauge field is supported entirely on the transverse components $\mathbf{A}_\perp = (A^1, A^2)$. This gauge respects the $\mathbb{Z}_2$ symmetry that swaps the two wedges ($z^+ \leftrightarrow z^-$) simultaneously with ($R \leftrightarrow L$), and therefore treats both observers symmetrically by construction. The residual gauge freedom is fixed by requiring $\partial_i A^i = 0$ in the transverse plane, leaving precisely the two physical helicity degrees of freedom $\epsilon^i_{(\lambda)}$ with $\lambda = \pm 1$; consistency with Maxwell's equations and the absence of over-determination are verified in Appendix 7.

The lack of manifest four-dimensional covariance in equation (13) is a consequence of aligning the gauge with the pair of physical null directions selected by the source, rather than an arbitrary choice. The full Lorentz group is broken to the little group preserving the pair $\{k_R^\mu, k_L^\mu\}$, which is the two-dimensional Euclidean group $ISO(2)$ acting on the transverse plane. This is precisely the little group of massless particles, and the two helicity eigenstates $\lambda = \pm 1$ are its irreducible representations. All physical results are independent of the gauge choice.

3.3 The Bilocal Source

In double-null gauge the electromagnetic field is characterised by its transverse components alone. The bilocal driving source generalises the scalar construction to

$$J^{ij}(x, y) = \lambda\, \epsilon^{i*}_{(-1)}\, \epsilon^j_{(+1)}\, f_L(x)\, f_R(y), \tag{14}$$

where λ is a coupling constant, f_L and f_R are normalised mode profiles supported on the left and right wedges respectively, and $\epsilon^i_{(\lambda)}$ are the circular polarisation vectors

$$\epsilon^i_{(+1)} = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\ i\end{pmatrix}, \qquad \epsilon^i_{(-1)} = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\ -i\end{pmatrix}. \tag{15}$$

The index structure of (14) assigns helicity $+1$ to the right-moving mode and helicity -1 to the left-moving mode: the pair is *anticorrelated* in helicity. This is the unique assignment consistent with angular momentum conservation at the emission vertex and with the $\mathbb{Z}_2$ wedge-swap symmetry, under which $\epsilon^i_{(+1)} \leftrightarrow \epsilon^i_{(-1)}$.

The sourced Lagrangian reads

$$\begin{aligned}\mathcal{L}_{\text{sourced}} &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \int J^{ij}(x,y)\, A^+_i(x)\, A^+_j(y)\, d^4x\, d^4y \;+\; \text{h.c.} \\ &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \Big(a^{L\dagger}_{-k,(-1)}\, a^{R\dagger}_{k,(+1)} \;+\; a^{L}_{-k,(-1)}\, a^{R}_{k,(+1)} \Big),\end{aligned} \tag{16}$$

where $a^{L\dagger}_{-k,(-1)}$ and $a^{R\dagger}_{k,(+1)}$ are creation operators for left-wedge helicity -1 and right-wedge helicity $+1$ modes respectively. The interaction term in (16) is a two-mode squeezing interaction at frequency ω, now resolved by helicity: it is the direct vector-field analogue of the scalar squeezing interaction [18], with the anticorrelated polarisation structure made explicit. The photon pair carries total angular momentum zero, consistent with emission from a scalar source at $\mathcal{P}$.

Each photon carries energy $E = \hbar\omega$ and, as a mode of the electromagnetic field, constitutes precisely one quantum of action $\hbar$ per oscillation cycle. The thrust quantum of Section 2 is thereby identified with a single photon: the minimum actionable event is the emission and detection of one such quantum.

The Unruh–Landauer bound is saturated not at the level of an individual photon but at the level of the source rate. The acceleration a does not fix the frequency ω of each emitted quantum; rather, it determines the rate $\dot{N}$ at which the bilocal source must produce entangled pairs in order to fully[1] implement that acceleration, with the total power $\dot{N}\hbar\omega$ matching the Unruh thermal flux. Each individual photon carries exactly one quantum of action $\hbar$ and one binary helicity outcome, one bit in the sense of one binary alternative, carrying $\ln 2$ nats, and these are paired by the construction independently of a. The Landauer bound then operates as a consistency condition on the ensemble: the minimum energy cost per bit acquired, summed over the rate of acquisition, equals the power required to sustain the acceleration.

3.4 Polarisation Bogoliubov Transformation

We now show that the polarisation labels survive the Rindler–Unruh mode decomposition, so that the thermal ensemble retains its helicity structure and the anticorrelation of the emitted pair descends unambiguously to the level of individual Rindler microstates.

In the transverse plane, the electromagnetic field in double-null gauge decomposes into two independent scalar-like sectors labelled by helicity $\lambda = \pm 1$. For each sector, the right-wedge field expands as

$$A^i_\lambda(z) = \int_0^\infty \frac{d\Omega}{2\pi} \frac{1}{\sqrt{2\Omega}}\, \epsilon^i_{(\lambda)} \left[b^R_{\Omega,\lambda}\, u^R_\Omega(z) + b^{R\dagger}_{\Omega,\lambda}\, u^{R*}_\Omega(z) \right], \tag{17}$$

where $u^R_\Omega(z)$ are the right-wedge Rindler modes at Rindler frequency Ω, and $b^R_{\Omega,\lambda}$, $b^{R\dagger}_{\Omega,\lambda}$ are the corresponding annihilation and creation operators satisfying

$$\left[b^R_{\Omega,\lambda},\, b^{R\dagger}_{\Omega',\lambda'} \right] = \delta(\Omega - \Omega')\, \delta_{\lambda\lambda'}. \tag{18}$$

An identical expansion holds in the left wedge with $R \to L$ and $\lambda \to -\lambda$, reflecting the anticorrelated helicity assignment. Because each helicity sector is independently a scalar-like sector, the Bogoliubov transformation relating Minkowski and Rindler modes takes the same form in each sector. For helicity λ at Rindler frequency Ω, the Unruh modes are

$$c_{\Omega,\lambda} = \cosh\theta_\Omega\, b^R_{\Omega,\lambda} - \sinh\theta_\Omega\, b^{L\dagger}_{\Omega,-\lambda}, \tag{19}$$

[1] The bilocal source may be only part of the acceleration, so $\dot{N}$ is actually a lower bound.

where $\tanh\theta_\Omega = e^{-\pi\Omega c/a}$ encodes the Unruh distribution at acceleration a, and $-\lambda$ on the left-wedge operator reflects the anticorrelation. The transformation (19) is the standard Bogoliubov relation for a massless field [13, 19], now carrying a helicity label throughout. The Minkowski vacuum in the Rindler basis reads, for each helicity sector,

$$|0\rangle_M^{(\lambda)} = \frac{1}{\cosh\theta_\Omega}\sum_{n=0}^{\infty}\tanh^n\theta_\Omega\,|n_{\Omega,\lambda}\rangle_R\,|n_{\Omega,-\lambda}\rangle_L, \tag{20}$$

and the full vacuum is the product over all Ω and both helicities:

$$|0\rangle_M = \prod_{\Omega,\lambda}|0\rangle_M^{(\lambda)}. \tag{21}$$

Equation (20) makes explicit that the Minkowski vacuum is a sum over entangled helicity-anticorrelated pairs at each Rindler frequency. The bilocal source selects a single term from this sum at fixed Ω and fixed helicity pair $(\lambda, -\lambda) = (+1, -1)$, yielding the single-microstate

$$|\Psi_\Omega\rangle_M = |1_{\Omega,+1}\rangle_R\,|1_{\Omega,-1}\rangle_L + \mathcal{O}(\text{higher order}). \tag{22}$$

$\mathcal{R}$'s detection of helicity $+1$ and $\mathcal{L}$'s detection of helicity -1 are therefore perfectly anticorrelated: each measurement event yields one classical bit, and the two bits are the complement of one another.

The key structural result is that the helicity label λ is conserved under the Bogoliubov transformation: it enters equation (19) as a spectator index and exits unchanged on the Unruh operators $c_{\Omega,\lambda}$. Thermality, when it arises from coarse-graining over the ensemble of microstates (22), is therefore helicity-resolved: each helicity sector thermalises independently, and the anticorrelation between wedges is preserved at every level of the construction.

3.5 Summary

The passage from the scalar to the vector field construction introduces three structural improvements. First, the photon carrying $E = \hbar\omega$ and one quantum of action per cycle identifies the thrust quantum of Section 2 with a concrete physical object, closing the Landauer acquisition gap. Second, the double-null gauge condition, motivated by the pair of null directions defined by the emission event, treats both Rindler wedges symmetrically and exposes the $\mathbb{Z}_2$ swap symmetry as the organising principle of the construction. Third, the anticorrelated helicity structure, conserved through the Bogoliubov transformation, provides each observer with one classical bit whose values are perfectly complementary: the two bits together constitute the correlated classical record that, in the AS pp-wave picture of Section 5, is imprinted as geometric information onto the horizon.

4 Matter as Substrate

The Unruh–Landauer relation of Section 2 constrains the minimal energy required to register information, but does not specify where that information resides physically. The present section addresses this gap by identifying matter as the substrate in which such information is stored, with each act of acquisition corresponding to the deposition of a discrete quantum of action into the detector's degrees of freedom. We show how information is encoded in matter and energy and is reflected in the horizon geometry. This requires a physical account of what it means for matter to *carry* information.

A massive particle accumulates phase along its worldline at a rate set by its rest energy. The Compton frequency relation $f = E/h$ assigns a natural frequency to any mass m, and Bremermann's computational limit [20] independently recovers the same rate $f = mc^2/h$

as the maximum clock rate, or Nyquist rate, of any physical system with that energy E. These two results agree because they are both expressions of the same underlying fact: a quantum of action $\hbar$ is the minimal unit of physical evolution, and mc^2/h is the rate at which a mass m consumes such quanta.

Each such quantum corresponds to one sample[2] of the phase $e^{iS/\hbar}$ in the path integral, and classical trajectories emerge from the constructive interference of these samples across paths. Paths separated by less than one sample of action, less than $\hbar$, do not produce resolvable interference fringes: the phase difference is insufficient to complete a full cycle of constructive and destructive interference, so the trajectory cannot be distinguished at sub-sample resolution. The trajectory is resolved only down to the scale set by $\hbar/E$, which is precisely the Heisenberg uncertainty at energy E. This corresponds to a natural ultraviolet cutoff given by the energy E of the system itself. Higher energies only come into play when external systems interact with the energy E system.

This gives matter a precise sense in which it is a *substrate for information*: it accumulates samples, and each sample is a discrete unit of action, Lorentz-invariant and recorded in the particle's proper time. Matter is persistent memory in the operational sense that its history of samples is physically encoded in its state and cannot be erased without an energy cost governed by Landauer's principle. The maximum rate of this memory is bounded by Bremermann.

> *The signal being sampled at the Bremermann rate is the internal quantum state of the particle, its spin, helicity, and other quantum numbers, and the dynamic range is* $\log_2$ *of the dimension of the internal Hilbert space.*

It is worth pointing out that these concepts are additive. The union of two systems with masses m_1 and m_2 and internal Hilbert spaces of dimensions d_1 and d_2 has a sampling rate of

$$\frac{mc^2}{\hbar} = \frac{(m_1 + m_2)c^2}{\hbar} = \frac{m_1 c^2}{\hbar} + \frac{m_2 c^2}{\hbar}, \tag{23}$$

and a dynamic range of

$$\log_2(d) = \log_2(\dim(H_1 \otimes H_2)) = \log_2(d_1 d_2) = \log_2(d_1) + \log_2(d_2), \tag{24}$$

for joint mass $m = m_1 + m_2$ and joint dimension $d = d_1 d_2$. Both the sampling rate and the information capacity are therefore extensive quantities, consistent with their interpretation as properties of matter that compose naturally under union.

The distinction between massless and massive degrees of freedom is therefore operational: massless quanta transmit information, while massive systems retain it, accumulating the action quanta that constitute physical memory. A massless particle, the photon, has no rest frame, no proper time, and accumulates no action between emission and absorption. It does not sample. It therefore carries no persistent memory and cannot serve as a substrate. Its role in the present construction is precisely that of a pure information carrier. A single photon carries exactly two internal irreducible constituents, distinct in kind:

- **One quantum of action** $\hbar$: the physical transfer, deposited as energy $\hbar\omega$ into the detector, incrementing its Bremermann sample rate by $\Delta f = \hbar\omega/h$ for as long as the energy remains in the mass. This increment is the physical trace of the interaction in the detector's matter-memory.
- **One bit of polarisation**: the informational outcome, the classical record of which helicity, $\lambda = +1$ or $\lambda = -1$ — was absorbed. This bit persists in the detector's state for as long as the detector retains the record. Erasure of this bit carries a minimum energy cost governed by Landauer's principle, regardless of how the record was originally established.

[2] We use *sample* throughout to denote a single zero crossing of the phase $e^{iS/\hbar}$; readers may equivalently read this as *event* or *tick*. It is only defined upto the Heisenberg resolution.

4.1 Types of sample

Not every sample produces information, and not all information is produced the same way. We distinguish three classes of sample, ordered by their informational character.

- **Absorption samples** are measurement events: a photon is detected, one quantum of action $\hbar$ is deposited persistently into the detector's matter-memory, and one helicity outcome is written as a classical bit. The Landauer cost $\Delta E \geq k_B T \ln 2$ is paid immediately and in full on acquisition. This is the sample category that drives the bilocal construction: $\mathcal{R}$ and $\mathcal{L}$ each undergo exactly one absorption sample per emitted pair, and it is these samples whose gravitational imprint is modelled by the AS pp-wave in Section 5.
- **Free samples** are pure phase accumulation events: a massive particle propagates along its worldline, incrementing its sample count at the Bremermann rate mc^2/h, without writing any bit into any detector's state. No Landauer cost is incurred, no classical record is produced, and no observer acquires information. These are the samples that constitute mass and proper time in the background sense, the substrate on which absorption and emission events act.
- **Emission samples** are the time-reverse of absorption: a bit stored in matter-memory is released, the corresponding quantum of action is re-emitted as radiation, and the Landauer cost is paid on the way *out* rather than on the way in. This is the category Landauer's principle most directly governs in its original erasure formulation.

A moving mirror realizes absorption and emission as reflection: it receives a photon, stores the helicity bit transiently in its matter degrees of freedom during a period of free sample accumulation, and re-emits the photon, paying the erasure cost at emission. An ensemble of such mirrors, say two silvered mirrors back to back, can cycle through absorption, free accumulation, and emission without ever producing a permanent classical record.

It is worth noting that the absorption/free/emission classification is observer-relative in the same sense as quantum state reduction itself. What constitutes an irreversible absorption sample for one observer may be a free sample for an external observer. If $\mathcal{R}$ truly forgets the helicity outcome, in the sense that his matter-memory becomes disentangled from the bit, the photon can interfere with its partner at a future $\mathcal{P}$. The classification is not a property of the interaction itself but of the observer's epistemic access to the outcome. True irreversibility is approached asymptotically as the bit becomes correlated with an increasing number of environmental degrees of freedom, which is the standard decoherence picture now applied to the sample classification.

The $\hbar$ is what makes the sample physical; the bit is what makes it meaningful to an observer. These are distinct: the action quantum is deposited regardless of whether a persistent record is produced, while the bit exists only relative to an observer for whom the outcome is accessible. That a single photon carries one of each in the minimal case is the reason it is the natural unit of the construction.

4.2 Penrose's Zig-Zag Mechanism

The matter substrate argument of the preceding subsection established that massive particles sample at the Bremermann rate mc^2/h and that each sample is a quantum of action $\hbar$ accumulated in the path integral phase $e^{iS/\hbar}$. But it left open a prior question: *why* does matter sample at all? The zig-zag[3] construction of Penrose and Dirac supplies a concrete physical answer, and in doing so reveals that the bilocal microstate construction

[3]Penrose calls these particle *zig-zag* particles, we are tempted to call them *tick-tock* particles, but only in a footnote.

of Section 3 is not merely analogous to the origin of mass, it is structurally identical to it at the level of individual events.

4.2.1 The Dirac 4-Spinor as a Bilocal System

The Lorentz algebra decomposes as

$$\mathfrak{so}(3,1) \cong \mathfrak{su}(2) \times \mathfrak{su}(2), \tag{25}$$

where the two $\mathfrak{su}(2)$ factors correspond to the two helicities: the first is generated by boosts combined with counter-clockwise rotations, and the second by boosts combined with clockwise rotations [21]. A Dirac 4-spinor decomposes under this algebra into two Weyl 2-spinors of opposite helicity $\lambda = \pm\frac{1}{2}$ and equal and opposite momenta. Penrose [22] proposes a physical interpretation of this decomposition: the two 2-spinors are real physical objects, each moving instantaneously at the speed of light in opposite directions, interacting at regular intervals via the Higgs field at the De Broglie frequency.

The structural parallel with the bilocal construction is very close:

- The two Weyl 2-spinors, moving at c in opposite directions with opposite helicity, correspond to the two photons emitted by $\mathcal{P}$ into the left and right Rindler wedges.
- The Higgs field plays the role of $\mathcal{R}$ and $\mathcal{L}$: it is the common mediator whose interaction reflects the entangled pair and sets the interaction rate.
- Between Higgs interactions, each 2-spinor is massless and moves at c, carrying no persistent memory, pure information carriers, exactly as the photons in the bilocal construction.
- The Higgs interaction event itself is the measurement event: it deposits one quantum of action $\hbar$, flips the active helicity from $+\frac{1}{2}$ to $-\frac{1}{2}$ and back, and produces one sample of proper time.

Mass emerges from the regularity of these interactions: a particle is massive because it undergoes repeated Higgs interactions at the De Broglie frequency, and its mass is proportional to that frequency via $E = \hbar\omega$. The Bremermann sample rate mc^2/h is therefore not a coincidence or a heuristic, it is the Higgs interaction rate, the rate at which the zig-zag mechanism deposits quanta of action into the particle's proper time.

4.2.2 Mass as Accumulated Measurement History

Between Higgs interactions, the two 2-spinors are massless and moving at c in opposite null directions. Each spinor therefore has a local Rindler structure relative to the other: from the perspective of the right-moving spinor, the left-moving spinor is behind a causal horizon, and vice versa. The Higgs interaction is the event that collapses this local horizon and produces one sample, one unit of proper time, exactly as the photon absorption by $\mathcal{R}$ collapses the Rindler horizon and produces one sample in the bilocal construction.

This gives a precise sense in which mass is an accumulated measurement history. Each Higgs interaction is a measurement event: it deposits one quantum of action $\hbar$ into the particle's matter-memory, persistently incrementing its sample rate by $\Delta f = \Delta E/h$. The particle's mass is the total accumulated record of all such interactions since its formation. Matter is not just the substrate for information; its mass encodes the accumulated history of its Higgs interaction events.

4.2.3 Proper Time as a Foliation of Micro-Horizons

The series of Higgs interaction events along a particle's worldline is a one-dimensional foliation of the proper time line by discrete events, each of which is a micro-horizon separating the causal past from the causal future of that interaction, see Figure 3. This



Figure 3: A 1-dimensional (proper) time line. The red, yellow, and green dots are 0-dimensional events, in the past, present and future respectively. A screen is pictured at the present.

is the one-dimensional analogue of the Rindler horizon separating left and right wedges in the bilocal construction, and it matches the holographic screen picture of Verlinde [2]: each screen is a surface at which information is registered and a sample of proper time is produced.

The zig-zag mechanism therefore provides a physical grounding for the matter substrate argument at the level of individual particles. The bilocal construction of Section 3 is the macroscopic version of the same structure: $\mathcal{R}$ and $\mathcal{L}$ play the role of the Higgs field, the two photons play the role of the two Weyl spinors, and they produce samples of proper time in the zig-zag picture. The difference in scale is large but the logical structure is identical.



Figure 4: These figures were adapted from [22]. They show his interpretation of the Dirac equation and the Higgs mechanism for an electron.

4.3 Matter as a Physical Channel

The sampling rate and dynamic range of matter can be combined into a channel capacity in the Shannon sense. For a system of mass m with internal Hilbert space dimension d, the maximum rate of information flow is

$$C = \underbrace{\frac{mc^2}{h}}_{\text{bandwidth}} \times \underbrace{\log_2(d)}_{\text{dynamic range}} \qquad \text{(bits per second)}, \tag{26}$$

where the bandwidth is the Bremermann sampling rate and the dynamic range is the information content per sample. The product C is the *channel capacity* of the matter substrate: the maximum rate at which physical interactions can write information into its degrees of freedom.

The three sample types of Section 4 correspond to three distinct regimes of the data processing inequality $H(\text{out}) \leq H(\text{in})$, which states that information cannot be created by physical processes, only preserved or lost:

$$\text{Free:} \quad H(\text{out}) = H(\text{in}), \tag{27}$$

$$\text{Absorption:} \quad H(\text{field, out}) < H(\text{field, in}), \tag{28}$$

$$\text{Emission:} \quad H(\text{detector, out}) < H(\text{detector, in}). \tag{29}$$

Free samples saturate the inequality: no information is lost or gained, the channel runs at capacity without transmitting anything. Absorption samples strictly decrease the field's information content as it flows into the detector: the channel is being *used*, information is being written into matter-memory at a rate bounded by C. Emission samples strictly decrease the detector's information content as the stored bit is released back into the field: the channel runs in reverse, and the Landauer cost is the energy price of that reversal.

The data processing inequality is therefore the information-theoretic statement of the thermodynamic arrow of time in this framework: absorption and emission are the two directions of persistent information flow, and free samples are the reversible background against which both are measured. The full quantum channel structure of the Bogoliubov transformation, its classical capacity in the Holevo–Schumacher–Westmoreland, we leave for future work.

The Bekenstein and Bremermann bounds are therefore the same constraint viewed from orthogonal directions: Bekenstein counts information capacity spatially, as area in Planck units, while Bremermann counts it temporally, as samples per second. Integrating the channel capacity over the natural causal timescale $\tau \sim Gm/c^3$ gives

$$C \cdot \tau = \frac{mc^2}{h} \cdot \frac{Gm}{c^3} = \frac{Gm^2}{hc} \sim H_{\text{Bekenstein}}, \tag{30}$$

up to numerical factors and the dynamic range $\log_2(d)$. The spacelike and timelike information bounds are therefore consistent faces of the same underlying constraint.

4.4 Horizon Separation

The horizon enters this picture not as a storage medium but as a boundary condition. It marks the surface at which the matter-memory on one side is causally disconnected from the matter-memory on the other. Horizon entropy, in this account, is not a property of the horizon surface itself but a count of the samples that have been written into the matter on each side, bounded by Bremermann from above and by Landauer from below. The holographic appearance of this count, proportional to area rather than volume, is a consequence of the Unruh–Landauer and Bekenstein relations derived in Section 2 for a spherically symmetric mass distribution and does not require holography as an independent postulate.

This reframing has a direct consequence for the emergent spatial separation developed in Section 6 below. The correlated measurement outcomes of $\mathcal{R}$ and $\mathcal{L}$ are not encoded in the horizon geometry in the first instance, they are encoded in the matter of their respective detectors, each of which has absorbed one photon and increased its Bremermann sampling rate by $\Delta f = \hbar\omega/h$. The geometric correlations between the two wedges, realized as correlated AS pp-wave imprints in Section 5, are downstream of this matter-level encoding. The bridge, if it exists, is built from correlated samples, not from the horizon itself.

5 pp-Wave Geometry

WLOG we introduce coordinates for the photon traveling from $\mathcal{P}$ $(-1, 0)$ to $\mathcal{R}$ $(0, 1)$. Let $u = t - z + 1$ and $v = t + z + 1$, where the photon propagates along the null plane $u = 0$, z is the longitudinal direction, and (x, y) are transverse coordinates. Then $(u, v) = (0, 0)$ is $\mathcal{P}$'s location, $(u, v) = (0, 2)$ is $\mathcal{R}$'s location, and $(u, v) = (0, 1)$ is the intersection of the photon with the past right Rinder horizon. The situtation is reversed for the left moving photon, we swap u and v to describe photon moving from $\mathcal{P}$ to $\mathcal{L}$.

5.1 The AS pp-Wave and Its Effects

Test particles at fixed transverse positions (x, y) experience two effects as the shock front crosses $u = 0$:

1. **Longitudinal shift.** The v coordinate is displaced by

$$\Delta v \propto -\ln r^2, \qquad r^2 = x^2 + y^2. \tag{31}$$

   This shift is lightlike: no proper time elapses for a particle crossing the shock, but it is displaced along the null direction. Near the axis ($r \to 0$) the shift diverges, encoding the concentration of energy along the photon's trajectory.

2. **Transverse focusing.** The gradient of the longitudinal shift produces a transverse velocity

$$\Delta \mathbf{v}_\perp \propto \nabla \ln r \propto \frac{1}{r}, \tag{32}$$

   directed toward the axis. Test particles are pulled toward the centre of the wave, regardless of their initial transverse separation.

5.2 The Longitudinal Shift Makes Space

The longitudinal shift deserves closer attention, because it carries a meaning beyond a mere coordinate displacement. The photon emitted by $\mathcal{P}$ propagates along the null direction; as it crosses the Rindler horizon, the AS shift displaces the future light cone of every test particle it passes. For the photon traveling into the right wedge, the shift pushes the right wedge's geometry in the $+v$ direction; for the entangled partner traveling into the left wedge, the shift pushes the left wedge in the $-v$ direction. The two wedges are displaced *away from each other* along the longitudinal null direction. See Figure 5.



Figure 5: The green arrows show the longitudinal shift that occurs where the photons cross the horizon. Test particles, shown in yellow, are pushed outwards away from $z = 0$ due to the creation of space along the null lines originating from $\mathcal{P}$.

At the intersection of the shock front $u = 0$ with the past right Rindler horizon at $(u, v) = (0, 1)$, the transverse (x, y) plane is identified with the horizon area elements $dA = dx\,dy$. The longitudinal shift $\Delta v \propto -\ln r^2$ displaces the horizon outward in the v direction, pushing the right wedge geometry away from the horizon. The entangled partner photon produces a symmetric shift in the u direction at the left Rindler horizon, pushing the left wedge away from its horizon. The two wedges are therefore displaced away from their respective horizons in opposite null directions.

5.3 Horizon Area Increment from a Single Photon

The transverse focusing of the AS pp-wave produces a calculable increment in horizon area. We derive this using the Raychaudhuri equation on the null congruence generating the Rindler horizon, following Jacobson [1], but applied here to a single microstate rather than a thermodynamic average.

For a null congruence with generators k^μ and expansion θ, the Raychaudhuri equation gives

$$\frac{d\theta}{du} = -\tfrac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} - 8\pi G\, T_{\mu\nu}k^\mu k^\nu. \tag{33}$$

Integrating over the horizon patch in the linearised regime and using $k^\mu = \delta^\mu_u$, the area change is

$$\Delta A = 8\pi G \int T_{uu}\, du\, d^2x_\perp. \tag{34}$$

In a strict AS construction, the source is a point particle and $T_{\mu\nu}$ is distributional, concentrated on the axis. The present construction provides a natural regularisation: the photon is a normalizable mode of the electromagnetic field in double-null gauge, with a smooth transverse profile set by the wavelength $\sim c/\omega$. The electromagnetic stress-energy T_{uu} is well-defined throughout, integrates to $\hbar\omega$ over the transverse plane by energy conservation, and recovers the strict AS limit as $\omega \to \infty$ with the logarithmic divergence regulated at the scale c/ω.

Substituting into equation (34) gives

$$\boxed{\Delta A = 8\pi G\,\hbar\omega = 8\pi G\,\Delta E,} \tag{35}$$

and applying the Bekenstein–Hawking relation yields the corresponding entropy increment

$$\Delta S = \frac{\Delta A}{4\ell_P^2} = \frac{2\pi\,\Delta E}{\hbar\omega/\hbar} = 2\pi, \tag{36}$$

in natural units, consistent with one Planck-area patch per detection event. This derivation uses only the null energy condition and energy conservation on the horizon; it does not invoke spherical symmetry or the heuristic Unruh–Landauer argument of Section 2, and is therefore native to the Rindler geometry of the bilocal construction. The horizon area grows by one discrete, calculable increment per photon absorbed, microstate by microstate, without reference to bulk thermodynamic averages.

The longitudinal shift $\Delta v \propto -\ln r^2$ produced by each AS pp-wave at the Rindler horizon is not merely a coordinate displacement: it is a supertranslation in the sense of Bondi–Metzner–Sachs (BMS) symmetry. Hawking, Perry, and Strominger [7] showed explicitly that a black hole is supertranslated by the passage of an asymmetric shockwave, and the AS longitudinal shift is precisely the linearised supertranslation field on the horizon. Carlip [**?**] demonstrated that the near-horizon diffeomorphisms of a generic black hole are enhanced to a BMS_3 algebra whose central charge determines the Bekenstein–Hawking entropy, but his programme identifies the symmetry without specifying the physical process that excites individual modes. The bilocal construction supplies that process: each measurement event produces one AS pp-wave, which deposits one supertranslation at the

Rindler horizon, exciting one mode of the BMS_3 algebra. The entropy count is therefore not merely consistent with Carlip's result — the microstates being counted are the measurement events themselves.

5.4 Coarse-Graining and the Emergence of Thermality

In a steady-state scenario, $\mathcal{P}$ continuously emits entangled pairs, and $\mathcal{L}$ and $\mathcal{R}$ continuously detect them. Each detection is an AS pp-wave microstate with a definite, non-thermal description in the global Minkowski frame: a specific photon energy, a specific transverse position, a specific longitudinal shift. Coarse-graining over this ensemble recovers the Unruh thermal spectrum at temperature $T = a/2\pi$, consistent with the vacuum decomposition of equation (**??**). The thermodynamic description of the horizon emerges from this averaging; it is not a feature of the individual microstates.

This completes the microstate programme. Each entropy increment in the Bekenstein–Hawking area law corresponds to a single AS pp-wave interaction. Each pp-wave interaction corresponds to a single measurement event in the bilocal construction. Each measurement event both encodes a spatial bit and generates the spatial separation that bit refers to. The gravitational, thermodynamic, information-theoretic, and geometric descriptions of the horizon are unified at the level of individual microstates, and thermality, entropy, and space itself emerge together from the coarse-grained aggregate of these events.

5.5 The Microstate Action and the Gravitational Boundary Term

The microstate action $S_{\mathrm{micro}} = \hbar\omega$, identified with the electromagnetic stress-energy integrated over the null plane, has a precise relationship to the gravitational boundary action. On a stationary Rindler horizon the unperturbed expansion of the null generators vanishes, $\theta = 0$, so the Gibbons–Hawking–York boundary term [23] is sourced entirely by the AS pp-wave perturbations:

$$S_{GHY}\big|_{\mathcal{H}} = \frac{c^4}{8\pi G}\int_{\mathcal{H}} \theta\, d^2x_\perp\, du = \frac{c^4}{8\pi G}\sum_i \Delta A_i = \frac{c^4}{8\pi G}\sum_i 8\pi G\hbar\omega_i = \sum_i \hbar\omega_i = \sum_i S_{\mathrm{micro},i}. \tag{37}$$

The gravitational boundary action and the matter microstate action are therefore two accountings of the same underlying measurement events: one geometric, one operational. This identifies the GHY boundary term as the natural gravitational counterpart of the matter action at the microstate level, and is consistent with programmes of Jacobson [1] and Padmanabhan [24] arguing that the boundary term rather than the bulk Einstein–Hilbert action is the fundamental gravitational degree of freedom. The present construction realises that claim at the level of individual microstates, with each boundary contribution identified with a specific quantum measurement event.

The equality (37) admits an independent derivation from the matter substrate perspective of Section 4, which provides additional intuition. Each absorption sample deposits one quantum of action $\hbar$ into the detector's matter-memory and increments the detector mass by $\Delta m = \hbar\omega/c^2$. By the Schwarzschild relation $r_s = 2Gm/c^2$, this mass increment shifts the horizon radius by

$$\Delta r_s = \frac{2G\hbar\omega}{c^4}, \tag{38}$$

and the corresponding horizon area increment is

$$\Delta A = 8\pi r_s\, \Delta r_s = 8\pi G\hbar\omega, \tag{39}$$

in precise agreement with eq. (35), now derived entirely from the matter sample picture rather than from the Raychaudhuri equation. The two derivations approach the same

result from opposite directions: the geometric derivation proceeds top-down, from the AS pp-wave stress-energy through the null congruence to the area increment; the matter derivation proceeds bottom-up, from the individual sample through the mass increment to the same area. Their agreement is not a coincidence but a consistency condition on the framework: the geometric and operational descriptions are two accountings of the same microstate event, and $\Delta A = 8\pi G\hbar\omega$ is the relation that connects them. Summing over all absorption samples then gives eq. (37) directly, with each $\hbar\omega$ contribution to S_{GHY} corresponding to one quantum of action deposited into matter-memory.

6 Entanglement as Spatial Information

Each measurement produces a polarisation outcome[4], H or V, at each wedge, perfectly anticorrelated between $\mathcal{L}$ and $\mathcal{R}$. But consider a different question: not *what* was observed, but *where*. The question "who observed H?" has a binary answer: $\mathcal{L}$ (left wedge) or $\mathcal{R}$ (right wedge). This is a *spatial bit*, a bit that encodes not a polarisation state but a location in space.

Each measurement event has exactly two irreducible constituents: one quantum of action $\hbar$ written persistently into the detector's matter-memory as a series of samples, and one bit of polarisation registered as a classical outcome. The entanglement between $\mathcal{R}$ and $\mathcal{L}$ is consumed in producing these two records, an increase in sampling rate and one bit at each wedge. What remains after measurement is not a remnant of the entanglement but its classical residue: the perfect anticorrelation between the two bits, and the correlated pair of samples written into the matter on each side of the horizon. The spatial bit is precisely the polarisation bit now read as a location rather than a spin state: the same bit that records *what* was observed also records *where*, because in this construction the two wedges carry opposite helicities by design. This double duty is not merely convenient but forced: the anticorrelated helicity assignment of the bilocal source makes the two wedges distinguishable by construction, so the spatial bit is already latent in the source geometry before any measurement occurs.

The information is carried by the matter of the detectors, not by the horizon. As established in Section 4, the horizon is a boundary that makes the left/right distinction sharp; it is the matter on each side that accumulates the samples and holds the classical record. $\mathcal{R}$'s detector has absorbed one photon, increased its sampling rate, and registered helicity ± 1. $\mathcal{L}$'s detector has absorbed one photon, increased its sampling rate, and registered helicity ∓ 1. The two records are perfectly complementary by the anticorrelated structure of the bilocal source, and their complementarity is encoded in matter before it is reflected in geometry.

This reframing allows us to realize this entanglement-induced spatial separation concretely. As established in Section 5, the longitudinal shift of the AS pp-wave creates a new null interval between the two wedges; a region of spacetime that did not exist, in the sense of being causally accessible, before the photon passed. The bit of spatial information encoded in the left/right measurement outcome, "who observed H?", refers to precisely this separation: the measurement event does not merely *record* a spatial distinction, it *creates* the spatial separation that the bit describes. The two wedges are therefore connected not by the horizon as an abstract geometric object but by the physical correlations between the energy-momentum impulses deposited in the matter on each side.

The spatial bit is therefore not merely a record of which side the photon landed on. It is a label for which of the two correlated geometric deformations was realized, and the two labels together reconstruct the bridge. This is an operational realization of entanglement-induced spatial separation at the level of individual microstates: the entanglement between $\mathcal{L}$ and $\mathcal{R}$'s outcomes is not a stand-in for the bridge but is physically instantiated in the

[4] We are analyzing the case where $\mathcal{L}$ and $\mathcal{R}$ both use the $H/\ V$ basis.

pair of correlated samples whose geometric imprint *is* the bridge. The bridge is built from matter, sample by sample.

This is the sense in which the construction realizes ER=EPR at the level of individual microstates. Maldacena and Susskind's correspondence [4] identifies entangled pairs with Einstein–Rosen bridges at the level of bulk thermodynamics. Here the EPR side is the entangled photon pair emitted by $\mathcal{P}$, whose quantum correlations are consumed by the two measurement events. The ER side is the pair of correlated AS longitudinal shifts in opposite null directions, which together constitute a geometric bridge between the two wedges: a region of null separation that did not exist before the photon pair crossed the horizons. The spatial bit — the answer to "who observed H?" — is the classical record of which side of the bridge each measurement occurred on. The correspondence is not an analogy: the entanglement is converted, by measurement, into a persistent geometric imprint, and the bridge is the imprint. What is new here is that both sides of the correspondence are calculable at the single-microstate level, with a definite energy-momentum source and a definite geometric output, rather than established by a bulk thermodynamic argument.

The anticorrelated helicity structure gives the bridge an orientation absent in the scalar case: the two sides carry opposite handedness, and whether this constitutes a topological invariant of the transverse geometry is an open question we leave for future work.

7 Appendix: Consistency of the Double-Null Gauge Condition

We verify that the double-null gauge condition $A_+ = A_- = 0$ is consistent with Maxwell's equations and leaves exactly two physical degrees of freedom.

7.1 Coordinates and Notation

In double-null coordinates (x^+, x^-, x^1, x^2) the Minkowski metric reads

$$ds^2 = -2\,dx^+\,dx^- + (dx^1)^2 + (dx^2)^2, \tag{40}$$

so that $\partial_+ = \partial/\partial x^+$, $\partial_- = \partial/\partial x^-$, and the d'Alembertian is

$$\Box = -2\partial_+\partial_- + \partial_i\partial^i, \qquad i = 1, 2. \tag{41}$$

The gauge field has components $A_\mu = (A_+, A_-, A_1, A_2)$ and transforms as

$$A_\mu \to A_\mu + \partial_\mu \Lambda \tag{42}$$

under a gauge transformation with parameter $\Lambda(x^+, x^-, x^1, x^2)$.

7.2 Imposing the Gauge Conditions in Stages

The single gauge parameter Λ is a function of all four coordinates. We use it in three stages, each consuming a different functional dependence.

Stage 1: $A_+ = 0$. Requiring $A_+ + \partial_+\Lambda = 0$ determines Λ as a function of x^+:

$$\Lambda(x^+, x^-, x^i) = -\int^{x^+} A_+(s, x^-, x^i)\,ds + \Lambda_0(x^-, x^i), \tag{43}$$

where $\Lambda_0(x^-, x^i)$ is an arbitrary residual function independent of x^+.

Stage 2: $A_- = 0$. Under the residual freedom $\Lambda_0(x^-, x^i)$, the A_- component transforms as $A_- \to A_- + \partial_- \Lambda_0$. Setting this to zero determines Λ_0 as a function of x^-:

$$\Lambda_0(x^-, x^i) = -\int^{x^-} A_-(s, x^i)\, ds + \Lambda_1(x^i), \tag{44}$$

where $\Lambda_1(x^i)$ is a residual function of the transverse coordinates only.

Stage 3: Transverse Coulomb condition $\partial^i A_i = 0$. Under the remaining freedom $\Lambda_1(x^i)$, the transverse components transform as $A_i \to A_i + \partial_i \Lambda_1$. Imposing $\partial^i A_i = 0$ requires

$$\nabla_\perp^2 \Lambda_1 = -\partial^i A_i, \tag{45}$$

which is a Poisson equation in the transverse plane with a unique solution (up to harmonic functions, fixed by boundary conditions). This exhausts the gauge freedom entirely.

The three conditions have consumed the full gauge parameter Λ in stages, with no over-determination: each stage uses a strictly different functional dependence of Λ.

7.3 Physical Degrees of Freedom

After imposing $A_+ = A_- = 0$ and $\partial^i A_i = 0$, the remaining components are A_1 and A_2 subject to one differential constraint $\partial^i A_i = 0$. In momentum space this reads $k^i A_i = 0$.

For our specific construction, the photons propagate in the $\pm z$ direction, so their wave-vectors are purely null: $k_\perp^i = 0$ for $i = 1, 2$. The transverse Coulomb condition $k^i A_i = 0$ is therefore trivially satisfied for any A_1, A_2, and both transverse components are independent. The two circular polarisation vectors

$$\epsilon^i_{(+1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}, \qquad \epsilon^i_{(-1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix} \tag{46}$$

are the two independent physical degrees of freedom, corresponding to helicity $\lambda = \pm 1$. This is the correct count for a massless spin-1 field in four dimensions.

The trivial satisfaction of the transverse Coulomb condition relies on the photons propagating purely in the $\pm z$ direction with vanishing transverse momentum $k_\perp^i = 0$. For photons with nonzero transverse momentum the condition $k^i A_i = 0$ imposes a non-trivial constraint and the degree of freedom count requires revisiting.

7.4 Consistency with Maxwell's Equations

With $A_+ = A_- = 0$ and $\partial^i A_i = 0$, the non-zero field strength components are

$$F_{+i} = \partial_+ A_i, \qquad F_{-i} = \partial_- A_i, \qquad F_{ij} = \partial_i A_j - \partial_j A_i, \qquad F_{+-} = 0. \tag{47}$$

Maxwell's equations $\partial^\mu F_{\mu\nu} = J_\nu$ decompose as follows.

Transverse components $\nu = i$:

$$-2\partial_+ \partial_- A_i + \partial^j (\partial_j A_i - \partial_i A_j) = J_i. \tag{48}$$

Using $\partial^i A_i = 0$, the second term simplifies to $\partial^j \partial_j A_i$, giving the standard wave equation

$$\boxed{\Box A_i = J_i.} \tag{49}$$

Null components $\nu = \pm$:

$$-\partial^i \partial_\pm A_i = J_\pm. \tag{50}$$

These are constraint equations relating the null source components to the transverse field. For our bilocal source, which is purely transverse, $J_+ = J_- = 0$, and the constraints reduce to

$$\partial^i \partial_\pm A_i = 0. \tag{51}$$

Differentiating the transverse Coulomb condition $\partial^i A_i = 0$ with respect to $x^\pm$ gives $\partial^i(\partial_\pm A_i) = 0$, so these constraints are automatically satisfied. ✓

7.5 Summary

The double-null gauge condition $A_+ = A_- = 0$ supplemented by the transverse Coulomb condition $\partial^i A_i = 0$:

1. uses the gauge freedom of Λ completely and without over-determination, via a three-stage procedure consuming the x^+-, x^--, and x^i-dependent parts of Λ in sequence;
2. leaves exactly two physical degrees of freedom, the helicity eigenstates $\lambda = \pm 1$, consistent with the known count for a massless spin-1 field;
3. reduces Maxwell's equations to the standard transverse wave equation $\Box A_i = J_i$, with the null constraint equations automatically satisfied for a purely transverse source.

The gauge is therefore internally consistent and does not over-constrain the physical degrees of freedom. The $\mathbb{Z}_2$ wedge-swap symmetry is preserved throughout: conditions on A_+ and A_- are imposed symmetrically, and neither null direction is privileged over the other.

8 Conclusion

9 Acknowledgements

I'm also grateful to Claude and ChatGPT for helpful discussions.

References

[1] T. Jacobson, "Thermodynamics of spacetime: the einstein equation of state," *Physical review letters*, vol. 75, no. 7, p. 1260, 1995.

[2] E. Verlinde, "On the origin of gravity and the laws of newton," *Journal of High Energy Physics*, vol. 2011, no. 4, pp. 1–27, 2011.

[3] M. Van Raamsdonk, "Building up space–time with quantum entanglement," *International Journal of Modern Physics D*, vol. 19, no. 14, pp. 2429–2435, 2010.

[4] J. Maldacena and L. Susskind, "Cool horizons for entangled black holes," *Fortschritte der Physik*, vol. 61, no. 9, pp. 781–811, 2013.

[5] T. Dray and G. Hooft, "The gravitational shock wave of a massless particle," *Nuclear physics B*, vol. 253, pp. 173–188, 1985.

[6] A. Strominger and C. Vafa, "Microscopic origin of the bekenstein-hawking entropy," *Physics Letters B*, vol. 379, no. 1-4, pp. 99–104, 1996.

[7] S. W. Hawking, M. J. Perry, and A. Strominger, "Soft hair on black holes," *Physical Review Letters*, vol. 116, no. 23, p. 231301, 2016.

[8] R. Penrose, "On gravity's role in quantum state reduction," *General relativity and gravitation*, vol. 28, no. 5, pp. 581–600, 1996.

[9] L. Diosi, "A universal master equation for the gravitational violation of quantum mechanics," *Physics letters A*, vol. 120, no. 8, pp. 377–381, 1987.

[10] K. Player, "Driving the unruh response," *arXiv preprint arXiv:2509.16710*, 2025.

[11] C. Rovelli, "Relational quantum mechanics," *International journal of theoretical physics*, vol. 35, no. 8, pp. 1637–1678, 1996.

[12] C. A. Fuchs, N. D. Mermin, and R. Schack, "An introduction to qbism with an application to the locality of quantum mechanics," *American Journal of Physics*, vol. 82, no. 8, pp. 749–754, 2014.

[13] W. G. Unruh, "Notes on black-hole evaporation," *Physical Review D*, vol. 14, no. 4, p. 870, 1976.

[14] E. P. Wigner, "Remarks on the mind-body question," in *Philosophical reflections and syntheses*, pp. 247–260, Springer, 1995.

[15] J. A. Wheeler, "Information, physics, quantum: The search for links," *Feynman and computation*, pp. 309–336, 2018.

[16] R. Landauer, "Irreversibility and heat generation in the computing process," *IBM journal of research and development*, vol. 5, no. 3, pp. 183–191, 1961.

[17] J. D. Bekenstein, "Black holes and entropy," *Phys. Rev. D*, vol. 7, pp. 2333–2346, Apr 1973.

[18] C. C. Gerry and P. L. Knight, *Introductory quantum optics*. Cambridge university press, 2023.

[19] N. D. Birrell and P. C. W. Davies, "Quantum fields in curved space," 1984.

[20] H. J. Bremermann, "Optimization through evolution and recombination.," *Self-organizing systems*, pp. 93–106, 1962.

[21] L. H. Ryder, *Quantum field theory*. Cambridge university press, 1996.

[22] R. Penrose, *The road to reality*. Random house, 2006.

[23] G. W. Gibbons and S. W. Hawking, "Action integrals and partition functions in quantum gravity," *Physical Review D*, vol. 15, no. 10, p. 2752, 1977.

[24] T. Padmanabhan, "Thermodynamical aspects of gravity: new insights," *Reports on Progress in Physics*, vol. 73, no. 4, p. 046901, 2010.