Measurement Events as Gravitational Microstates

Kevin Player*

April 18, 2026

Abstract

Thermodynamic approaches to gravity, from Jacobson to Verlinde, relate spacetime dynamics to horizon entropy but do not specify the microscopic carriers of entropy increments. We take as primitive that quantum measurement necessarily involves a physical interaction with energy–momentum transfer between system and detector. Within this setting, we model such interactions as localized Aichelburg–Sexl (AS) pp-wave perturbations of spacetime geometry. Each measurement event then induces a discrete shift in null horizon coordinates, contributing additively to horizon area, and coarse-graining over the resulting ensemble reproduces the Bekenstein–Hawking area law. A bilocal realization using entangled photon pairs yields correlated AS shifts on opposite Rindler horizons, providing an operational realization of entanglement as spatial connectivity in the spirit of ER=EPR. Within this construction, the gravitational boundary term on the horizon is identified with the accumulated energy transfer associated with measurement events, linking matter dynamics and horizon entropy at the level of individual microstates.

1 Introduction

Two landmark results establish that gravity and Unruh thermodynamics are deeply intertwined. Jacobson showed that the Einstein equations can be derived from thermodynamic identities applied to local Rindler horizons, with horizon entropy playing the role of the fundamental quantity from which spacetime dynamics emerge [1]. Verlinde extended this perspective, arguing that gravitational forces arise from entropy gradients on holographic screens, recasting gravity as an entropic rather than a fundamental interaction [2]. Both frameworks, however, operate at the level of bulk thermodynamics and leave open a prior question: what physical process underlies each individual entropy increment?

We argue that the answer is quantum measurement. Each act of information acquisition by a physical observer constitutes a discrete microstate event, and the collective backreaction of such events, together with the causal correlations they induce, gives rise to gravitational effects. The argument proceeds in three steps. First, any observation is mediated by a physical interaction: information transfer necessarily involves an exchange of energy and momentum between the observed system and the detector, a discrete impulse. Second, by the equivalence principle, this impulse is locally indistinguishable in that from the effect of a gravitational field: no local measurement can distinguish the recoil associated with a quantum measurement event from acceleration in a gravitational potential. Third, when such impulses are modelled geometrically, via Aichelburg–Sexl (AS) pp-wave solutions, as we explore, they imprint energy and information directly onto horizon geometry, and coarse-graining over the resulting ensemble recovers standard horizon thermodynamics.

* *kjplaye@gmail.com*

Modeling the spacetime response to such localized energy–momentum transfer events using Aichelburg–Sexl (AS) pp-wave geometries follows the gravitational shockwave framework of Dray, 't Hooft [3], and others. Their construction uses AS solutions sourced by infalling classical particles to define horizon scattering amplitudes; the key difference here is that the source is a quantum entangled pair emitted from a common ancestor $\mathcal{P}$, and the information acquisition framing connects the geometric shift directly to a classical bit acquired by an observer.

What distinguishes the present construction from all of these approaches is the combination of three features that do not appear together elsewhere in the literature: a non-thermal bilocal source that selects individual microstates from the Unruh ensemble rather than assuming thermality as a primitive; the identification of each microstate with a specific quantum measurement event carrying a calculable energy-momentum transfer; and an operational realization of entanglement-induced spatial separation at the single-microstate level with a concrete geometric imprint. Each of these features appears in isolation in various forms across the literature, but their combination, and in particular the grounding of horizon geometry in individual measurement events rather than bulk thermodynamic averages, appears to be new.

1.1 Conceptual Roadmap

This stands in sharp contrast to the Diósi–Penrose objective collapse programme [4, 5], which postulates that gravity induces an objective, observer-independent collapse of the quantum state. In that framework, gravitational self-interaction is taken to define a physical timescale for wavefunction reduction, leading to a dynamical modification of standard quantum mechanics. The present work does not adopt objective collapse or any modification of unitary evolution. Instead, we treat quantum measurement as a physical interaction mediated by energy–momentum transfer, and interpret any apparent reduction of the state as an effective description arising from such processes. The contrast between the two approaches is illustrated in Figure 1.



Figure 1: Solid arrows denote "substrate" relations: $A \to B$ means every instance of B is physically realized by an instance of A, for a given observer. The dashed arrow denotes the Diósi–Penrose postulate, where the relation is claimed to hold objectively rather than observer-relatively.

To make this concrete, we build on the microstate construction of [6], which realises individual terms of the Unruh thermal ensemble via bilocal driving sources. The present paper applies and extends that construction. Entangled pairs emitted from a localized source propagate into left and right Rindler wedges; polarization measurement events are associated with discrete energy–momentum transfer and yield correlated classical outcomes. Each such microstate carries a calculable gravitational imprint, modelled by an AS pp-wave solution, and coarse-graining over the ensemble reproduces standard horizon thermodynamics. This construction also provides an operational realization of entanglement-induced spatial separation at the level of individual microstates: entanglement is mapped

into geometric correlations via the associated gravitational backreaction, and each measurement event contributes a discrete, calculable increment to horizon entropy through its associated energy–momentum transfer and gravitational backreaction.

2 Motivating Heuristics

Landauer's principle associates a minimal energy cost with the acquisition of information [7]. For an accelerated observer, the relevant temperature is the Unruh temperature [8]. Substituting one into the other eliminates temperature entirely, yielding a relation between information acquisition, energy transfer, and acceleration (a version of the BHT first law, equation (3)). This is not merely an algebraic convenience: it is a concrete dictionary between the thermal description natural to a Rindler observer and the non-thermal global Minkowski description, with the apparent thermality of the vacuum emerging as a consequence of the observer's acceleration rather than as a fundamental property of the state. In this way we derive a sequence of standard results in black hole thermodynamics as a consequence of Newton's laws.

2.1 The Unruh-Landauer Relation

Landauer's principle suggests that the acquisition of ΔH bits of information by a physical observer requires a minimum energy transfer

$$\Delta E \geq k_B T \, \Delta H. \tag{1}$$

We note that Landauer's principle is conventionally stated for information erasure rather than acquisition. Here we apply it to acquisition, which is the physically relevant process for an observer registering a measurement outcome. The bound has the same form in both cases; we apply it to acquisition on the grounds that registering a measurement outcome is physically equivalent to a bit being written into the detector's state, which is the process Landauer's principle governs.

For an observer undergoing proper acceleration a, the Unruh effect associates a temperature

$$T = \frac{\hbar a}{2\pi k_B c} \tag{2}$$

with the local vacuum. Substituting into Landauer's bound eliminates $k_B T$, yielding

$$\boxed{\Delta E \geq \frac{\hbar a}{2\pi c} \, \Delta H.} \tag{3}$$

Temperature has dropped out entirely: equation (3) is a temperature-independent relation between energy, acceleration, and information, suitable for the Minkowski vacuum. Acceleration sets the energy scale of information.

The Unruh–Landauer bound is saturated at one bit when the photon wavelength equals the horizon distance up to a factor of $2\pi/\ln 2$:

$$\lambda_{\text{sat}} = \frac{2\pi}{\ln 2} \cdot \frac{c^2}{a} = \frac{2\pi}{\ln 2} \cdot d_{\mathcal{H}}, \tag{4}$$

where $d_{\mathcal{H}} = c^2/a$ is the proper distance to the Rindler horizon. The saturation photon therefore fits within the causal diamond of the measurement event: its wavelength is commensurate with the horizon scale, too short to be super-horizon and too long to deposit more than one bit. This identifies the horizon distance as the natural length scale of a single-bit measurement event, and the Rindler horizon as the surface at which the Landauer bound is geometrically realized.

2.2 Momentum Transfer and Acceleration

In what follows, standard results emerge as consequences of this single relation and Newton's laws. The energy transfer ΔE implied by equation (3) is carried by a physical quantum — matter or radiation — mediating the measurement interaction. We refer to this as the *thrust quantum* of effective mass m_t. (This will actually be a pure energy carrier in our later microstate example.) For a relativistic carrier, the associated momentum transfer is

$$\Delta p = \frac{\Delta E}{c}. \tag{5}$$

A measurement event localized to a spatial scale r is bounded by a causal diamond: the intersection of the future light cone of its initiation and the past light cone of its completion. The temporal extent of this diamond is $\Delta t = 2r/c$, fixed entirely by the causal geometry of the event rather than by any dynamical assumption. We identify the interaction time with this natural causal duration, so that

$$a = \frac{\Delta p}{m_d \Delta t} = \frac{\Delta E}{2 m_d r}. \tag{6}$$

This perspective is closely related in spirit to Erik Verlinde's entropic gravity [2], where forces arise from entropy gradients, but differs in that the present construction grounds each entropy increment in a discrete measurement event with an associated energy-momentum transfer, rather than in a coarse-grained thermodynamic gradient.

2.3 The Schwarzschild Radius

We match the acceleration inferred from the measurement-induced impulse to the Newtonian gravitational field in order to identify the characteristic length scale associated with the detector. The thrust quantum carries energy ΔE and produces an acceleration a on the detector. Identifying this with the Newtonian gravitational acceleration due to the mass m_t at distance r.

$$a = \frac{G m_t}{r^2}, \tag{7}$$

and substituting equation (6) with $\Delta E = m_t c^2$, we obtain

$$\frac{m_t c^2}{2 m_d r} = \frac{G m_t}{r^2}. \tag{8}$$

Solving for r:

$$r = \frac{2 G m_d}{c^2} \equiv r_s, \tag{9}$$

which is precisely the Schwarzschild radius of the detector mass m_d.

It is notable that the resulting length scale is set by the detector mass m_d, rather than the source mass m_t. This reflects the fact that the construction is inherently observer-centred: the acceleration entering the argument is the response of the detector to the measurement-induced impulse, and the associated scale is therefore the horizon radius relevant to that observer. In this sense, the quantity r_s should be interpreted not as the Schwarzschild radius of the source, but as the characteristic horizon scale associated with the detector as a physical system undergoing measurement. This is consistent with the observer-dependence of horizon structure in relativistic settings, as exemplified by the Unruh effect, where acceleration alone gives rise to a causal horizon.

2.4 Surface Gravity

Evaluating the acceleration at $r = r_s$ gives the surface gravity of the detector:

$$\kappa = \frac{G m_d}{r_s^2} = \frac{c^4}{4 G m_d}, \tag{10}$$

in agreement with the standard Schwarzschild result. This is the acceleration that appears in the Unruh temperature associated with the detector's horizon, $T = \hbar\kappa/2\pi k_B c$, providing an internal consistency check on the framework.

2.5 Bekenstein-Hawking Entropy

Returning to the Unruh-Landauer relation (3) and substituting the surface gravity κ for a, the entropy increment associated with an energy increment $\Delta E = c^2 \Delta m_d$ is

$$\Delta H \leq \frac{2\pi c}{\hbar\kappa}\Delta E = \frac{8\pi G m_d}{\hbar c^3} \cdot c^2 \Delta m_d. \tag{11}$$

Expressing this in terms of the change in Schwarzschild area $\Delta A = 32\pi G^2 m_d \Delta m_d / c^4$:

$$\Delta H \leq \frac{\Delta A}{4\ell_P^2}, \qquad \ell_P^2 = \frac{\hbar G}{c^3}, \tag{12}$$

which is the Bekenstein-Hawking area law. Each measurement event contributes a discrete entropy increment proportional to the area imprinted on the horizon by the associated energy-momentum transfer.

This derivation parallels Jacob Bekenstein's original argument [9], in which the entropy increase of a black hole is inferred from the minimal energy required to localize and drop a system across the horizon; here, the same structure emerges from the minimal energy cost of registering information at the horizon scale.

The two bounds are saturated simultaneously when $a = c^2/R$ — which is exactly the condition for being *at a horizon*. The horizon is therefore not just where thermodynamics happens to work out, it is the unique locus where the minimum cost of information acquisition equals the maximum information content of the region.

2.6 Summary

Beginning from the single relation (3), and using only Newton's laws and the equivalence principle, we have recovered the Schwarzschild radius, the surface gravity, and the Bekenstein-Hawking area law as consequences of the observation $\rightarrow$ acceleration identification. These are standard results, but their derivation here from the physical cost of information acquisition rather than from thermodynamic postulates is the point.

The thrust quantum introduced in this section, the physical carrier of the measurement-induced impulse, will be given a explicit microstate realization in the sections that follow. It is identified with one of a pair of entangled photons modeled as Aichelburg-Sexl pp-waves crossing past Rindler horizons in the Unruh picture. In this case the back reaction of a single force carrier in one wedge is correlated with an equal and opposite reaction in the other from a common source.

3 Bilocal Microstate Construction

Let $\hbar = c = 1$.

The heuristic arguments of Section 2 establish a relationship between information acquisition, acceleration, and gravitational effects, but leave open the question of what a single microstate looks like concretely. In standard treatments, the Unruh effect is derived by coupling a detector to a pre-existing acceleration and reading off a thermal response from the vacuum. Following [6], we invert this logic: we place an interaction in the past, where a source injects non-thermal entangled excitations into the field, and show that the observer's accelerated response can be driven by these source-induced excitations. The thermal ensemble then emerges as a coarse-grained aggregate of such microstates, rather than being a primitive property of the vacuum.

Each microstate, equation (23) upcomming, is individually non-thermal: it carries a definite helicity pair and a definite Rindler frequency Ω, neither of which is fixed by the Unruh temperature. Thermality is not a property of any single microstate but of the ensemble; it emerges when one asks only which wedge a detection occurred in, discarding the specific microstate identity, and is a consequence of coarse-graining rather than a premise of the construction.

We now upgrade this construction from a spin-0 scalar field in the original source [6] to the spin-1 electromagnetic field. This is not merely a technical generalisation. As we show, the photon carries a unit of action $\hbar$ per cycle alongside its energy $E = \hbar\omega$, which closes the Landauer gap left open in Section 2: in each wedge, the minimum actionable event in the construction is precisely a single photon, and its polarization state constitutes the classical bit acquired by each observer from detection. The anticorrelated helicities of the entangled pair are the natural completion of the $\mathbb{Z}_2$ swap symmetry between wedges, and as we discuss in Section 4, they imprint a definite geometric signature onto the horizon.

3.1 A Source and Three Observers

Three observers are placed at spacetime positions $(t, z) = (0, 1)$, $(0, -1)$, and $(-1, 0)$, named $\mathcal{R}$ - *Right*, $\mathcal{L}$ - *Left*, and $\mathcal{P}$ - *Past* respectively. $\mathcal{P}$ serves as the common causal ancestor of $\mathcal{R}$ and $\mathcal{L}$ (see Figure 2).



Figure 2: A source in the past, $\mathcal{P}$, emits an entangled photon pair that accelerates $\mathcal{R}$ into the right Rindler wedge and $\mathcal{L}$ into the left Rindler wedge. The two null directions of propagation define the z^+ and z^- axes of the double-null coordinate system.

$\mathcal{P}$ emits an entangled pair of spin-1 massless photons with momenta $\pm k$ and Rindler frequency $\omega = \omega_k$. The momenta are perfectly balanced, so the source exhibits no net back-reaction in this minimal model. The two photons propagate into the left and right Rindler wedges, where $\mathcal{L}$ and $\mathcal{R}$ detect them respectively, each registering a helicity eigenvalue and acquiring one classical bit of information.

The three-party structure $\mathcal{P}$–$\mathcal{R}$–$\mathcal{L}$ is a tripartite entanglement: neither $\mathcal{R}$ nor $\mathcal{L}$ alone holds sufficient information to reconstruct the emission event at $\mathcal{P}$, but their two complementary bits together do, making $\mathcal{P}$ the logical AND of the pair and providing a natural microstate-level realization of interior reconstruction.

3.2 Double-Null Gauge

The emission event naturally defines two null directions: the right-moving photon has wave-vector k_R^μ and the left-moving photon has wave-vector $k_L^\mu = -k_R^\mu$. These two null directions are precisely the z^+ and z^- axes of double-null coordinates

$$z^\pm = \frac{1}{\sqrt{2}}(t \pm z), \tag{13}$$

with transverse coordinates $\mathbf{z}_\perp = (z^1, z^2)$. Working in this frame we impose the symmetric double-null gauge condition

$$A_+(z) = 0, \qquad A_-(z) = 0, \tag{14}$$

so that the gauge field is supported entirely on the transverse components $\mathbf{A}_\perp = (A^1, A^2)$. This gauge respects the $\mathbb{Z}_2$ symmetry that swaps the two wedges ($z^+ \leftrightarrow z^-$) simultaneously with ($R \leftrightarrow L$), and therefore treats both observers symmetrically by construction. The residual gauge freedom is fixed by requiring $\partial_i A^i = 0$ in the transverse plane, leaving precisely the two physical helicity degrees of freedom $\epsilon^i_{(\lambda)}$ with $\lambda = \pm 1$; consistency with Maxwell's equations and the absence of over-determination are verified in Appendix 6.

The lack of manifest four-dimensional covariance in equation (14) is a consequence of aligning the gauge with the pair of physical null directions selected by the source, rather than an arbitrary choice. The full Lorentz group is broken to the little group preserving the pair $\{k^\mu_R, k^\mu_L\}$, which is the two-dimensional Euclidean group $ISO(2)$ acting on the transverse plane. This is precisely the little group of massless particles, and the two helicity eigenstates $\lambda = \pm 1$ are its irreducible representations. All physical results are independent of the gauge choice.

3.3 The Bilocal Source

In double-null gauge the electromagnetic field is characterised by its transverse components alone. The bilocal driving source generalises the scalar construction to

$$J^{ij}(x, y) = \lambda\, \epsilon^{i*}_{(-1)}\, \epsilon^j_{(+1)}\, f_L(x)\, f_R(y), \tag{15}$$

where λ is a coupling constant, f_L and f_R are normalised mode profiles supported on the left and right wedges respectively, and $\epsilon^i_{(\lambda)}$ are the circular polarization vectors

$$\epsilon^i_{(+1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}, \qquad \epsilon^i_{(-1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}. \tag{16}$$

The index structure of (15) assigns helicity $+1$ to the right-moving mode and helicity -1 to the left-moving mode: the pair is *anticorrelated* in helicity. This is the unique assignment consistent with angular momentum conservation at the emission vertex and with the $\mathbb{Z}_2$ wedge-swap symmetry, under which $\epsilon^i_{(+1)} \leftrightarrow \epsilon^i_{(-1)}$.

The sourced Lagrangian reads

$$\begin{aligned} \mathcal{L}_{\text{sourced}} &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \int J^{ij}(x, y)\; A^+_i(x)\, A^+_j(y)\; d^4x\, d^4y \;+\; \text{h.c.} \\ &= \mathcal{L}_{\text{free}} + \tfrac{1}{2}\lambda \Big(a^{L\dagger}_{-k,(-1)}\, a^{R\dagger}_{k,(+1)} \;+\; a^L_{-k,(-1)}\, a^R_{k,(+1)} \Big), \end{aligned} \tag{17}$$

where $a^{L\dagger}_{-k,(-1)}$ and $a^{R\dagger}_{k,(+1)}$ are creation operators for left-wedge helicity -1 and right-wedge helicity $+1$ modes respectively. The interaction term in (17) is a two-mode squeezing interaction at frequency ω, now resolved by helicity: it is the direct vector-field analogue of the scalar squeezing interaction [10], with the anticorrelated polarization structure made explicit. The photon pair carries total angular momentum zero, consistent with emission from a scalar source at $\mathcal{P}$.

Each photon carries energy $E = \hbar\omega$ and, as a mode of the electromagnetic field, constitutes precisely one quantum of action $\hbar$ per oscillation cycle. The thrust quantum of Section 2 is thereby identified with a single photon: the minimum actionable event is the emission and detection of one such quantum.

The Unruh–Landauer bound is saturated not at the level of an individual photon but at the level of the source rate. The acceleration a does not fix the frequency ω of each emitted quantum; rather, it determines the rate $\dot{N}$ at which the bilocal source must

produce entangled pairs in order to fully[1] implement that acceleration, with the total power $\dot{N}\hbar\omega$ matching the Unruh thermal flux. Each individual photon carries exactly one quantum of action $\hbar$ and one binary helicity outcome, one bit in the sense of one binary alternative, carrying $\ln 2$ nats, and these are paired by the construction independently of a. The Landauer bound then operates as a consistency condition on the ensemble: the minimum energy cost per bit acquired, summed over the rate of acquisition, equals the power required to sustain the acceleration.

3.4 Polarization Bogoliubov Transformation

We now show that the polarization labels survive the Rindler–Unruh mode decomposition, so that the thermal ensemble retains its helicity structure and the anticorrelation of the emitted pair descends unambiguously to the level of individual Rindler microstates.

In the transverse plane, the electromagnetic field in double-null gauge decomposes into two independent scalar-like sectors labelled by helicity $\lambda = \pm 1$. For each sector, the right-wedge field expands as

$$A^i_\lambda(z) = \int_0^\infty \frac{d\Omega}{2\pi} \frac{1}{\sqrt{2\Omega}} \epsilon^i_{(\lambda)} \left[b^R_{\Omega,\lambda}\, u^R_\Omega(z) + b^{R\dagger}_{\Omega,\lambda}\, u^{R*}_\Omega(z) \right], \tag{18}$$

where $u^R_\Omega(z)$ are the right-wedge Rindler modes at Rindler frequency Ω, and $b^R_{\Omega,\lambda}$, $b^{R\dagger}_{\Omega,\lambda}$ are the corresponding annihilation and creation operators satisfying

$$\left[b^R_{\Omega,\lambda},\, b^{R\dagger}_{\Omega',\lambda'} \right] = \delta(\Omega - \Omega')\, \delta_{\lambda\lambda'}. \tag{19}$$

An identical expansion holds in the left wedge with $R \to L$ and $\lambda \to -\lambda$, reflecting the anticorrelated helicity assignment. Because each helicity sector is independently a scalar-like sector, the Bogoliubov transformation relating Minkowski and Rindler modes takes the same form in each sector. For helicity λ at Rindler frequency Ω, the Unruh modes are

$$c_{\Omega,\lambda} = \cosh\theta_\Omega\, b^R_{\Omega,\lambda} - \sinh\theta_\Omega\, b^{L\dagger}_{\Omega,-\lambda}, \tag{20}$$

where $\tanh\theta_\Omega = e^{-\pi\Omega c/a}$ encodes the Unruh distribution at acceleration a, and $-\lambda$ on the left-wedge operator reflects the anticorrelation. The transformation (20) is the standard Bogoliubov relation for a massless field [8, 11], now carrying a helicity label throughout. The Minkowski vacuum in the Rindler basis reads, for each helicity sector,

$$|0\rangle^{(\lambda)}_M = \frac{1}{\cosh\theta_\Omega} \sum_{n=0}^{\infty} \tanh^n\theta_\Omega\, |n_{\Omega,\lambda}\rangle_R\, |n_{\Omega,-\lambda}\rangle_L, \tag{21}$$

and the full vacuum is the product over all Ω and both helicities:

$$|0\rangle_M = \prod_{\Omega,\lambda} |0\rangle^{(\lambda)}_M. \tag{22}$$

Equation (21) makes explicit that the Minkowski vacuum is a sum over entangled helicity-anticorrelated pairs at each Rindler frequency. The bilocal source selects a single term from this sum at fixed Ω and fixed helicity pair $(\lambda, -\lambda) = (+1, -1)$, yielding the single-microstate

$$|\Psi_\Omega\rangle_M = |1_{\Omega,+1}\rangle_R\, |1_{\Omega,-1}\rangle_L + \mathcal{O}(\text{higher order}). \tag{23}$$

$\mathcal{R}$'s detection of helicity $+1$ and $\mathcal{L}$'s detection of helicity -1 are therefore perfectly anticorrelated: each measurement event yields one classical bit, and the two bits are the complement of one another.

[1] The bilocal source may be only part of the acceleration, so $\dot{N}$ is actually a lower bound.

The key structural result is that the helicity label λ is conserved under the Bogoliubov transformation: it enters equation (20) as a spectator index and exits unchanged on the Unruh operators $c_{\Omega,\lambda}$. Thermality, when it arises from coarse-graining over the ensemble of microstates (23), is therefore helicity-resolved: each helicity sector thermalises independently, and the anticorrelation between wedges is preserved at every level of the construction.

3.5 Summary

The passage from the scalar to the vector field construction introduces three structural improvements. First, the photon carrying $E = \hbar\omega$ and one quantum of action per cycle identifies the thrust quantum of Section 2 with a concrete physical object, closing the Landauer acquisition gap. Second, the double-null gauge condition, motivated by the pair of null directions defined by the emission event, treats both Rindler wedges symmetrically and exposes the $\mathbb{Z}_2$ swap symmetry as the organising principle of the construction. Third, the anticorrelated helicity structure, conserved through the Bogoliubov transformation, provides each observer with one classical bit whose values are perfectly complementary: the two bits together constitute the correlated classical record that, in the AS pp-wave picture of Section 4, is imprinted as geometric information onto the horizon.

4 pp-Wave Geometry

WLOG we introduce coordinates for the photon traveling from $\mathcal{P}$ $(-1, 0)$ to $\mathcal{R}$ $(0, 1)$. Let $u = t - z + 1$ and $v = t + z + 1$, where the photon propagates along the null plane $u = 0$, z is the longitudinal direction, and (x, y) are transverse coordinates. Then $(u, v) = (0, 0)$ is $\mathcal{P}$'s location, $(u, v) = (0, 2)$ is $\mathcal{R}$'s location, and $(u, v) = (0, 1)$ is the intersection of the photon with the past right Rinder horizon. The situtation is reversed for the left moving photon, we swap u and v to describe photon moving from $\mathcal{P}$ to $\mathcal{L}$.

4.1 The AS pp-Wave and Its Effects

Test particles at fixed transverse positions (x, y) experience two effects as the shock front crosses $u = 0$:

1. **Longitudinal shift.** The v coordinate is displaced by

$$\Delta v \propto -\ln r^2, \qquad r^2 = x^2 + y^2. \tag{24}$$

   This shift is lightlike: no proper time elapses for a particle crossing the shock, but it is displaced along the null direction. Near the axis ($r \to 0$) the shift diverges, encoding the concentration of energy along the photon's trajectory.

2. **Transverse focusing.** The gradient of the longitudinal shift produces a transverse velocity

$$\Delta \mathbf{v}_\perp \propto \nabla \ln r \propto \frac{1}{r}, \tag{25}$$

   directed toward the axis. Test particles are pulled toward the centre of the wave, regardless of their initial transverse separation.

4.2 The Longitudinal Shift Makes Space

The longitudinal shift deserves closer attention, because it carries a meaning beyond a mere coordinate displacement. The photon emitted by $\mathcal{P}$ propagates along the null direction; as it crosses the Rindler horizon, the AS shift displaces the future light cone of every test particle it passes. For the photon traveling into the right wedge, the shift

pushes the right wedge's geometry in the $+v$ direction; for the entangled partner traveling into the left wedge, the shift pushes the left wedge in the $-v$ direction. The two wedges are displaced *away from each other* along the longitudinal null direction. See Figure 3.



Figure 3: The green arrows show the longitudinal shift that occurs where the photons cross the horizon. Test particles, shown in yellow, are pushed outwards away from $z = 0$ due to the creation of space along the null lines originating from $\mathcal{P}$.

At the intersection of the shock front $u = 0$ with the past right Rindler horizon at $(u, v) = (0, 1)$, the transverse (x, y) plane is identified with the horizon area elements $dA = dx\,dy$. The longitudinal shift $\Delta v \propto -\ln r^2$ displaces the horizon outward in the v direction, pushing the right wedge geometry away from the horizon. The entangled partner photon produces a symmetric shift in the u direction at the left Rindler horizon, pushing the left wedge away from its horizon. The two wedges are therefore displaced away from their respective horizons in opposite null directions.

4.3 Transverse Focusing Increments Horizon Area

The transverse focusing of the AS pp-wave produces a calculable increment in horizon area. We derive this using the Raychaudhuri equation on the null congruence generating the Rindler horizon, following Jacobson [1], but applied here to a single microstate rather than a thermodynamic average.

For a null congruence with generators k^μ and expansion θ, the Raychaudhuri equation gives

$$\frac{d\theta}{du} = -\tfrac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} - 8\pi G\, T_{\mu\nu}k^\mu k^\nu. \tag{26}$$

Integrating over the horizon patch in the linearised regime and using $k^\mu = \delta^\mu_u$, the area change is

$$\Delta A = 8\pi G \int T_{uu}\, du\, d^2x_\perp. \tag{27}$$

In a strict AS construction, the source is a point particle and $T_{\mu\nu}$ is distributional, concentrated on the axis. The present construction provides a natural regularization: the photon is a normalizable mode of the electromagnetic field in double-null gauge, with a smooth transverse profile set by the wavelength $\sim c/\omega$. The electromagnetic stress-energy T_{uu} is well-defined throughout, integrates to $\hbar\omega$ over the transverse plane by energy conservation, and recovers the strict AS limit as $\omega \to \infty$ with the logarithmic divergence regulated at the scale c/ω.

Substituting into equation (27) gives

$$\boxed{\Delta A = 8\pi G\,\hbar\omega = 8\pi G\,\Delta E.} \tag{28}$$

The horizon area grows by one discrete, calculable increment per photon absorbed, microstate by microstate, without reference to bulk thermodynamic averages.

Applying the Bekenstein–Hawking relation as a heuristic, valid for Schwarzschild but taken here as an ansatz for the Rindler case, yields a corresponding entropy increment

$$\Delta S = \frac{\Delta A}{4\ell_P^2} = \frac{2\pi\,\Delta E}{\hbar\omega/\hbar} = 2\pi, \tag{29}$$

in natural units, consistent with one Planck-area patch per detection event.

The longitudinal shift $\Delta v \propto -\ln r^2$ produced by each AS pp-wave at the Rindler horizon is not merely a coordinate displacement: it is a supertranslation in the sense of Bondi–Metzner–Sachs (BMS) symmetry. Hawking, Perry, and Strominger [12] showed explicitly that a black hole is supertranslated by the passage of an asymmetric shockwave, and the AS longitudinal shift is precisely the linearised supertranslation field on the horizon. Carlip [**?**] demonstrated that the near-horizon diffeomorphisms of a generic black hole are enhanced to a BMS_3 algebra whose central charge determines the Bekenstein–Hawking entropy, but his programme identifies the symmetry without specifying the physical process that excites individual modes. The bilocal construction supplies that process: each measurement event produces one AS pp-wave, which deposits one supertranslation at the Rindler horizon, exciting one mode of the BMS_3 algebra. The entropy count is therefore not merely consistent with Carlip's result, the microstates being counted are the measurement events themselves.

4.4 Action and the Gravitational Boundary

Each absorption sample deposits one quantum of action $\hbar$ into the detector's matter-memory. The total action accumulated across all measurement events is therefore

$$S_{\text{matter}} = \sum_i \hbar\omega_i, \tag{30}$$

where the sum is over all absorption samples, each contributing energy $\hbar\omega_i$ to the detector mass. This is the matter action counted sample by sample: one quantum of action per event, one event per emitted pair.

The same events have a geometric description. By eq. (28), each absorption sample contributes a horizon area increment $\Delta A_i = 8\pi G\hbar\omega_i$, so the total area increment summed over all events is

$$\sum_i \Delta A_i = 8\pi G \sum_i \hbar\omega_i = 8\pi G\, S_{\text{matter}}. \tag{31}$$

This geometric sum has a precise identification in the gravitational action. On a stationary Rindler horizon the unperturbed expansion of the null generators vanishes, $\theta = 0$, so the Gibbons–Hawking–York boundary term [13] is sourced entirely by the AS pp-wave perturbations:

$$S_{GHY}\big|_{\mathcal{H}} = \frac{c^4}{8\pi G}\int_{\mathcal{H}} \theta\, d^2x_\perp\, du = \frac{c^4}{8\pi G}\sum_i \Delta A_i = \frac{c^4}{8\pi G}\cdot 8\pi G \sum_i \hbar\omega_i = \sum_i \hbar\omega_i = S_{\text{matter}}. \tag{32}$$

The matter action and the gravitational boundary action are therefore equal, event by event, on the stationary Rindler horizon. This is not a bulk result: the bulk Einstein–Hilbert action vanishes on-shell, and it is the boundary term that carries the physical content. The microstate construction identifies what that boundary content is: each contribution to S_{GHY} corresponds to one quantum of action deposited into matter-memory by one measurement event.

This result is consistent with and extends programmes of Jacobson [1] and Padmanabhan [14], which argue that the gravitational boundary term rather than the bulk Einstein–Hilbert action is the fundamental gravitational degree of freedom. Those programmes

establish this at the level of thermodynamic averages. The present construction realises the same identification at the level of individual microstates, with each boundary contribution sourced by a specific quantum measurement event rather than a bulk average. The standard result emerges from summing over the ensemble; the microstate picture supplies what the thermodynamic programmes leave open: the physical process underlying each boundary contribution.

5 Entanglement as Spatial Information

The bilocal source prepares an anticorrelated Bell state (23). We consider the simplest case: both $\mathcal{R}$ and $\mathcal{L}$ measure in the same basis and both obtain a definite outcome, which we label H_+ or H_-. In this case their outcomes are perfectly anticorrelated by the Bell state structure, independently of which basis is chosen. The question

"who observed H_+?"

then has a binary answer: $\mathcal{L}$ (left wedge) or $\mathcal{R}$ (right wedge). This is a *spatial bit*, a bit that encodes not a polarization state but a location in space.

Each measurement event has exactly two irreducible constituents: one quantum of action $\hbar$ written persistently into the detector's matter-memory as a series of samples, and one bit of polarization registered as a classical outcome. The entanglement between $\mathcal{R}$ and $\mathcal{L}$ is consumed in producing these two records, an increase in sampling rate and one bit at each wedge. What remains after measurement is not the entanglement itself but its classical residue: the perfect anticorrelation between the two bits, and the correlated pair of samples written into the matter on each side of the horizon. The spatial bit is precisely the polarization bit now read as a location rather than a spin state: the same bit that records *what* was observed also records *where*, because in this construction the two wedges carry opposite helicities by design. This double duty is not merely convenient but forced: the anticorrelated helicity assignment of the bilocal source makes the two wedges distinguishable by construction, so the spatial bit is already latent in the source geometry before any measurement occurs.

The information is carried by the matter of the detectors, not by the horizon. As established in Section **??**, the horizon is a boundary that makes the left/right distinction sharp; it is the matter on each side that accumulates the samples and holds the classical record. $\mathcal{R}$'s detector has absorbed one photon, increased its sampling rate, and registered polarization H_+/H_-. $\mathcal{L}$'s detector has absorbed one photon, increased its sampling rate, and registered polarization H_-/H_+. The two records are perfectly complementary by the anticorrelated structure of the bilocal source, and their complementarity is encoded in matter before it is reflected in geometry.

5.1 ER = EPR

The Unruh vacuum is the thermofield double state of the left and right Rindler wedges [15]: equation (21) is precisely this state, with the helicity label λ carried through. This places the present construction squarely within the ER=EPR setting of Maldacena and Susskind rather than merely analogous to it — the ER=EPR correspondence is the ambient mathematical structure within which the bilocal construction operates.

The bilocal source selects one term from the thermofield double sum at fixed Ω and fixed helicity pair, yielding the single microstate (23). The EPR side is the entangled photon pair emitted by $\mathcal{P}$, whose quantum correlations are consumed by the two measurement events. The ER side is the pair of correlated AS longitudinal shifts in opposite null directions: the geometric imprint of one term in the thermofield double sum, calculable and explicit. The spatial bit — the answer to "who observed H_+?" — is the classical

record of which side of the bridge each measurement occurred on. The bridge is not a stand-in for the entanglement but its physical instantiation, built from correlated matter samples, one quantum of action at a time.

What is new here is that both sides of the correspondence are calculable at the single-microstate level, with a definite energy-momentum source and a definite geometric output, rather than established by a bulk thermodynamic argument. The anticorrelated helicity structure gives the bridge an orientation absent in the scalar case: the two sides carry opposite handedness, and whether this constitutes a topological invariant of the transverse geometry is an open question we leave for future work.

This completes the microstate programme. Each entropy increment in the Bekenstein–Hawking area law corresponds to a single AS pp-wave interaction. Each pp-wave interaction corresponds to a single measurement event in the bilocal construction. Each measurement event both encodes a spatial bit and generates the spatial separation that bit refers to. The gravitational, thermodynamic, information-theoretic, and geometric descriptions of the horizon are unified at the level of individual microstates, and thermality, entropy, and space itself emerge together from the coarse-grained aggregate of these events.

6 Appendix: Consistency of the Double-Null Gauge Condition

We verify that the double-null gauge condition $A_+ = A_- = 0$ is consistent with Maxwell's equations and leaves exactly two physical degrees of freedom.

6.1 Coordinates and Notation

In double-null coordinates (x^+, x^-, x^1, x^2) the Minkowski metric reads

$$ds^2 = -2\,dx^+\,dx^- + (dx^1)^2 + (dx^2)^2, \tag{33}$$

so that $\partial_+ = \partial/\partial x^+$, $\partial_- = \partial/\partial x^-$, and the d'Alembertian is

$$\Box = -2\partial_+\partial_- + \partial_i\partial^i, \qquad i = 1, 2. \tag{34}$$

The gauge field has components $A_\mu = (A_+, A_-, A_1, A_2)$ and transforms as

$$A_\mu \to A_\mu + \partial_\mu \Lambda \tag{35}$$

under a gauge transformation with parameter $\Lambda(x^+, x^-, x^1, x^2)$.

6.2 Imposing the Gauge Conditions in Stages

The single gauge parameter Λ is a function of all four coordinates. We use it in three stages, each consuming a different functional dependence.

Stage 1: $A_+ = 0$**.** Requiring $A_+ + \partial_+\Lambda = 0$ determines Λ as a function of x^+:

$$\Lambda(x^+, x^-, x^i) = -\int^{x^+} A_+(s, x^-, x^i)\,ds + \Lambda_0(x^-, x^i), \tag{36}$$

where $\Lambda_0(x^-, x^i)$ is an arbitrary residual function independent of x^+.

Stage 2: $A_- = 0$**.** Under the residual freedom $\Lambda_0(x^-, x^i)$, the A_- component transforms as $A_- \to A_- + \partial_-\Lambda_0$. Setting this to zero determines Λ_0 as a function of x^-:

$$\Lambda_0(x^-, x^i) = -\int^{x^-} A_-(s, x^i)\,ds + \Lambda_1(x^i), \tag{37}$$

where $\Lambda_1(x^i)$ is a residual function of the transverse coordinates only.

Stage 3: Transverse Coulomb condition $\partial^i A_i = 0$. Under the remaining freedom $\Lambda_1(x^i)$, the transverse components transform as $A_i \to A_i + \partial_i \Lambda_1$. Imposing $\partial^i A_i = 0$ requires

$$\nabla_\perp^2 \Lambda_1 = -\partial^i A_i, \tag{38}$$

which is a Poisson equation in the transverse plane with a unique solution (up to harmonic functions, fixed by boundary conditions). This exhausts the gauge freedom entirely.

The three conditions have consumed the full gauge parameter Λ in stages, with no over-determination: each stage uses a strictly different functional dependence of Λ.

6.3 Physical Degrees of Freedom

After imposing $A_+ = A_- = 0$ and $\partial^i A_i = 0$, the remaining components are A_1 and A_2 subject to one differential constraint $\partial^i A_i = 0$. In momentum space this reads $k^i A_i = 0$.

For our specific construction, the photons propagate in the $\pm z$ direction, so their wave-vectors are purely null: $k_\perp^i = 0$ for $i = 1, 2$. The transverse Coulomb condition $k^i A_i = 0$ is therefore trivially satisfied for any A_1, A_2, and both transverse components are independent. The two circular polarization vectors

$$\epsilon^i_{(+1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}, \qquad \epsilon^i_{(-1)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix} \tag{39}$$

are the two independent physical degrees of freedom, corresponding to helicity $\lambda = \pm 1$. This is the correct count for a massless spin-1 field in four dimensions.

The trivial satisfaction of the transverse Coulomb condition relies on the photons propagating purely in the $\pm z$ direction with vanishing transverse momentum $k_\perp^i = 0$. For photons with nonzero transverse momentum the condition $k^i A_i = 0$ imposes a non-trivial constraint and the degree of freedom count requires revisiting.

6.4 Consistency with Maxwell's Equations

With $A_+ = A_- = 0$ and $\partial^i A_i = 0$, the non-zero field strength components are

$$F_{+i} = \partial_+ A_i, \qquad F_{-i} = \partial_- A_i, \qquad F_{ij} = \partial_i A_j - \partial_j A_i, \qquad F_{+-} = 0. \tag{40}$$

Maxwell's equations $\partial^\mu F_{\mu\nu} = J_\nu$ decompose as follows.

Transverse components $\nu = i$:

$$-2\partial_+ \partial_- A_i + \partial^j (\partial_j A_i - \partial_i A_j) = J_i. \tag{41}$$

Using $\partial^i A_i = 0$, the second term simplifies to $\partial^j \partial_j A_i$, giving the standard wave equation

$$\boxed{\Box A_i = J_i.} \tag{42}$$

Null components $\nu = \pm$:

$$-\partial^i \partial_\pm A_i = J_\pm. \tag{43}$$

These are constraint equations relating the null source components to the transverse field. For our bilocal source, which is purely transverse, $J_+ = J_- = 0$, and the constraints reduce to

$$\partial^i \partial_\pm A_i = 0. \tag{44}$$

Differentiating the transverse Coulomb condition $\partial^i A_i = 0$ with respect to $x^\pm$ gives $\partial^i (\partial_\pm A_i) = 0$, so these constraints are automatically satisfied. ✓

6.5 Summary

The double-null gauge condition $A_+ = A_- = 0$ supplemented by the transverse Coulomb condition $\partial^i A_i = 0$:

1. uses the gauge freedom of Λ completely and without over-determination, via a three-stage procedure consuming the x^+-, x^--, and x^i-dependent parts of Λ in sequence;
2. leaves exactly two physical degrees of freedom, the helicity eigenstates $\lambda = \pm 1$, consistent with the known count for a massless spin-1 field;
3. reduces Maxwell's equations to the standard transverse wave equation $\Box A_i = J_i$, with the null constraint equations automatically satisfied for a purely transverse source.

The gauge is therefore internally consistent and does not over-constrain the physical degrees of freedom. The $\mathbb{Z}_2$ wedge-swap symmetry is preserved throughout: conditions on A_+ and A_- are imposed symmetrically, and neither null direction is privileged over the other.

7 Conclusion

The construction developed above can be situated within several broader programmes. The ER=EPR correspondence of Maldacena and Susskind [15] is the ambient mathematical structure: the Unruh vacuum is the thermofield double state, and the bilocal source selects individual non-thermal microstates from it, with the EPR side being the entangled photon pair and the ER side being the pair of correlated AS longitudinal shifts in opposite null directions. Van Raamsdonk's argument [16] that entanglement holds bulk geometry together is realised here operationally: each measurement event converts entanglement into a geometric imprint, and the spatial bit encodes which side of the bridge each measurement occurred on. The boundary excitations of Carlip's near-horizon BMS algebra [**?**] are identified with AS supertranslations sourced by individual measurement events, and the discrete area increments of Bianchi [**?**] correspond to the calculable $\Delta A = 8\pi G\hbar\omega$ of Section 4, now with a definite dynamical origin in a quantum measurement event. More broadly, the construction realises Wheeler's "it from bit" programme [21] at the level of individual microstates: each bit acquired by a physical observer is backed by a measurement-induced energy-momentum transfer whose gravitational imprint incrementally builds horizon geometry.

Susskind and Maldacena's ER=EPR correspondence [15] identifies entangled pairs with Einstein–Rosen bridges at the level of bulk thermodynamics; we realize this correspondence operationally at the level of a single microstate, with a calculable geometric imprint sourced by a specific bilocal emission event. The Unruh vacuum is precisely the thermofield double state of the left and right Rindler wedges, placing our present construction squarely within the ER=EPR context.

Van Raamsdonk [16] argued that entanglement between boundary degrees of freedom is what holds bulk geometry together, and that reducing entanglement tears spacetime apart, a result that motivates our identification of the ER=EPR bridge with correlated measurement events, though his framework operates with thermal boundary states rather than individual non-thermal microstates.

Microstate counting for black holes has been approached from several directions. Strominger and Vafa [17] achieved exact entropy counting for extremal black holes using D-brane configurations in string theory, establishing that black hole entropy has a microscopic origin, but the microstates are defined algebraically in a high-energy string framework far removed from the operational measurement picture developed here. Carlip [**?**] derived black hole entropy from the Virasoro algebra of diffeomorphisms at the horizon boundary, identifying microstates with boundary excitations; our AS pp-wave imprints

may provide a physical realization of these boundary excitations in terms of measurement events a connection we make precise in Section 4: the AS longitudinal shift is a horizon supertranslation in the sense of Hawking, Perry, and Strominger [12], and the bilocal construction supplies the physical process, quantum measurement, that excites each mode of the algebra.

Bianchi's loop quantum gravity microstate counting [**?**] assigns discrete area increments to individual spin network punctures at the horizon, structurally analogous to our result that each measurement event contributes a discrete AS-wave area increment; the difference is that our punctures have a definite dynamical origin in a quantum measurement event rather than a kinematical definition in the LQG Hilbert space. Interior reconstruction via the Papadodimas–Raju programme [**?**] addresses how degrees of freedom behind the horizon are encoded in exterior operators, a question closely related to our spatial bit argument in Section 5. Their framework requires thermality of the boundary state, whereas our construction works microstate by microstate with a non-thermal source; it would be interesting to determine whether our classical spatial record, the anticorrelated helicity bits registered by right and left wedges, realizes the interior reconstruction map at the single-microstate level, but we leave this comparison for future work.

7.1 Subjectivity and Existing Ontologies

A closely related perspective is provided by relational and information-theoretic interpretations of quantum theory. Carlo Rovelli's relational quantum mechanics [18] and the QBist programme associated with Christopher Fuchs [19] both treat the quantum state as observer-dependent, encoding information rather than objective physical reality. In these frameworks, measurement is understood as an update of an observer's knowledge, not a universal physical collapse. The Unruh effect [8] provides a direct physical analogue of this observer-relativity: particle content and thermality themselves become observer-dependent notions, tied to acceleration. Our construction extends this shared principle across domains, identifying quantum state reduction, thermality, and gravitational effects as different aspects of observer-relative descriptions grounded in underlying physical interactions. This viewpoint also sharpens the conceptual tension highlighted by Eugene Wigner [20], where different observers may consistently assign different physical descriptions to the same process, suggesting that the resulting geometric imprints in our framework may likewise be observer-dependent. Finally, the present approach may be viewed as a concrete dynamical realization of Wheeler's "it from bit" programme [21]. Where Wheeler's proposal remained largely conceptual, discrete acts of information acquisition giving rise to physical law, we identify an explicit dynamical mechanism: each bit acquired by a physical observer is backed by a measurement-induced energy-momentum transfer, in our example, modelled geometrically as an AS pp-wave, whose gravitational imprint incrementally builds spacetime structure. The bit does not merely describe the geometry; it produces it.

8 Acknowledgements

I am also grateful to Claude for helpful discussions, noting with some irony that each such exchange constitutes, in the framework of this paper, a sequence of measurement events whose Landauer cost was paid by the data centers hosting them.

References

[1] T. Jacobson, "Thermodynamics of spacetime: the einstein equation of state," *Physical review letters*, vol. 75, no. 7, p. 1260, 1995.

[2] E. Verlinde, "On the origin of gravity and the laws of newton," *Journal of High Energy Physics*, vol. 2011, no. 4, pp. 1–27, 2011.

[3] T. Dray and G. Hooft, "The gravitational shock wave of a massless particle," *Nuclear physics B*, vol. 253, pp. 173–188, 1985.

[4] R. Penrose, "On gravity's role in quantum state reduction," *General relativity and gravitation*, vol. 28, no. 5, pp. 581–600, 1996.

[5] L. Diosi, "A universal master equation for the gravitational violation of quantum mechanics," *Physics letters A*, vol. 120, no. 8, pp. 377–381, 1987.

[6] K. Player, "Driving the unruh response," *arXiv preprint arXiv:2509.16710*, 2025.

[7] R. Landauer, "Irreversibility and heat generation in the computing process," *IBM journal of research and development*, vol. 5, no. 3, pp. 183–191, 1961.

[8] W. G. Unruh, "Notes on black-hole evaporation," *Physical Review D*, vol. 14, no. 4, p. 870, 1976.

[9] J. D. Bekenstein, "Black holes and entropy," *Phys. Rev. D*, vol. 7, pp. 2333–2346, Apr 1973.

[10] C. C. Gerry and P. L. Knight, *Introductory quantum optics*. Cambridge university press, 2023.

[11] N. D. Birrell and P. C. W. Davies, "Quantum fields in curved space," 1984.

[12] S. W. Hawking, M. J. Perry, and A. Strominger, "Soft hair on black holes," *Physical Review Letters*, vol. 116, no. 23, p. 231301, 2016.

[13] G. W. Gibbons and S. W. Hawking, "Action integrals and partition functions in quantum gravity," *Physical Review D*, vol. 15, no. 10, p. 2752, 1977.

[14] T. Padmanabhan, "Thermodynamical aspects of gravity: new insights," *Reports on Progress in Physics*, vol. 73, no. 4, p. 046901, 2010.

[15] J. Maldacena and L. Susskind, "Cool horizons for entangled black holes," *Fortschritte der Physik*, vol. 61, no. 9, pp. 781–811, 2013.

[16] M. Van Raamsdonk, "Building up space–time with quantum entanglement," *International Journal of Modern Physics D*, vol. 19, no. 14, pp. 2429–2435, 2010.

[17] A. Strominger and C. Vafa, "Microscopic origin of the bekenstein-hawking entropy," *Physics Letters B*, vol. 379, no. 1-4, pp. 99–104, 1996.

[18] C. Rovelli, "Relational quantum mechanics," *International journal of theoretical physics*, vol. 35, no. 8, pp. 1637–1678, 1996.

[19] C. A. Fuchs, N. D. Mermin, and R. Schack, "An introduction to qbism with an application to the locality of quantum mechanics," *American Journal of Physics*, vol. 82, no. 8, pp. 749–754, 2014.

[20] E. P. Wigner, "Remarks on the mind-body question," in *Philosophical reflections and syntheses*, pp. 247–260, Springer, 1995.

[21] J. A. Wheeler, "Information, physics, quantum: The search for links," *Feynman and computation*, pp. 309–336, 2018.