

BRGY. WAWA, TAGUIG HEALTH CENTER

# BABY VACCINES

**PRENATAL CHECK UP 8AM-12PM(THURSDAY) AFTERNOON CONSULTATIONS**



Vaccine	Minimum Age at 1st Dose	Number of Doses	Dose	Minimum Interval Between Doses	Route	Reason
Bacillus Calmette-Guérin	Birth or any time after birth	1 dose	0.05 mL	none	Intradermal	BCG given at earliest possible age protects the possibility of TB meningitis and other TB infections in which infants are prone
Pentavalent Vaccine		3 doses	0.5 mL	4weeks	Intramuscular	Protection against multiple diseases iphtheria, tetanus, pertussis (whooping cough), hepatitis B, and Haemophiles influenzae type b (Hib).
Oral Polio Vaccine	6 weeks old	3 doses	2-3 drops	4 weeks	Oral	The extent of protection against polio is increased the earlier the OPV is given.
Hepatitis B Vaccine	At birth	3 doses	0.5 mL	4 weeks interval	Intramuscular	An early start of Hepatitis B vaccine reduces the chance of being infected and becoming a carrier.
Measles Vaccine	9 months old	1 dose	0.5 mL	none	Subcutaneous	At least 85% of measles can be prevented by immunization at this age.