

# Stochastic Calculus

Author: Kellen Kanarios

Mentor: Ethan Zell

University of Michigan

April 11, 2022

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Note

Intuition for a **measure**:

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Note

Intuition for a **measure**:

1. In  $\mathbb{R}$ : length,

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Note

Intuition for a **measure**:

1. In  $\mathbb{R}$ : length,
2. In  $\mathbb{R}^2$ : area,

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Note

Intuition for a **measure**:

1. In  $\mathbb{R}$ : length,
2. In  $\mathbb{R}^2$ : area,
3. In  $\mathbb{R}^3$ : volume.

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

Definition translated into english

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Definition translated into english

1. Unbeknownst to us, someone chooses a random  $\omega \in \Omega$ . Then we see the  $X(\omega) \in \mathbb{R}$ .



# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Definition translated into english

1. Unbeknownst to us, someone chooses a random  $\omega \in \Omega$ . Then we see the  $X(\omega) \in \mathbb{R}$ .
2. We cannot see the corresponding  $\omega \in \Omega$ , but the  $X(\omega) \in \mathbb{R}$  gives us partial information about  $\omega$ .

# Background

## Definition

A real-valued random variable  $X$  is a mapping  $X : \Omega \rightarrow \mathbb{R}$ . Here,  $\Omega$  is the **sample space** and  $\mathbb{P}$  is the measure of the **probability space**, such that  $\mathbb{P}(\Omega) = 1$ .

## Definition translated into english

1. Unbeknownst to us, someone chooses a random  $\omega \in \Omega$ . Then we see the  $X(\omega) \in \mathbb{R}$ .
2. We cannot see the corresponding  $\omega \in \Omega$ , but the  $X(\omega) \in \mathbb{R}$  gives us partial information about  $\omega$ .
3.  $\mathbb{P}$  tells us how likely subsets  $A \subseteq \Omega$  are to occur.

# Background

## Example

Consider the case where you flip a coin. Using our previous definition, this could be described as  $\Omega = \{\text{heads}, \text{tails}\}$  and

$$X(\omega) = \begin{cases} 1, & \text{if } \omega = \text{heads} \\ -1, & \text{if } \omega = \text{tails} \end{cases} \quad \text{where } \omega \in \Omega.$$

This would yield the familiar notation of  $\mathbb{P}(X = 1) = .5$  and  $\mathbb{P}(X = -1) = .5$  for a fair coin.

# Background

## Definition

A **stochastic process** is a family of random variables indexed by a time parameter  $t \geq 0$ .

# Background

## Definition

A **stochastic process** is a family of random variables indexed by a time parameter  $t \geq 0$ .

## Example

A sequence of coin flips. At each time  $t$  your process corresponds to a random variable (aka coinflip)  $X_t$ .

# Wiener Process

## Definition

A stochastic process  $W$  is called a **Wiener process** if the follow conditions hold:

# Wiener Process

## Definition

A stochastic process  $W$  is called a **Wiener process** if the follow conditions hold:

1.  $W_0 = 0$ ,

# Wiener Process

## Definition

A stochastic process  $W$  is called a **Wiener process** if the follow conditions hold:

1.  $W_0 = 0$ ,
2. The process  $W$  has independent increments,




# Wiener Process

## Definition

A stochastic process  $W$  is called a **Wiener process** if the follow conditions hold:

1.  $W_0 = 0$ ,
2. The process  $W$  has independent increments,
3. For  $s < t$  the random variable  $W_t - W_s$  has the Gaussian distribution  $\mathcal{N}(0, t - s)$ ,




# Wiener Process

## Definition

A stochastic process  $W$  is called a **Wiener process** if the follow conditions hold:

1.  $W_0 = 0$ ,
2. The process  $W$  has independent increments,
3. For  $s < t$  the random variable  $W_t - W_s$  has the Gaussian distribution  $\mathcal{N}(0, t - s)$ ,



4.  $W$  has continuous trajectories.

# Wiener Process

## Theorem

*A Wiener trajectory is with probability one, nowhere differentiable, and it has locally infinite total variation.*




Figure: Wiener trajectory

# Quadratic Variation

## Definition

Let  $X$  be a stochastic process. Suppose  $P$  is a partition of  $[0, t]$  denoted  $t_k$  and let  $\|P\|$  be the mesh of the partition then the **quadratic variation** of  $X$  is defined to be:

$$[X]_t = \lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (X_{t_k} - X_{t_{k-1}})^2.$$

# Quadratic Variation

## Definition

Let  $X$  be a stochastic process. Suppose  $P$  is a partition of  $[0, t]$  denoted  $t_k$  and let  $\|P\|$  be the mesh of the partition then the **quadratic variation** of  $X$  is defined to be:

$$[X]_t = \lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (X_{t_k} - X_{t_{k-1}})^2.$$

## Comments

Note that Quadratic Variation itself is a stochastic process. An intuitive way to think about quadratic variation is the internal clock of a process, describing how randomness accumulates over time.

# Quadratic Variation of Wiener Process

## Theorem

*Quadratic variation of a Wiener Process on the interval  $[0, t]$  is  $t$ .*

# Quadratic Variation of Wiener Process

## Theorem

*Quadratic variation of a Wiener Process on the interval  $[0, t]$  is  $t$ .*

## Proof.

Let us fix a point  $t$  and subdivide the interval  $[0, t]$  into  $n$  equally large subintervals of the form  $[k\frac{t}{n}, (k+1)\frac{t}{n}]$ , where  $k = 0, 1, \dots, n-1$ . Then the quadratic variation on  $P$  is:

$$[W]_t^P = \sum_{i=1}^n (W_{i\frac{t}{n}} - W_{(i-1)(\frac{t}{n})})^2.$$

# Quadratic Variation of Wiener Process

## Theorem

*Quadratic variation of a Wiener Process on the interval  $[0, t]$  is  $t$ .*

## Proof.

Let us fix a point  $t$  and subdivide the interval  $[0, t]$  into  $n$  equally large subintervals of the form  $[k\frac{t}{n}, (k+1)\frac{t}{n}]$ , where  $k = 0, 1, \dots, n-1$ . Then the quadratic variation on  $\bar{P}$  is:

$$[W]_t^P = \sum_{i=1}^n (W_{i\frac{t}{n}} - W_{(i-1)(\frac{t}{n})})^2.$$

Therefore,

$$\mathbb{E} [[W]_t^P] = \sum_{i=1}^n \mathbb{E} \left[ (W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})})^2 \right].$$



# Quadratic Variation of Wiener Process

## Theorem

$$\text{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}(X))^2.$$

# Quadratic Variation of Wiener Process

Proof cont.

Since  $W_{(i)(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ ,

$$\mathbb{E}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = 0.$$

# Quadratic Variation of Wiener Process

Proof cont.

Since  $W_{(i)(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ ,

$$\mathbb{E}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = 0.$$

It follows that:

$$(\mathbb{E} [W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}])^2 = 0.$$

# Quadratic Variation of Wiener Process

Proof cont.

Since  $W_{(i)(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ ,

$$\mathbb{E}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = 0.$$

It follows that:

$$(\mathbb{E} [W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}])^2 = 0.$$

Thus,  $\text{Var}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = E([W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}]^2).$

# Quadratic Variation of Wiener Process

## Proof cont.

Since  $W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ ,

$$\mathbb{E}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = 0.$$

It follows that:

$$(\mathbb{E} [W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}])^2 = 0.$$

$$\text{Thus, } \text{Var}(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}) = E\left([W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}]^2\right).$$

Since  $\mathbb{E}[(W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})})^2] = \text{Var} [W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}]$ , we can write:

$$\mathbb{E} [ [W]_t^P ] = \mathbb{E} \left[ \sum_{k=1}^n (W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})})^2 \right] = \text{Var} \left[ \sum_{k=1}^n [W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}] \right].$$

# Quadratic Variation of Wiener Process

Stats fact

$$\begin{aligned} \text{Var} \left[ \sum_{k=1}^n \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right] \right] &= \sum_{k=1}^n \text{Var} \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right] \\ &\quad + \sum_{i \neq \ell}^n \text{Cov} \left( \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right], \left[ W_{\ell(\frac{t}{n})} - W_{(\ell-1)(\frac{t}{n})} \right] \right) \end{aligned}$$

# Quadratic Variation of Wiener Process

## Proof cont.

Since Wiener increments are independent of each other,

$$\text{Cov} \left( \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right], \left[ W_{\ell(\frac{t}{n})} - W_{(\ell-1)(\frac{t}{n})} \right] \right) = 0.$$

# Quadratic Variation of Wiener Process

## Proof cont.

Since Wiener increments are independent of each other,

$$\text{Cov} \left( \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right], \left[ W_{\ell(\frac{t}{n})} - W_{(\ell-1)(\frac{t}{n})} \right] \right) = 0.$$

Therefore,

$$E \left[ [W]_t^P \right] = \sum_{k=1}^n \text{Var} \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right]$$



# Quadratic Variation of Wiener Process

## Proof cont.

Since Wiener increments are independent of each other,

$$\text{Cov} \left( \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right], \left[ W_{\ell(\frac{t}{n})} - W_{(\ell-1)(\frac{t}{n})} \right] \right) = 0.$$

Therefore,

$$E \left[ [W]_t^P \right] = \sum_{k=1}^n \text{Var} \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right]$$

Using  $W_{(i)(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ :

$$\begin{aligned} \mathbb{E} \left[ [W]_t^P \right] &= \sum_{k=1}^n \left( i(\frac{t}{n}) - (i-1)(\frac{t}{n}) \right) \\ &= t \end{aligned}$$

# Quadratic Variation of Wiener Process

## Proof cont.

Since Wiener increments are independent of each other,

$$\text{Cov} \left( \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right], \left[ W_{\ell(\frac{t}{n})} - W_{(\ell-1)(\frac{t}{n})} \right] \right) = 0.$$

Therefore,

$$\mathbb{E} \left[ [W]_t^P \right] = \sum_{k=1}^n \text{Var} \left[ W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \right]$$

Using  $W_{(i)(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})} \in \mathcal{N}(0, i(\frac{t}{n}) - (i-1)(\frac{t}{n}))$ :

$$\begin{aligned} \mathbb{E} \left[ [W]_t^P \right] &= \sum_{k=1}^n \left( i(\frac{t}{n}) - (i-1)(\frac{t}{n}) \right) \\ &= t \end{aligned}$$

Now we must show  $\mathbb{E}([W]_t^P) = [W]_t = t$ .

# Quadratic Variation of Wiener Process

## Stats fact

If  $\text{Var}(X) = 0$  then  $X = \mathbb{E}[X]$ .

# Quadratic Variation of Wiener Process

Proof.

Therefore, we must show that :

$$\text{Var}\left(\lim_{\|P\|\rightarrow 0} [W]_t^P\right) = 0.$$

Or equivalently,

$$\text{Var}\left(\lim_{n\rightarrow\infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) = 0,$$

# Quadratic Variation of Wiener Process

Proof.

Therefore, we must show that :

$$\text{Var}\left(\lim_{\|P\|\rightarrow 0} [W]_t^P\right) = 0.$$

Or equivalently,

$$\text{Var}\left(\lim_{n\rightarrow\infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) = 0,$$

and we showed earlier:

$$\text{Var}\left(\lim_{n\rightarrow\infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) = \lim_{n\rightarrow\infty} \sum_{i=1}^n \text{Var}\left(\left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right).$$

# Quadratic Variation of Wiener Process

Proof.

Also from earlier:

$$\begin{aligned} \text{Var}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbb{E}\left(\left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^4\right) \\ &\quad - \mathbb{E}\left(\left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right)^2. \end{aligned}$$

# Quadratic Variation of Wiener Process

Stats fact

$$E[(X - \mu)^p] = \begin{cases} 0, & \text{if } p \text{ is odd} \\ \sigma^p (p-1)!!, & \text{if } p \text{ is even.} \end{cases}$$

# Quadratic Variation of Wiener Process

Proof.

Also from earlier:

$$\begin{aligned} \text{Var}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbb{E}\left(\left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^4\right) \\ &\quad - \mathbb{E}\left(\left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right)^2. \end{aligned}$$

Therefore,

$$\begin{aligned} \text{Var}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 3 \left(i(\frac{t}{n}) - (i-1)(\frac{t}{n})\right)^2 \\ &\quad - \left(i(\frac{t}{n}) - (i-1)(\frac{t}{n})\right)^2. \end{aligned}$$



# Quadratic Variation of Wiener Process

Proof.

Simplifying,

$$\begin{aligned}\text{Var}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 2 \frac{t^2}{n^2}, \\ &= \lim_{n \rightarrow \infty} 2 \frac{t^2}{n}, \\ &= 0.\end{aligned}$$

# Quadratic Variation of Wiener Process

Proof.

Simplifying,

$$\begin{aligned}\text{Var}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[W_{i(\frac{t}{n})} - W_{(i-1)(\frac{t}{n})}\right]^2\right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 2 \frac{t^2}{n^2}, \\ &= \lim_{n \rightarrow \infty} 2 \frac{t^2}{n}, \\ &= 0.\end{aligned}$$

Thus,

$$[W]_t = \mathbb{E}([W]_t) = \mathbb{E}\left(\lim_{\|P\| \rightarrow 0} [W]_t^P\right).$$

# Quadratic Variation of Wiener Process

Proof.

By dominating convergence theorem (outside scope),

$$\begin{aligned}\mathbb{E} \left( \lim_{\|P\| \rightarrow 0} [W]_t^P \right) &= \lim_{\|P\| \rightarrow 0} \mathbb{E} \left( [W]_t^P \right), \\ &= \lim_{\|P\| \rightarrow 0} t, \\ &= t.\end{aligned}$$



# Quadratic Variation of Wiener Process

## Proof.

By dominating convergence theorem (outside scope),

$$\begin{aligned}\mathbb{E} \left( \lim_{\|P\| \rightarrow 0} [W]_t^P \right) &= \lim_{\|P\| \rightarrow 0}, \mathbb{E} \left( [W]_t^P \right), \\ &= \lim_{\|P\| \rightarrow 0} t, \\ &= t.\end{aligned}$$

Thus,  $[W]_t = t$ .



# Quadratic Variation of Wiener Process

## Implications

This motivates us to write:

$$\int_0^t (dW_t)^2 = t.$$

Or equivalently,

$$(dW_t)^2 = dt.$$

This will come up frequently later as we transition into Ito calculus.

# The Stochastic Integral

## Motivation

Let  $h_t$  be a stochastic process that represents our trading strategy and let  $W_t$  be the price of the stock at the given time. Then

$$\int_0^t h_t dW_t$$

would be our gains or losses from this strategy.

# The Stochastic Integral

## The Problem

Integrals of the form  $\int_0^t g_s dW_s$  for some stochastic process  $g_s$ .

# The Stochastic Integral

## The Problem

Integrals of the form  $\int_0^t g_s dW_s$  for some stochastic process  $g_s$ .

## Possible Solution 1

Try to define like the Riemman Integral:

1. 
$$\sum_{k=1}^n g_s(W_{t_{k+1}} - W_{t_k}),$$



# The Stochastic Integral

## The Problem

Integrals of the form  $\int_0^t g_s dW_s$  for some stochastic process  $g_s$ .

## Possible Solution 1

Try to define like the Riemman Integral:

1.  $\sum_{k=1}^n g_s(W_{t_{k+1}} - W_{t_k}),$
2. Not possible due to locally unbounded variation of a Wiener Process.

# The Stochastic Integral

## Definition

A stochastic process is **simple** on  $[a, b]$  when there exists deterministic points in time  $a = t_0 < t_1 < \cdots < t_n = b$  such that  $g_s = g_{t_k}$  for all  $s \in [t_k, t_{k+1}]$ .

# The Stochastic Integral

## Definition

A stochastic process is **simple** on  $[a, b]$  when there exists deterministic points in time  $a = t_0 < t_1 < \dots < t_n = b$  such that  $g_s = g_{t_k}$  for all  $s \in [t_k, t_{k+1}]$ .

## Real Solution

Now we can define the stochastic integral by

$$\int_a^b g_s dW_s = \sum_{k=0}^{n-1} g_{t_k} [W_{t_{k+1}} - W_{t_k}],$$

for some simple stochastic process  $g_s$ .

# The Stochastic Integral

## Generalized Version

By the **simple approximation theorem** (outside of scope), for some stochastic process  $g_s$  there exists a sequence of simple functions  $g_s^n$ , such that  $g_s^n \rightarrow g_s$ . Therefore, we define our integral for non-simple functions as:

$$\int_a^b g_s dW_t = \lim_{n \rightarrow \infty} \int_a^b g_s^n dW_t.$$

# The Stochastic Integral

## Generalized Version

By the **simple approximation theorem** (outside of scope), for some stochastic process  $g_s$  there exists a sequence of simple functions  $g_s^n$ , such that  $g_s^n \rightarrow g_s$ . Therefore, we define our integral for non-simple functions as:

$$\int_a^b g_s dW_t = \lim_{n \rightarrow \infty} \int_a^b g_s^n dW_t.$$

## Something to think about

How might we evaluate  $\int_0^t W_s dW_s$ ?

# Stochastic Differential Equations

## Definition

An ito process,  $X_t$ , is a process that can be represented as:

$$X_t = X_0 + \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s.$$

# Stochastic Differential Equations

## Definition

An ito process,  $X_t$ , is a process that can be represented as:

$$X_t = X_0 + \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s.$$

We often use the notation:

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

# Stochastic Differential Equations

## Definition

An Ito process,  $X_t$ , is a process that can be represented as:

$$X_t = X_0 + \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s.$$

We often use the notation:

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

This is known as a stochastic differential equation, and a useful tool for solving these is Ito's lemma, which we will see shortly.



# Ito's Multiplication Table

## Definition

**Ito's multiplication table** is:

	$dt$	$dW_t$
$dt$	0	0
$dW_t$	0	$dt$

We have already shown the only interesting result:  $(dW_t)^2 = dt$ .

# Ito's Multiplication Table

## Definition

**Ito's multiplication table** is:

	$dt$	$dW_t$
$dt$	0	0
$dW_t$	0	$dt$

We have already shown the only interesting result:  $(dW_t)^2 = dt$ .

## Quick sketch.

Suppose you are integrating on the interval  $[a, b]$ . Partition  $[a, b]$  into  $N$  increments. Then  $dt \sim \frac{1}{N}$  and  $(dt)^2 \sim \frac{1}{N^2}$ . It follows that:

$$\sum_{i=1}^N \frac{1}{N^2} = \frac{1}{N},$$
$$\lim_{N \rightarrow \infty} \sum_{i=1}^N \frac{1}{N^2} = \lim_{N \rightarrow \infty} \frac{1}{N} = \int_a^b (dt)^2 = 0.$$

# Ito's Multiplication Table

## Quick sketch.

Similarly, suppose you are integrating on the interval  $[a, b]$ . Partition  $[a, b]$  into  $N$  increments. Since  $dW_t = \sqrt{dt}$ ,  $dt \sim \frac{1}{N}$  and  $(dt)(dW_t) \sim \frac{1}{N^{3/2}}$ . It follows that:

$$\sum_{i=1}^N \frac{1}{N^{3/2}} = \frac{1}{N^{1/2}},$$

$$\lim_{N \rightarrow \infty} \sum_{i=1}^N \frac{1}{N^{3/2}} = \lim_{N \rightarrow \infty} \frac{1}{N^{1/2}} = \int_a^b (dt)(dW_t) = 0.$$



# Ito's Multiplication Table

## Quick sketch.

Similarly, suppose you are integrating on the interval  $[a, b]$ . Partition  $[a, b]$  into  $N$  increments. Since  $dW_t = \sqrt{dt}$ ,  $dt \sim \frac{1}{N}$  and  $(dt)(dW_t) \sim \frac{1}{N^{3/2}}$ . It follows that:

$$\sum_{i=1}^N \frac{1}{N^{3/2}} = \frac{1}{N^{1/2}},$$
$$\lim_{N \rightarrow \infty} \sum_{i=1}^N \frac{1}{N^{3/2}} = \lim_{N \rightarrow \infty} \frac{1}{N^{1/2}} = \int_a^b (dt)(dW_t) = 0.$$



This completes the multiplication table and provides the foundation for introducing Ito's lemma.

# Ito's Lemma

## Theorem (Ito's formula)

*Assume that process  $X$  has a stochastic differential given by*

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

*Define the process  $Z$  by  $Z(t) = f(t, X_t)$ . Then  $Z$  has a stochastic differential given by*

$$df(t, X_t) = \left\{ \frac{\partial f}{\partial t}(t, X_t) + \mu_t \frac{\partial f}{\partial x}(t, X_t) + \frac{1}{2} \sigma_t^2 \frac{\partial^2 f}{\partial x^2}(t, X_t) \right\} dt + \sigma \frac{\partial f}{\partial x}(t, X_t) dW_t.$$

# Ito's Lemma

## Heuristic proof.

Using  $f$  from the theorem, we will consider the second order Taylor expansion:

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (dX_t)^2 + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

# Ito's Lemma

## Heuristic proof.

Using  $f$  from the theorem, we will consider the second order Taylor expansion:

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (dX_t)^2 + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

By definition we have:

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

# Ito's Lemma

## Heuristic proof.

Using  $f$  from the theorem, we will consider the second order Taylor expansion:

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (dX_t)^2 + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

By definition we have:

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

so, we obtain

$$(dX_t)^2 = \mu_t^2 (dt)^2 + 2\mu_t \sigma_t (dt)(dW_t) + \sigma_t^2 (dW_t)^2.$$



# Ito's Lemma

## Heuristic proof.

Using  $f$  from the theorem, we will consider the second order Taylor expansion:

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (dX_t)^2 + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

By definition we have:

$$dX_t = \mu_t dt + \sigma_t dW_t.$$

so, we obtain

$$(dX_t)^2 = \mu_t^2 (dt)^2 + 2\mu_t \sigma_t (dt)(dW_t) + \sigma_t^2 (dW_t)^2.$$

From Ito's multiplication table we have:

$$(dX_t)^2 = \sigma_t^2 dt.$$

# Ito's Lemma

## Heuristic proof cont.

Substituting back for  $(dX_t)^2$ ,

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$



# Ito's Lemma

## Heuristic proof cont.

Substituting back for  $(dX_t)^2$ ,

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

Applying Ito's multiplication table one more time,

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$



# Ito's Lemma

## Heuristic proof cont.

Substituting back for  $(dX_t)^2$ ,

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{1}{2} \frac{\partial^2 f}{\partial t^2} (dt)^2 + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

Applying Ito's multiplication table one more time,

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dX_t + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{\partial^2 f}{\partial t \partial x} dt dX_t.$$

Now substituting in  $(dX_t)$ ,

$$\begin{aligned} df &= \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} (\mu_t dt + \sigma_t dW_t) + \frac{1}{2} \frac{\partial^2 f}{\partial x^2} (\sigma_t^2)(dt) + \frac{\partial^2 f}{\partial t \partial x} (\mu_t dt + \sigma_t dW_t) dt, \\ &= \left\{ \frac{\partial f}{\partial t}(t, X_t) + \mu_t \frac{\partial f}{\partial x}(t, X_t) + \frac{1}{2} \sigma_t^2 \frac{\partial^2 f}{\partial x^2}(t, X_t) \right\} dt + \sigma \frac{\partial f}{\partial x}(t, X_t) dW_t. \end{aligned}$$



# Ito's Lemma

## Example

Find  $d(W_t^2)$ .

# Ito's Lemma

## Example

Find  $d(W_t^2)$ .

## Solution

Define  $X$  by  $X_t = W_t$  and set  $f(t, x) = x^2$ . In terms of Ito's formula we have

$$\frac{\partial f}{\partial t}(t, x) = 0, \quad \frac{\partial f}{\partial x}(t, x) = 2x, \quad \frac{\partial^2 f}{\partial t^2}(t, x) = 2.$$

# Ito's Lemma

## Example

Find  $d(W_t^2)$ .

## Solution

Define  $X$  by  $X_t = W_t$  and set  $f(t, x) = x^2$ . In terms of Ito's formula we have

$$\frac{\partial f}{\partial t}(t, x) = 0, \quad \frac{\partial f}{\partial x}(t, x) = 2x, \quad \frac{\partial^2 f}{\partial t^2}(t, x) = 2.$$

Substituting,

$$d(W_t^2) = \left\{ 0 + 0 \cdot 2X_t + \frac{1}{2} \cdot 1 \cdot 2 \right\} dt + 1 \cdot 2X_t dW_t.$$

# Ito's Lemma

## Example

Find  $d(W_t^2)$ .

## Solution

Define  $X$  by  $X_t = W_t$  and set  $f(t, x) = x^2$ . In terms of Ito's formula we have

$$\frac{\partial f}{\partial t}(t, x) = 0, \quad \frac{\partial f}{\partial x}(t, x) = 2x, \quad \frac{\partial^2 f}{\partial t^2}(t, x) = 2.$$

Substituting,

$$d(W_t^2) = \left\{ 0 + 0 \cdot 2X_t + \frac{1}{2} \cdot 1 \cdot 2 \right\} dt + 1 \cdot 2X_t dW_t.$$

Thus,

$$d(W_t^2) = dt + 2W_t dW_t.$$



# Ito's Lemma

## Revisiting

Now if we integrate both side such that:

$$\int_0^t d(W_t^2) = \int_0^t dt + \int_0^t 2W_t dW_t.$$

# Ito's Lemma

## Revisiting

Now if we integrate both side such that:

$$\int_0^t d(W_t^2) = \int_0^t dt + \int_0^t 2W_t dW_t.$$

This gives us:

$$\int_0^t W_t dW_t = \frac{W_t^2}{2} - \frac{t}{2}.$$

# Real world application

## Definition

**Geometric Brownian Motion** is a stochastic process whose dynamics follow the stochastic differential equation

$$dX_t = \alpha X_t dt + \sigma X_t dW_t \text{ for some } \alpha, \beta \in \mathbb{R},$$

where  $dW_t$  is the infinitesimal increment of the Wiener process.

# Real world application

## Definition

**Geometric Brownian Motion** is a stochastic process whose dynamics follow the stochastic differential equation

$$dX_t = \alpha X_t dt + \sigma X_t dW_t \text{ for some } \alpha, \beta \in \mathbb{R},$$

where  $dW_t$  is the infinitesimal increment of the Wiener process.

Geometric Brownian Motion is one of the building blocks for the modeling of asset prices, and turns up naturally in many other places.

# Real world application

## Closed form

With our toolbox of Ito calculus and the ability to solve linear ODE's, it can be easily shown that the closed form of Geometric Brownian Motion is:

$$dX_t = X_t \left( \mu - \frac{\sigma^2}{2} \right) dt + \sigma dW_t.$$

# Real world application

## Closed form

With our toolbox of Ito calculus and the ability to solve linear ODE's, it can be easily shown that the closed form of Geometric Brownian Motion is:

$$dX_t = X_t \left( \mu - \frac{\sigma^2}{2} \right) dt + \sigma dW_t.$$

Now let's see how accurate this elementary model is when trying to predict the price of real stocks.

# Real world application

## Use in Finance

Black Scholes Model is one of the most widely used models of stock price behaviour and is built on Geometric Brownian motion.

## Why GBM?

Some of the arguments for GBM are:

# Real world application

## Use in Finance

Black Scholes Model is one of the most widely used models of stock price behaviour and is built on Geometric Brownian motion.

## Why GBM?

Some of the arguments for GBM are:

- ▶ The expected returns are independent of the value of the process, which is how stocks behave in reality,



# Real world application

## Use in Finance

Black Scholes Model is one of the most widely used models of stock price behaviour and is built on Geometric Brownian motion.

## Why GBM?

Some of the arguments for GBM are:

- ▶ The expected returns are independent of the value of the process, which is how stocks behave in reality,
- ▶ A GBM process only assumes positive values, just like real stock prices.

# Real world application

## Use in Finance

Black Scholes Model is one of the most widely used models of stock price behaviour and is built on Geometric Brownian motion.

## Why GBM?

Some of the arguments for GBM are:

- ▶ The expected returns are independent of the value of the process, which is how stocks behave in reality,
- ▶ A GBM process only assumes positive values, just like real stock prices.
- ▶ Calculations with GBM processes are relatively easy.

# Real world application

## Drawbacks

# Real world application

## Drawbacks

- ▶ In real stock prices, volatility changes over time (possibly stochastically), but in GBM, volatility is assumed constant.

# Real world application

## Drawbacks

- ▶ In real stock prices, volatility changes over time (possibly stochastically), but in GBM, volatility is assumed constant.
- ▶ In real life, stock prices often show jumps caused by unpredictable events or news, but in GBM, the path is continuous (no discontinuity).

# Real world application

## Drawbacks

- ▶ In real stock prices, volatility changes over time (possibly stochastically), but in GBM, volatility is assumed constant.
- ▶ In real life, stock prices often show jumps caused by unpredictable events or news, but in GBM, the path is continuous (no discontinuity).

This is why GBM is just a building block for more advanced models. For example, one that models volatility stochastically is a stochastic volatility model, which is built off GBM.