

# Logolas: New Frontiers in Logo Visualization

Enrichment Depletion Logos (EDLogo) | String Logos | Adaptive Scaling of Logos

Kushal K Dey\*

Dongyue Xie

Matthew Stephens\*

Department of Statistics, University of Chicago
\*Department of Human Genetics, University of Chicago

## **Project Page**

https://kkdey.github.io/Logolas-pages

## Gallery

https://kkdey.github.io/Logolas-pages/Gallery.html

### R package

Github Version kkdey/Logolas

Bioconductor Logolas

#### CONTACT

Kushal K Dey

Email: kkdey@uchicago.edu

Github: kkdey
Twitter: kanishkadey
Website: kkdey.github.io

### **ACKNOWLEDGEMENTS**

John D. Blischak, Hussein Al-Asadi, Alex E. White, Trevor Price


#### REFERENCES

Stephens, M.: False discovery rates: a new deal. Biostatistics 18 (2), 275–294 (2016)
Koch, C.M., et al.: The landscape of histone modifications across 1in five human cell lines.
Genome Research 17(6), 691–707 (2007)
Alexandrov, L., Nik-Zainal, G., Wedge, D., Campbell, P., Stratton, M.: Deciphering signatures of mutational processes operative in human cancer. Cell Reports 3(1), 246–259 (2013)
Kheradpour, P., Kellis, M.: Systematic discovery and characterization of regulatory motifs in encode tf binding experiments. Nucleic Acids Research, 1–12 (2013)

Joseph, A.P., Shingate, P., Upadhyay, A.K., Sowdhamini, R.: 3pfdb+: improved search protocol and update for the identification of representatives of protein sequence domain families. Database (Oxford) bau026 (2014) Standard sequence logos tend to highlight enrichment primarily.


We propose *EDLogo* plots that highlight both enrichment and depletion at a position.






PSSM logo plots are often used to visualize protein sequence logos, but *EDLogo* representation is more parsimonious and interpretable than the PSSM logo plot.


Bacterial transcription activator, effector binding domain protein PF06445 (motif 4, Start=153 Length=8).



An application of alphanumeric string symbols in *Logolas* is in modeling the composition of histone marks in different regions of the genome (sue to Koch et al 2007).




Depletion is a biological phenomenon. Several transcription factors of EBF-1 family show base depletion in the middle of the motif. An example provided below for EBF1-disc1 transcription factor.




Logolas allows the user to plot string logos. An example application of such string logos is in visualizing mutational signature profiles.

EDLogo representation can detect depletion of G to the right of C mutation dominant signatures, possibly reflecting CpG site bias. We illustrate using an example from Alexandrov et al (2013).



Logolas provides an adaptive scaling of the Position Weight Matrix based on number of aligned sequences.

This method is called Dirichlet adaptive shrinkage or dash.



Poster Template Designed by Genigraphics ©2012 1.800.790.4001 www.genigraphics.com