

STAT234: Lecture 7 - Type I, Type II and Power

Kushal K. Dey

Last class

Last class

- ▶ Type I error : Probability that we reject the null hypothesis when it is true (fair for statistician)

Last class

- ▶ Type I error : Probability that we reject the null hypothesis when it is true (fair for statistician)
- ▶ Type II error: Probability that we accept the null hypothesis when it is false (fair for the researcher)

Last class

- ▶ Type I error : Probability that we reject the null hypothesis when it is true (fair for statistician)
- ▶ Type II error: Probability that we accept the null hypothesis when it is false (fair for the researcher)
- ▶ Power:

$$\text{Power} = 1 - \text{Type II error}$$

It is the probability that we reject the null hypothesis when it is false.

Last class

- ▶ Type I error : Probability that we reject the null hypothesis when it is true (fair for statistician)
- ▶ Type II error: Probability that we accept the null hypothesis when it is false (fair for the researcher)
- ▶ Power:

$$\text{Power} = 1 - \text{Type II error}$$

It is the probability that we reject the null hypothesis when it is false.

We want to minimize Type I error and maximize Power

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Type II error

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Type II error = failing to convict a guilty person

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Type II error = failing to convict a guilty person

In practice, we first choose an α and consider only tests with probability of Type I error no greater than α .

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Type II error = failing to convict a guilty person

In practice, we first choose an α and consider only tests with probability of Type I error no greater than α . Then we select one that makes Type II error as small as possible (test with most power)

Two Types of Errors Revisited

Recall that the following four outcomes are possible when conducting a test:

Reality	Our Decision	
	H_0	H_a
H_0	✓ (Prob = $1 - \alpha$)	Type I Error (Prob = α)
H_a	Type II Error (Prob = β)	✓ (Prob = $1 - \beta$)

Type I error is generally **more serious** than type II error.

For example: H_0 : the person is not guilty, H_1 : the person is guilty

Type I error = convicting an innocent person

Type II error = failing to convict a guilty person

In practice, we first choose an α and consider only tests with probability of Type I error no greater than α . Then we select one that makes Type II error as small as possible (test with most power)

Hypothesis testing framework

Last time we considered the example,
 X_1, X_2, \dots, X_n be data coming from

$$X_i \sim N(\mu, \sigma^2)$$

and we assumed that we know σ^2 . In our model,

$$\sigma = 2$$

Hypothesis to test

$$H_0 : \mu = 0 \quad H_1 : \mu > 0$$

We want to choose smallest c such that $\bar{X} > c$ happens with probability greater than or equal to the Type I error (0.05)

$$\text{find smallest } c : Pr [\bar{X} > c | \mu = 0] < 0.05$$

Hypothesis testing framework

$$\text{Type I error : } Pr \left[Z > \frac{\sqrt{n}}{\sigma} c | \mu = 0 \right] = 0.05 \quad Z \sim N(0, 1)$$

From this, we know

$$\frac{\sqrt{n}}{\sigma} c = 1.644$$

Our case: $\sigma = 2$ and $n = 25$ resulted in $c = 0.658$. Power needs a μ under alternate hypothesis. If we assume $\mu = 1$,

$$\text{Power : } Pr [\bar{X} > 0.658 | \mu = 1]$$

$$\text{Power}(1) : Pr \left[Z > \frac{\sqrt{n}}{\sigma} (c - 1) \right] \quad Z \sim N(0, 1)$$

Hypothesis testing framework

The right hand side quantity

$$\frac{\sqrt{n}}{\sigma}(c - 1) = \frac{\sqrt{n}}{\sigma}c - \frac{\sqrt{n}}{\sigma}$$

Hypothesis testing framework

The right hand side quantity

$$\frac{\sqrt{n}}{\sigma}(c - 1) = \frac{\sqrt{n}}{\sigma}c - \frac{\sqrt{n}}{\sigma}$$

here at 5% level of significance,

$$\frac{\sqrt{n}}{\sigma}(c - 1) = 1.644 - \frac{\sqrt{n}}{\sigma}$$

$$\text{Power : } Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma} \right] \quad Z \sim N(0, 1)$$

Hypothesis testing framework

The right hand side quantity

$$\frac{\sqrt{n}}{\sigma}(c - 1) = \frac{\sqrt{n}}{\sigma}c - \frac{\sqrt{n}}{\sigma}$$

here at 5% level of significance,

$$\frac{\sqrt{n}}{\sigma}(c - 1) = 1.644 - \frac{\sqrt{n}}{\sigma}$$

$$\text{Power} : Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma} \right] \quad Z \sim N(0, 1)$$

As σ decreases or n increases, we have higher power.

Hypothesis testing framework

The right hand side quantity

$$\frac{\sqrt{n}}{\sigma}(c - 1) = \frac{\sqrt{n}}{\sigma}c - \frac{\sqrt{n}}{\sigma}$$

here at 5% level of significance,

$$\frac{\sqrt{n}}{\sigma}(c - 1) = 1.644 - \frac{\sqrt{n}}{\sigma}$$

$$\text{Power} : Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma} \right] \quad Z \sim N(0, 1)$$

As σ decreases or n increases, we have higher power.

But we do not control σ , we can only control n .

Hypothesis testing framework

The right hand side quantity

$$\frac{\sqrt{n}}{\sigma}(c - 1) = \frac{\sqrt{n}}{\sigma}c - \frac{\sqrt{n}}{\sigma}$$

here at 5% level of significance,

$$\frac{\sqrt{n}}{\sigma}(c - 1) = 1.644 - \frac{\sqrt{n}}{\sigma}$$

$$\text{Power} : Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma} \right] \quad Z \sim N(0, 1)$$

As σ decreases or n increases, we have higher power.

But we do not control σ , we can only control n .

Hypothesis testing framework

Suppose I say I want my power to be 0.99 at 5% level of significance.

Hypothesis testing framework

Suppose I say I want my power to be 0.99 at 5% level of significance.

We know that

$$\text{Type I error : } Pr \left[Z > \frac{\sqrt{n}}{\sigma} c \mid \mu = 0 \right] = 0.05 \quad Z \sim N(0, 1)$$

Hypothesis testing framework

Suppose I say I want my power to be 0.99 at 5% level of significance.

We know that

$$\text{Type I error : } Pr \left[Z > \frac{\sqrt{n}}{\sigma} c \mid \mu = 0 \right] = 0.05 \quad Z \sim N(0, 1)$$

From this, we know

$$\frac{\sqrt{n}}{\sigma} c = 1.644$$

Hypothesis testing framework

Suppose I say I want my power to be 0.99 at 5% level of significance.

We know that

$$\text{Type I error : } Pr \left[Z > \frac{\sqrt{n}}{\sigma} c \mid \mu = 0 \right] = 0.05 \quad Z \sim N(0, 1)$$

From this, we know

$$\frac{\sqrt{n}}{\sigma} c = 1.644$$

and

$$\text{Power : } Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma} \right] \quad Z \sim N(0, 1)$$

Hypothesis testing framework

$$1.644 - \frac{\sqrt{n}}{2} = -2.32$$

Hypothesis testing framework

$$1.644 - \frac{\sqrt{n}}{2} = -2.32$$

$$\sqrt{n} = 2 * (2.32 + 1.644) = 7.928$$

Hypothesis testing framework

$$1.644 - \frac{\sqrt{n}}{2} = -2.32$$

$$\sqrt{n} = 2 * (2.32 + 1.644) = 7.928$$

$$n \approx 65$$

Hypothesis testing framework

$$1.644 - \frac{\sqrt{n}}{2} = -2.32$$

$$\sqrt{n} = 2 * (2.32 + 1.644) = 7.928$$

$$n \approx 65$$

So just collecting 65 samples under this testing framework we have can give us a Power of 0.99 at $\mu = 1$ or Type II error as low as 0.01 given that we fix the Type I error to be 0.05.

Hypothesis testing framework

$$1.644 - \frac{\sqrt{n}}{2} = -2.32$$

$$\sqrt{n} = 2 * (2.32 + 1.644) = 7.928$$

$$n \approx 65$$

So just collecting 65 samples under this testing framework we have can give us a Power of 0.99 at $\mu = 1$ or Type II error as low as 0.01 given that we fix the Type I error to be 0.05.

So, we can decide on how many samples to choose by deciding on the Type I and Type II errors we want.

Hypothesis testing framework

But $\mu = 1$ is only one possible alternative for the hypothesis. So, what we would want to do is get a formula of power for each value of μ in the alternate space.

$$Power(\mu) : Pr \left[Z > \frac{\sqrt{n}}{\sigma}(c - \mu) \right] \quad Z \sim N(0, 1)$$

Since from Type I error argument, we showed that

$$\frac{\sqrt{n}}{\sigma}c = 1.644$$


$$Power(\mu) : Pr \left[Z > 1.644 - \frac{\sqrt{n}}{\sigma}\mu \right] \quad Z \sim N(0, 1)$$

This can be written as

$$Power(\mu) := \Phi \left(\frac{\sqrt{n}}{\sigma}\mu - 0.644 \right)$$


Hypothesis testing framework

We plot $\text{Power}(\mu)$ against μ . Say $n = 65$ and we know $\sigma = 2$.




Hypothesis testing framework

We plot $\text{Power}(\mu)$ against μ . Say $n = 25$ and we know $\sigma = 2$.



Hypothesis testing framework

We plot $\text{Power}(\mu)$ against μ . Say $n = 25$ and we know $\sigma = 0.1$.



Hypothesis testing framework

- ▶ As σ decreases (for fixed n), the power curve becomes steeper and reaches 1 early (Type II error decreases steeply to 0).

Hypothesis testing framework

- ▶ As σ decreases (for fixed n), the power curve becomes steeper and reaches 1 early (Type II error decreases steeply to 0).
- ▶ As n increases (for fixed σ), the power curve becomes steeper and reaches 1 early. As n decreases, the power curve takes more time to reach 1.

Hypothesis testing framework

- ▶ As σ decreases (for fixed n), the power curve becomes steeper and reaches 1 early (Type II error decreases steeply to 0).
- ▶ As n increases (for fixed σ), the power curve becomes steeper and reaches 1 early. As n decreases, the power curve takes more time to reach 1.
- ▶ for fixed σ , the power for higher n remains higher at each μ . For fixed n , the power for lower σ remains higher at each μ

Another important continuous distribution

Chi-Square Distribution

The chi-square distributions: χ_k^2

$$f(x) = \frac{x^{(k/2)-1} e^{-x/2}}{2^{k/2} \Gamma(k/2)} \quad x > 0, k \geq 1$$

Chi-Square Distribution

The chi-square distributions: χ_k^2

$$f(x) = \frac{x^{(k/2)-1} e^{-x/2}}{2^{k/2} \Gamma(k/2)} \quad x > 0, k \geq 1$$


$$E(X) = k \quad \text{Var}(X) = 2k$$

Chi-Square Distribution

The chi-square distributions: χ_k^2

$$f(x) = \frac{x^{(k/2)-1} e^{-x/2}}{2^{k/2} \Gamma(k/2)} \quad x > 0, k \geq 1$$

$$E(X) = k \quad \text{Var}(X) = 2k$$



Chi-Square Distribution

The chi-square distributions: χ_1^2

$$f(x) = \frac{x^{-1/2}e^{-x/2}}{\sqrt{2\pi}} \quad x > 0$$

Chi-Square Distribution

The chi-square distributions: χ_1^2

$$f(x) = \frac{x^{-1/2}e^{-x/2}}{\sqrt{2\pi}} \quad x > 0$$

Origin of the Chi-square distribution:

Let $Z \sim N(0, 1)$ and define $Y = Z^2$. Then, $Y \sim \chi_1^2$.

Chi-Square Distribution

The chi-square distributions: χ_1^2

$$f(x) = \frac{x^{-1/2}e^{-x/2}}{\sqrt{2\pi}} \quad x > 0$$

Origin of the Chi-square distribution:

Let $Z \sim N(0, 1)$ and define $Y = Z^2$. Then, $Y \sim \chi_1^2$.

In general, χ_n^2 is a sum of n i.i.d. standard normal distributions.

Chi-Square Distribution

Properties:

- ▶ We will need the **reproductive property of χ^2** :

Chi-Square Distribution

Properties:

- ▶ We will need the **reproductive property of χ^2** :

Let $X_1 \sim \chi_{n_1}^2, X_2 \sim \chi_{n_2}^2, \dots, X_k \sim \chi_{n_k}^2$ with X_1, X_2, \dots, X_k independent random variables. Define

$Y = X_1 + X_2 + \dots + X_k$. Then, $Y \sim \chi_n^2$ where $n = n_1 + n_2 + \dots + n_k$.

Chi-Square Distribution

Properties:

- ▶ We will need the **reproductive property of χ^2** :

Let $X_1 \sim \chi_{n_1}^2, X_2 \sim \chi_{n_2}^2, \dots, X_k \sim \chi_{n_k}^2$ with X_1, X_2, \dots, X_k independent random variables. Define

$Y = X_1 + X_2 + \dots + X_k$. Then, $Y \sim \chi_n^2$ where
 $n = n_1 + n_2 + \dots + n_k$.

- ▶ Conversely, $X_1 \sim \chi_{n_1}^2, X_1 + X_2 \sim \chi_{n_1+n_2}^2 \Rightarrow X_2 \sim \chi_{n_2}^2$.

Chi-Square Distribution

Properties:

- ▶ We will need the **reproductive property of χ^2** :

Let $X_1 \sim \chi_{n_1}^2, X_2 \sim \chi_{n_2}^2, \dots, X_k \sim \chi_{n_k}^2$ with X_1, X_2, \dots, X_k independent random variables. Define

$Y = X_1 + X_2 + \dots + X_k$. Then, $Y \sim \chi_n^2$ where $n = n_1 + n_2 + \dots + n_k$.

- ▶ Conversely, $X_1 \sim \chi_{n_1}^2, X_1 + X_2 \sim \chi_{n_1+n_2}^2 \Rightarrow X_2 \sim \chi_{n_2}^2$.
- ▶ If $X_i \sim N()$, then \bar{X} and S^2 are independent.

Chi-Square Distribution

Properties:

- ▶ We will need the **reproductive property of χ^2** :

Let $X_1 \sim \chi_{n_1}^2, X_2 \sim \chi_{n_2}^2, \dots, X_k \sim \chi_{n_k}^2$ with X_1, X_2, \dots, X_k independent random variables. Define

$Y = X_1 + X_2 + \dots + X_k$. Then, $Y \sim \chi_n^2$ where $n = n_1 + n_2 + \dots + n_k$.

- ▶ Conversely, $X_1 \sim \chi_{n_1}^2, X_1 + X_2 \sim \chi_{n_1+n_2}^2 \Rightarrow X_2 \sim \chi_{n_2}^2$.
- ▶ If $X_i \sim N()$, then \bar{X} and S^2 are independent.
Reason: $\text{Cov}(\bar{X}, X_i - \bar{X}) = 0 \Rightarrow \bar{X}$ and $X_i - \bar{X}$ are uncorrelated
 $\Rightarrow \bar{X}$ and $X_i - \bar{X}$ are independent.

Chi-Square Distribution

To get started, consider a random sample X_1, X_2, \dots, X_n from a $N(\mu, \sigma^2)$ population.

Chi-Square Distribution

To get started, consider a random sample X_1, X_2, \dots, X_n from a $N(\mu, \sigma^2)$ population. Let

$$Z_i = \frac{X_i - \mu}{\sigma} \quad \text{so that} \quad Z_i^2 \sim \chi_1^2 \quad \text{and } Z_i' \text{'s are independent.}$$

Chi-Square Distribution

To get started, consider a random sample X_1, X_2, \dots, X_n from a $N(\mu, \sigma^2)$ population. Let

$$Z_i = \frac{X_i - \mu}{\sigma} \quad \text{so that} \quad Z_i^2 \sim \chi_1^2 \quad \text{and } Z_i' \text{'s are independent.}$$

Then,

$$\sum_{i=1}^n \left[\frac{X_i - \mu}{\sigma} \right]^2 = \sum_{i=1}^n Z_i^2 = \sum_{i=1}^n \chi_1^2 = \chi_n^2.$$

Chi-Square Distribution

To get started, consider a random sample X_1, X_2, \dots, X_n from a $N(\mu, \sigma^2)$ population. Let

$$Z_i = \frac{X_i - \mu}{\sigma} \quad \text{so that} \quad Z_i^2 \sim \chi_1^2 \quad \text{and } Z_i' \text{'s are independent.}$$

Then,

$$\sum_{i=1}^n \left[\frac{X_i - \mu}{\sigma} \right]^2 = \sum_{i=1}^n Z_i^2 = \sum_{i=1}^n \chi_1^2 = \chi_n^2.$$

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{(n-1)}^2$$

Mean of Sampling Distribution of s^2

Since we have,

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{(n-1)}$$

for a random sample from a $N(\mu, \sigma^2)$ population, we have that

$$E \left[\frac{(n-1)S^2}{\sigma^2} \right] = E(\chi^2_{(n-1)}) = n - 1$$

Mean of Sampling Distribution of s^2

Since we have,

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{(n-1)}$$

for a random sample from a $N(\mu, \sigma^2)$ population, we have that

$$E\left[\frac{(n-1)S^2}{\sigma^2}\right] = E(\chi^2_{(n-1)}) = n-1$$

It follows that,

$$n-1 = E\left[\frac{(n-1)S^2}{\sigma^2}\right] = \frac{(n-1)}{\sigma^2} E(S^2) \Rightarrow E(s^2) = \sigma^2.$$

So, when sampling from a normal population, s^2 is on target (σ^2) on average.

Mean of Sampling Distribution of s^2

What is the mean of the sampling distribution for S^2 for samples from *any* population?

Mean of Sampling Distribution of s^2

What is the mean of the sampling distribution for S^2 for samples from *any* population?

First, show that $E \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] = (n-1)\sigma^2$

Mean of Sampling Distribution of s^2

What is the mean of the sampling distribution for S^2 for samples from *any* population?

First, show that $E \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] = (n-1)\sigma^2$

Then, we can conclude that

$$E(S^2) = E \left[\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \sigma^2$$

for samples from *any* population.

Mean of Sampling Distribution of s^2

What is the mean of the sampling distribution for S^2 for samples from *any* population?

First, show that $E \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] = (n-1)\sigma^2$

Then, we can conclude that

$$E(S^2) = E \left[\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \sigma^2$$

for samples from *any* population.

This is one reason why sample variance is defined by dividing by $n-1$ instead of n . If we divide by n , the resulting sample variance will tend to underestimate σ^2 on average.

Mean of Sampling Distribution of s^2

What is the mean of the sampling distribution for S^2 for samples from *any* population?

First, show that $E \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] = (n-1)\sigma^2$

Then, we can conclude that

$$E(S^2) = E \left[\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \sigma^2$$

for samples from *any* population.

This is one reason why sample variance is defined by dividing by $n-1$ instead of n . If we divide by n , the resulting sample variance will tend to underestimate σ^2 on average.

The “Student’s” T Probability Distribution

Let $Z \sim N(0, 1)$ and $Y \sim \chi^2_\nu$ with Z, Y **independent**. Then,
 $T = \frac{Z}{\sqrt{Y/\nu}}$ is said to have the T -distribution with ν degrees of freedom (df): $T \sim t_n$.

The “Student’s” T Probability Distribution

Let $Z \sim N(0, 1)$ and $Y \sim \chi^2_\nu$ with Z, Y **independent**. Then, $T = \frac{Z}{\sqrt{Y/\nu}}$ is said to have the T -distribution with ν degrees of freedom (df): $T \sim t_n$. p.d.f:

$$f(t) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\pi\nu}\Gamma(\frac{\nu}{2})} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}, -\infty < t < \infty$$

The “Student’s” T Probability Distribution

Let $Z \sim N(0, 1)$ and $Y \sim \chi^2_\nu$ with Z, Y **independent**. Then, $T = \frac{Z}{\sqrt{Y/\nu}}$ is said to have the T -distribution with ν degrees of freedom (df): $T \sim t_n$. p.d.f:

$$f(t) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\pi\nu}\Gamma(\frac{\nu}{2})} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}, -\infty < t < \infty$$


It can be shown that $E(T) = 0$ and $Var(T) = \nu/(\nu - 2)$. $\nu > 2$

The “Student’s” T Probability Distribution

Let $Z \sim N(0, 1)$ and $Y \sim \chi^2_\nu$ with Z, Y **independent**. Then, $T = \frac{Z}{\sqrt{Y/\nu}}$ is said to have the T -distribution with ν degrees of freedom (df): $T \sim t_n$. p.d.f:

$$f(t) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\pi\nu}\Gamma(\frac{\nu}{2})} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}, -\infty < t < \infty$$

It can be shown that $E(T) = 0$ and $Var(T) = \nu/(\nu - 2)$. $\nu > 2$




The “Student’s” T Probability Distribution

Let $Z \sim N(0, 1)$ and $Y \sim \chi^2_\nu$ with Z, Y **independent**. Then, $T = \frac{Z}{\sqrt{Y/\nu}}$ is said to have the T -distribution with ν degrees of freedom (df): $T \sim t_n$. p.d.f:

$$f(t) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\pi\nu}\Gamma(\frac{\nu}{2})} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}, -\infty < t < \infty$$

It can be shown that $E(T) = 0$ and $Var(T) = \nu/(\nu - 2)$. $\nu > 2$



Question: What's the T distribution when $\nu \rightarrow \infty$?

How is the T -distribution Related to Estimation?

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ popn.

We know that $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$.

How is the T -distribution Related to Estimation?

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ popn.

We know that $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$.

But what if σ^2 is not known?

How is the T -distribution Related to Estimation?

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ popn.

We know that $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$.

But what if σ^2 is not known? Use $s \approx \sigma$ to estimate it.

Consider

$$\begin{aligned} T &= \frac{\bar{X} - \mu}{s/\sqrt{n}} = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{s} = Z \frac{\sigma}{s} \quad (\text{where } Z \sim N(0, 1)) \\ &= \frac{Z}{s/\sigma} = \frac{Z}{\sqrt{s^2/\sigma^2}} \\ &= \frac{Z}{\sqrt{\frac{(n-1)s^2}{\sigma^2(n-1)}}} = \frac{Z}{\sqrt{\chi_{(n-1)}^2/(n-1)}} \sim t_{(n-1)} \end{aligned}$$

How is the T -distribution Related to Estimation?

Let X_1, X_2, \dots, X_n be a random sample from a $N(\mu, \sigma^2)$ popn.

We know that $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$.

But what if σ^2 is not known? Use $s \approx \sigma$ to estimate it.

Consider

$$\begin{aligned} T &= \frac{\bar{X} - \mu}{s/\sqrt{n}} = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{s} = Z \frac{\sigma}{s} \quad (\text{where } Z \sim N(0, 1)) \\ &= \frac{Z}{s/\sigma} = \frac{Z}{\sqrt{s^2/\sigma^2}} \\ &= \frac{Z}{\sqrt{\frac{(n-1)s^2}{\sigma^2(n-1)}}} = \frac{Z}{\sqrt{\chi_{(n-1)}^2/(n-1)}} \sim t_{(n-1)} \end{aligned}$$

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

So, even when the popn variance σ^2 is not known, we can still find probabilities for the sample mean \bar{X} for data from a normal popn.

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

So, even when the popn variance σ^2 is not known, we can still find probabilities for the sample mean \bar{X} for data from a normal popn.

We just replace σ by its estimate (s) in the standardization of \bar{X} and then look up probabilities from the $t_{(n-1)}$ distribution instead of the $N(0, 1)$ distribution.

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

So, even when the popn variance σ^2 is not known, we can still find probabilities for the sample mean \bar{X} for data from a normal popn.

We just replace σ by its estimate (s) in the standardization of \bar{X} and then look up probabilities from the $t_{(n-1)}$ distribution instead of the $N(0, 1)$ distribution.

The $t_{(n-1)}$ distribution is similar to the $N(0, 1)$, but with more spread (to account for the extra uncertainty in the estimate of σ .)

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

So, even when the popn variance σ^2 is not known, we can still find probabilities for the sample mean \bar{X} for data from a normal popn.

We just replace σ by its estimate (s) in the standardization of \bar{X} and then look up probabilities from the $t_{(n-1)}$ distribution instead of the $N(0, 1)$ distribution.

The $t_{(n-1)}$ distribution is similar to the $N(0, 1)$, but with more spread (to account for the extra uncertainty in the estimate of σ .)

As sample size increases, s gets closer to σ and the $t_{(n-1)}$ distribution gets closer to $N(0, 1)$.

How is the T -distribution Related to Estimation?

$$T = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{(n-1)}$$

So, even when the popn variance σ^2 is not known, we can still find probabilities for the sample mean \bar{X} for data from a normal popn.

We just replace σ by its estimate (s) in the standardization of \bar{X} and then look up probabilities from the $t_{(n-1)}$ distribution instead of the $N(0, 1)$ distribution.

The $t_{(n-1)}$ distribution is similar to the $N(0, 1)$, but with more spread (to account for the extra uncertainty in the estimate of σ .)

As sample size increases, s gets closer to σ and the $t_{(n-1)}$ distribution gets closer to $N(0, 1)$.