



4503 - Integer Game

Asia - Dhaka - 2009/2010

You are playing a single player game where you can convert one integer from another through a sequence of moves. The integer Y is reachable from X in a single move if the following is satisfied.

$$Y = \frac{X \times P_2^k}{P_1^k}, \text{ where } k \text{ is a positive integer, } P_1 \text{ and } P_2 \text{ are prime numbers and } X \text{ is divisible by } P_1^k.$$

For example 18 is reachable from 8 in one move, because you can divide 8 by 4 and then multiply by 9. But 6 is not reachable from 8. Given two integers A and B calculate the minimum number of moves necessary to transform A into B. Both A and B can be very large. So each of them is needed to be expressed as a

multiplication of a sequence of small integers: $A = \prod_{i=1}^N a_i$ and $B = \prod_{i=1}^M b_i$. Both of the sequences a_i and b_i will be given as inputs.

Input

First line of the input contains T ($1 \leq T \leq 40$) the number of test cases. Then T blocks of test cases follows. First line of the test case contains N ($1 \leq N \leq 300$) followed by N integers. N is the length of the sequence a_i and the following N integers form the sequence a_i . The second line starts with an integer M ($1 \leq M \leq 300$). M is the length of the sequence b_i and the following M integers form the sequence b_i . Each of integers in these two sequences will be between 2 and 200 (inclusive).

Output

For each case of input, print the serial of output followed by an integer: the minimum number of moves required to transform A to B. If it is impossible to transform A to B with any number of moves output -1 instead. If the minimum number of moves is greater than or equal to 20 print a -1 as well.

Sample Input Input

4

Case 1: 1

Output for Sample

1 4

Case 2: 1

1 9

Case 3: -1

2 2 2

Case 4: 3

2 3 3

1 8

1 6

2 3 2 1 1

3 2 7 2 5 1 3

Dhaka 2009-2010

Problemsetter: Abdullah al Mahmud

Special Thanks: Md. Arifuzzaman Arif