

Department of Physics, NTNU

Homework 6 TFY4210/FY8302 Quantum theory of solids Spring 2021.

Problem 1

In this problem, we will consider a system defined by a Hamiltonian

$$H = H_0 + V = \sum_{k,\sigma} \varepsilon_k c_{k,\sigma}^\dagger c_{k,\sigma} + V$$

where V is a two-body interaction term that renders the problem not exactly solvable. In class, we have seen how we can express the single-particle electron Green's function for such a system as

$$\begin{aligned} G(k, t - t') &= -i \langle \Psi(0) | T \left[\hat{c}_k(t) \hat{c}_k^\dagger(t') \right] | \Psi(0) \rangle \\ &= -i \frac{\langle \phi_0 | T \left[c_k(t) c_k^\dagger(t') S(\infty, -\infty) \right] | \phi_0 \rangle}{\langle \phi_0 | S(\infty, -\infty) | \phi_0 \rangle} \end{aligned}$$

where $S(\infty, -\infty)$ is given by

$$S(\infty, -\infty) = 1 + \sum_{n=1}^{\infty} \frac{(-i)^n}{n!} \int_{-\infty}^{\infty} dt_1 \cdots \int_{-\infty}^{\infty} dt_n T[V(t_1) \cdots V(t_n)]$$

and T is the time-ordering operator. The notation is otherwise the same as we have used in class. Assume now that the perturbation is given by the Hubbard-interaction

$$V = U \sum_i n_{i\uparrow} n_{i\downarrow} = \frac{U}{2} \sum_{i,\sigma} n_{i\sigma} n_{i-\sigma}$$

a) Write the Hubbard-interaction on the form

$$V = \sum_{k,k',q,\sigma,\sigma'} \tilde{V}(q, \sigma, \sigma') c_{k+q,\sigma}^\dagger c_{k'-q,\sigma'}^\dagger c_{k',\sigma'} c_{k,\sigma}$$

thereby specifying $\tilde{V}(q, \sigma, \sigma')$. (Note: this interaction is spin-dependent, contrary to what the case is for the standard density-density Coulomb-interaction or the effective electron-electron interaction mediated by phonons.)

b) Use the resulting interaction and calculate the leading order correction of the denominator in the second expression for $G(k, t - t')$, and give a diagrammatic representation for it along the same lines that we used for the electron-phonon coupling in class.

c) Calculate the leading order correction of the numerator in the second expression for $G(k, t - t')$, and give a diagrammatic representation for it along the same lines that we used for the electron-phonon coupling in class.

d) Show that, to leading order in V , the denominator cancels the disconnected diagrams appearing in the numerator.