





EEG beta power during planning predicts motor sequence initiation time

Martin Geiger¹, Atif Shahzad^{2,3}, Fei He⁴, Katja Kornysheva^{1,5}

¹Centre for Human Brain Health, School of Psychology, University of Birmingham; ²Smart Sensor Lab, Lambe Institute of Translational Research, College of Medicine, Nursing Health Sciences, University of Galway; ³Centre for Systems Modelling and Quantitative Biomedicine, University of Birmingham; ⁴Research Centre for Computational Science and Mathematical Modelling, Coventry University; ⁵Bangor Imaging Unit, Bangor University

Background

- Parkinson’s Disease (PD) patients experience impaired movement sequence initiation and bradykinesia (He et al., 2020).
- In their electroencephalogram (EEG), beta band event-related desynchronization (ERD) is typically reduced during movement (Mehler, 2022).
- We investigate how ERD preceding movement onset (motor planning) is linked to subsequent motor performance to provide targets for brain-computer interface (BCI) neurofeedback.



Conclusions

- Single-trial low beta power (13-20 Hz) is modulated during planning of motor sequence and significantly predicts behaviour
- ➔ We are implementing a neurofeedback BCI to train PD patients to voluntarily enhance beta ERD during planning to facilitate motor sequence execution.

References

He, S., Everest-Phillips, C., Clouter, A., Brown, P., & Tan, H. (2020). Neurofeedback-Linked Suppression of Cortical β Bursts Speeds Up Movement Initiation in Healthy Motor Control: A Double-Blind Sham-Controlled Study. *Journal of Neuroscience*, 40(20), 4021–4032. <https://doi.org/10.1523/JNEUROSCI.0208-20.2020>

Mehler, D. M. A. (2022). Turning markers into targets – scoping neural circuits for motor neurofeedback training in Parkinson’s disease. *Brain-Apparatus Communication: A Journal of Bacomics*, 1(1), 1–27. <https://doi.org/10.1080/27706710.2022.2061300>

