

# CLINICAL STUDY REPORT

**Name of Sponsor/Company:** NovaCura Biotech

**Name of Finished Product:** CLN-201

**Name of Active Ingredient:** CLN-201

**Study Title:** A Phase II, Randomized, Double-Blind, Placebo-Controlled, Multicenter Study to Evaluate the Efficacy, Safety, and Tolerability of CLN-201 in Adult Participants With Moderate Inflammatory Disease

**Protocol Number:** CLN-201-2026-01

**Development Phase:** Phase II

**Report Version:** 1.0

**Report Date:** 2026-05-16

**Confidentiality Statement:** This clinical study report contains confidential information of NovaCura Biotech. This document is intended for regulatory and authorized professional review only.

*ICH E3 note: A final CSR should also include names/titles of sponsor signatories, medical expert responsible for the report, and where applicable, investigator and coordinating investigator identification.*

## Table of Contents

---

1. Synopsis
2. Table of Contents
3. List of Abbreviations and Definition of Terms
4. Ethics
5. Investigators and Administrative Structure
6. Introduction
7. Study Objectives
8. Investigational Plan
  1. Overall Study Design and Plan
  2. Discussion of Study Design, Including Choice of Control Groups
  3. Selection of Study Population
  4. Treatments

5. Efficacy and Safety Variables
6. Data Quality Assurance
7. Statistical Methods Planned in the Protocol and Determination of Sample Size
8. Changes in the Conduct of the Study or Planned Analyses
9. Study Patients
  1. Disposition of Patients
  2. Protocol Deviations
10. Efficacy Evaluation
  1. Data Sets Analyzed
  2. Demographic and Other Baseline Characteristics
  3. Measurements of Treatment Compliance
  4. Efficacy Results and Tabulations
11. Safety Evaluation
  1. Extent of Exposure
  2. Adverse Events
  3. Deaths, Other Serious Adverse Events, and Other Significant Adverse Events
  4. Clinical Laboratory Evaluation
  5. Vital Signs, Physical Findings, and Other Observations Related to Safety
12. Discussion and Overall Conclusions
13. Tables, Figures, and Graphs Referred to but Not Included in the Text
14. Reference List
15. Appendices

## List of Abbreviations and Definition of Terms

| Abbreviation | Definition                                                |
|--------------|-----------------------------------------------------------|
| AE           | Adverse Event                                             |
| CSR          | Clinical Study Report                                     |
| GCP          | Good Clinical Practice                                    |
| ICH          | International Council for Harmonisation                   |
| IRB/IEC      | Institutional Review Board / Independent Ethics Committee |
| SAE          | Serious Adverse Event                                     |
| SAP          | Statistical Analysis Plan                                 |

## 1. Synopsis

| Item                    | Description                                                                                                                                                                                        |
|-------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Name of Sponsor/Company | NovaCura Biotech                                                                                                                                                                                   |
| Study Title             | A Phase II, Randomized, Double-Blind, Placebo-Controlled, Multicenter Study to Evaluate the Efficacy, Safety, and Tolerability of CLN-201 in Adult Participants With Moderate Inflammatory Disease |
| Protocol Number         | CLN-201-2026-01                                                                                                                                                                                    |
| Study Phase             | Phase II                                                                                                                                                                                           |
| Study Design            | Randomized, double-blind, placebo-controlled, parallel-group, multicenter study.                                                                                                                   |
| Study Population        | Adults 18 to 75 years of age with moderate inflammatory disease and stable background therapy where applicable.                                                                                    |
| Participants            | Screened: 148; Randomized: 120; Completed treatment: 110; Discontinued early: 10.                                                                                                                  |
| Treatment Groups        | CLN-201 once daily; matching placebo once daily.                                                                                                                                                   |
| Study Duration          | Screening up to 21 days; Treatment 12 weeks; Follow-up 4 weeks.                                                                                                                                    |
| Primary Objective       | To evaluate the efficacy of CLN-201 compared with placebo in reducing disease activity from baseline to Week 12.                                                                                   |
| Primary Endpoint        | Change from baseline to Week 12 in disease activity score.                                                                                                                                         |
| Efficacy Summary        | CLN-201 demonstrated greater improvement than placebo in the primary endpoint, with favorable trends in secondary endpoints.                                                                       |
| Safety Summary          | CLN-201 was generally well tolerated. No new major safety signals were identified.                                                                                                                 |
| Conclusion              | The study met its primary objective and supports continued clinical development of CLN-201.                                                                                                        |

## 2. Ethics

The study was conducted in accordance with the ethical principles originating in the Declaration of Helsinki, ICH Good Clinical Practice, and applicable local regulatory requirements.

Approval from the relevant Institutional Review Board or Independent Ethics Committee was obtained before initiation of study-specific procedures at each participating site.

Written informed consent was obtained from each participant before the conduct of any study-specific procedures.

*Add study-specific IRB/IEC names, approval dates, and whether any protocol amendments required re-approval.*

### 3. Investigators and Administrative Structure

---

This multicenter study was conducted at qualified investigational sites under the supervision of appropriately trained investigators.

*Insert list of investigators, study centers, coordinating investigator (if applicable), contract research organization, central laboratory, bioanalytical laboratory, statistical support provider, and any steering/data monitoring committees.*

### 4. Introduction

---

CLN-201 is an investigational product under development for the treatment of inflammatory disease. This study was designed to evaluate the efficacy, safety, and tolerability of CLN-201 in adult participants with moderate inflammatory disease.

The rationale for the study was based on the need for improved therapeutic options and prior evidence suggesting potential clinical benefit of CLN-201 in reducing disease activity.

*Add background nonclinical and clinical experience, mechanism of action, and rationale for selected dose, duration, and endpoint strategy.*

### 5. Study Objectives

---

#### 5.1 Primary Objective

To evaluate the efficacy of CLN-201 compared with placebo in reducing disease activity from baseline to Week 12.

#### 5.2 Secondary Objectives

- To further assess the effect of CLN-201 on symptoms of inflammatory disease.

- To evaluate the effect of CLN-201 on inflammatory biomarkers, including C-reactive protein.
- To assess quality-of-life outcomes and clinical response criteria.
- To evaluate the safety and tolerability of CLN-201.

## 6. Investigational Plan

---

### 6.1 Overall Study Design and Plan

This was a randomized, double-blind, placebo-controlled, parallel-group, multicenter Phase II study. Eligible participants were randomized in a 1:1 ratio to receive CLN-201 or matching placebo orally once daily for 12 weeks.

Assessments were performed during screening, baseline, Weeks 2, 4, 8, and 12, and at follow-up approximately 4 weeks after the last dose.

### 6.2 Discussion of Study Design, Including Choice of Control Groups

A placebo-controlled design was selected to provide an unbiased assessment of the efficacy and safety of CLN-201 while minimizing confounding influences. Blinding and randomization were used to reduce bias in efficacy and safety assessments.

### 6.3 Selection of Study Population

#### *Inclusion Criteria*

- Adults 18 to 75 years of age.
- Diagnosis of moderate inflammatory disease as defined in the protocol.
- Stable background therapy where applicable.
- Able and willing to provide written informed consent.

#### *Exclusion Criteria*

- Any clinically significant condition that could interfere with study participation or interpretation of results.
- Relevant uncontrolled comorbidities or active infection.
- Other protocol-defined exclusions.

*Replace with exact protocol-defined inclusion/exclusion criteria and specify withdrawal/discontinuation criteria.*

### 6.4 Treatments

Participants received CLN-201 once daily or matching placebo once daily for 12 weeks.

*Add dosage strength, formulation, batch information if required, route of administration, packaging, storage, treatment accountability, randomization procedure, blinding maintenance, and rules for dose interruption/discontinuation.*

## 6.5 Efficacy and Safety Variables

### *Efficacy Variables*

- **Primary efficacy variable:** Change from baseline to Week 12 in disease activity score.
- **Secondary efficacy variables:** Symptom improvement, inflammatory biomarker changes, quality-of-life outcomes, and clinical response rates at Week 12.

### *Safety Variables*

- Adverse events and serious adverse events
- Clinical laboratory parameters
- Vital signs
- Physical examination findings
- Other protocol-specified safety assessments

## 6.6 Data Quality Assurance

Data were reviewed using standard quality control procedures to support completeness, consistency, and accuracy.

*Add monitoring approach, source data verification strategy, database quality control, coding dictionaries used, and audit/inspection information if applicable.*

## 6.7 Statistical Methods Planned in the Protocol and Determination of Sample Size

The primary analysis compared change from baseline to Week 12 in disease activity score between the CLN-201 and placebo groups.

*Add formal sample size justification, analysis populations (e.g., ITT, safety, per-protocol), statistical models, covariates, handling of missing data, multiplicity control, significance thresholds, and any interim analyses.*

## 6.8 Changes in the Conduct of the Study or Planned Analyses

No major deviations from the overall planned study conduct are described in this draft report.

*Add all protocol amendments, analysis changes, endpoint changes, or operational deviations from planned conduct.*

## 7. Study Patients

---

### 7.1 Disposition of Patients

A total of 148 participants were screened and 120 were randomized. Of the randomized participants, 110 completed treatment and 10 discontinued early.

| Disposition Category | Count |
|----------------------|-------|
| Screened             | 148   |
| Randomized           | 120   |
| Completed Treatment  | 110   |
| Discontinued Early   | 10    |

Reasons for early discontinuation included adverse events, withdrawal of consent, loss to follow-up, or investigator decision.

### 7.2 Protocol Deviations

*Insert summary of important protocol deviations and identify whether any participants were excluded from specific analysis sets.*

## 8. Efficacy Evaluation

---

### 8.1 Data Sets Analyzed

*Specify intent-to-treat/full analysis set, per-protocol set, safety set, and any other analysis populations with participant counts.*

### 8.2 Demographic and Other Baseline Characteristics

Demographic and baseline characteristics were generally balanced between treatment groups. The mean age was 46 years and 62% of participants were female.

*Add full baseline tables including disease severity, duration, region, race/ethnicity if collected, relevant baseline biomarkers, and prior/concomitant therapies.*

### 8.3 Measurements of Treatment Compliance

*Add treatment exposure/compliance summary, such as pill count compliance, treatment duration by arm, dose interruptions, and dose reductions if applicable.*

## 8.4 Efficacy Results and Tabulations

### ***Primary Efficacy Result***

At Week 12, participants treated with CLN-201 showed greater improvement in disease activity score than placebo.

| Measure                                                        | CLN-201 | Placebo |
|----------------------------------------------------------------|---------|---------|
| Mean change from baseline to Week 12 in disease activity score | -3.8    | -2.1    |

**Treatment difference:** -1.7 in favor of CLN-201.

### ***Secondary Efficacy Results***

- Greater symptom improvement with CLN-201
- Reduction in inflammatory biomarkers including C-reactive protein
- Improved quality-of-life outcomes
- Higher proportion meeting clinical response criteria at Week 12

*Add confidence intervals, p-values, model outputs, sensitivity analyses, subgroup analyses, and supporting tables/figures.*



## 9. Safety Evaluation

### 9.1 Extent of Exposure

*Add number of doses received, mean/median exposure duration, cumulative dose, and exposure summaries by treatment arm.*

### 9.2 Adverse Events

Overall, CLN-201 was generally well tolerated. Most adverse events were mild to moderate.

| Safety Measure                               | CLN-201 | Placebo |
|----------------------------------------------|---------|---------|
| Participants with at least one adverse event | 38%     | 31%     |
| Serious adverse events                       | 2%      | 1%      |

#### Most common adverse events:

- Headache
- Nausea
- Fatigue

### 9.3 Deaths, Other Serious Adverse Events, and Other Significant Adverse Events

No new major safety signals were identified.

*Explicitly state number of deaths, details of SAEs, discontinuations due to AEs, and clinically significant adverse events of special interest.*

### 9.4 Clinical Laboratory Evaluation

Routine laboratory tests did not reveal clinically meaningful treatment-related safety concerns.

*Add hematology, chemistry, urinalysis summaries, shift tables, notable abnormalities, and clinically significant findings.*

### 9.5 Vital Signs, Physical Findings, and Other Observations Related to Safety

Vital signs and physical examination findings did not reveal clinically meaningful treatment-related safety concerns.

*Add ECGs or other safety assessments if performed.*

## 10. Discussion and Overall Conclusions

---

This Phase II study met its primary objective. CLN-201 improved disease activity compared with placebo over 12 weeks. Secondary endpoints were generally supportive of benefit. The safety profile was acceptable and consistent with prior knowledge of the investigational product.

Overall, these findings support continued clinical development of CLN-201.

*Expand discussion to include benefit-risk interpretation, consistency across endpoints, limitations, protocol deviations affecting interpretation, and generalizability.*

## 11. Tables, Figures, and Graphs Referred to but Not Included in the Text

---

- Participant disposition figure
- Baseline characteristics table
- Primary endpoint statistical output table
- Secondary endpoint summary tables
- Treatment-emergent adverse event summary tables
- Laboratory summary tables

## 12. Reference List

---

*Insert scientific references, protocol references, statistical analysis plan references, and other cited documents.*

## 13. Appendices

---

- Study protocol and amendments
- Sample case report form
- List of investigators and study administrative structure
- Ethics committee information and representative written information for patient informed consent
- Statistical analysis plan
- Patient data listings
- Summary tables
- Safety listings

*For full ICH E3 compliance, appendices should include all required supporting documentation and data presentations referenced in the body of the CSR.*