



# Metric spaces

Kalle Kytölä

February 9, 2021



# Contents

Foreword	v
Symbols indicating material beyond core content:	v
Chapter I. Foundations: set theory and logic	1
I.1. Basics of set theory	2
Sets and elements	2
Subsets	4
Inverted relation symbols	5
Familiar sets of numbers	5
Intervals	6
Operations with sets	7
Cartesian products	9
I.2. Functions	10
Surjective, injective, and bijective functions	12
Compositions of functions	13
Restricting the domain	14
Images and preimages under functions	14
I.3. Logic and related notation	17
Quantifiers	17
Implications and equivalences	19
Negations	20
I.4. ♥ In depth topics about the set theory	23
Chapter II. Real numbers	25
II.1. Absolute value, distances between numbers, and triangle inequality	25
II.2. Sequences on the real line	27
Sequences of real numbers	27
Monotonicity properties of number sequences	27
Boundedness properties of number sequences	28
II.3. Limits of sequences on the real line	29
The definition of the limit of a sequence	29
Basic properties of limits of sequences	29
Rules of calculation with limits	33
II.4. Density of rational and irrational numbers	35
Density of rational numbers	35
Density of irrational numbers	35
II.5. Axioms of the real numbers	36
On the field axioms and order axioms	37
Completeness axiom	38
II.6. ♥ In depth topics about the real numbers	42
Chapter III. Sequences and functions on the real line	45
III.1. Real number sequences	45

Subsequences	45
III.2. Functions of real variable	48
Continuity of a function of real variable	48
Operations on continuous functions	51
III.3. Continuous functions on a closed interval	54
Chapter IV. Normed spaces and inner product spaces	59
IV.1. Vector spaces	59
Axioms of real vector spaces	60
First examples of real vector spaces	60
Vector subspaces	62
IV.2. Inner product spaces	64
Axioms of inner product spaces	64
Examples of inner product spaces	65
The norm induced by an inner product	67
Angles and orthogonality	69
IV.3. Normed spaces	71
Axioms of normed spaces	71
Examples of normed spaces	72
Chapter V. Metric spaces	77
V.1. Axioms of metric spaces	77
V.2. Examples of metric spaces	78
Metric inherited to subsets	80
V.3. First notions in metric spaces	82
Balls	82
Diameter and distance	83
V.4. Open and closed sets	85
Open sets	85
Open sets in subspaces of metric spaces	87
Closed sets	88
Warning: clopen sets!	89
V.5. ♥ Interior, exterior, and boundary	90
V.6. ♥ Closure	91
Chapter VI. Continuity of functions	93
VI.1. Definition of continuity	94
Lipschitz continuity	95
♥ Other quantitative notions of continuity	100
VI.2. Characterization of continuity	101
VI.3. Exercises	104
Chapter VII. Topology and homeomorphisms	107
VII.1. ♥ General topology	107
VII.2. Homeomorphism	107
Homeomorphism as mappings that preserve topology	110
VII.3. Equivalence of metrics	111
VII.4. Product spaces	112
Metrics on product space	114
Equivalence of the product space metrics	114
Projections	114

Chapter VIII. Sequences in metric spaces	117
VIII.1. Sequences and limits in metric spaces	117
VIII.2. Characterization of closure and closed sets	119
VIII.3. Characterization of continuity	121
Operations on continuous real-valued functions	123
VIII.4. Sequences in product spaces	123
Chapter IX. Function sequences	125
IX.1. Pointwise convergence and uniform convergence	125
IX.2. Preservation of continuity in uniform limits	130
IX.3. Function series	131
♥ Power series	131
♥ Fourier series	132
IX.4. ♥ Uniform convergence and integration	133
Chapter X. Completeness	135
X.1. Cauchy sequences	135
X.2. Completeness	138
X.2.1. Definition of completeness	138
X.2.2. Completeness of the real line	139
X.2.3. Important complete spaces	139
X.2.4. Completeness for subspaces	141
X.3. Banach's fixed point theorem	141
Iterating a self-map	141
The contraction mapping principle	143
Applications	144
X.4. ♥ Hilbert spaces and Banach spaces	147
Chapter XI. Compactness	149
XI.1. Sequential compactness	149
XI.1.1. Definition and examples of sequential compactness	149
XI.1.2. Compactness versus completeness	149
XI.1.3. Compactness and closed subsets	150
XI.1.4. Preservation of compactness under continuous mappings	150
XI.1.5. Continuous real valued functions on compact spaces	150
XI.2. Compact subsets in Euclidean spaces	150
XI.3. *Compactness of product spaces	150
XI.4. Covering compactness	151
XI.5. Uniform continuity	151
XI.5.1. *Application to Riemann-integrability	151
XI.5.2. Exercises	151
Chapter XII. Connectedness	153
XII.1. Definition and examples of connectedness	153
XII.1.1. Connected components	153
XII.2. Connected subsets of the real axis	153
XII.3. Preservation of connectedness under continuous mappings	154
XII.4. Path connectedness	154
XII.4.1. *Simply connectedness	155
Appendix A. Further topics in set theory	157
A.1. ♥ Cardinalities of sets	157

An observation about finite sets	157
Cardinality comparison and equal cardinalities	158
Countable and uncountable infinite sets	159
Appendix B. Further topics about real numbers	163
B.1. ♥ Field axioms and order axioms of the real numbers	163
♥ Field axioms	163
♥ Order axioms	165
B.2. ♥ Equivalence of the formulations of the completeness axiom	166
B.3. ♥ Cantor set	168
B.4. ♥ Applications of supremum and infimum	172
♥ Arc length of a curve	172
♥ Infinite sums with non-negative terms	173
♥ Riemann integration	174
Appendix. Index	177
Appendix. References	181
Bibliography	181

## Foreword

These lecture notes are primarily intended for the core B.Sc. level mathematics course **MS-E1541** *Metric Spaces* at Aalto University.

The structure of these notes is largely based on an earlier version of the course taught by Pekka Alestalo, and parts of the textbook [Väi99]. The notes are still in an entirely preliminary and incomplete form, and I plan to frequently update them during the current course. You could help me — and perhaps more importantly the students who will use this material — by sending comments about mistakes, misprints, needs for clarification, etc., to me via the course’s Zulip chat forum or by email ([kalle.kytola@aalto.fi](mailto:kalle.kytola@aalto.fi)).

As textbooks for the course, [Väi99] matches the contents and level very well, and is extremely well written, but it only exists in Finnish and Swedish. As English textbooks we recommend for example parts of [Car00] or [Rud76] (both of these books cover significantly more material than the present course).

## Symbols indicating material beyond core content:

### Section symbols

- ♥ stands for sections that are **optional** for the purposes of the present course; they are intended for those who prepare to study mathematics further or seek in depth understanding.

### Exercise symbols

- ✓ stands for exercises which should be quick, easy, and **routine**, in view of the preceding definitions and/or basic results.
- ⋯ stands for exercises whose solution can be **lengthy** in comparison to regular ones, but not necessarily more difficult.
- # stands for exercises that are more **challenging** than regular ones; they may require some creativity and a good command of various topics in basic mathematics, in addition to the newly introduced material.





## Lecture I

### Foundations: set theory and logic

#### *Study tips*

This chapter is long...Most of the material should, however, be familiar to you already. This more extensive background chapter is intended to serve two purposes:

- If there are things you are not yet familiar with, you can still catch up now and follow the rest of the course.
- We fix some notations and terminology that will be used in the subsequent chapters. If you are already very familiar with set theory and logic, perhaps just quickly check the terms in **bold** and the related notation.

Among the topics of this chapter, the images and preimages of sets under functions (the last part of Section I.2) is perhaps the only one likely to not have been covered in prior basic courses. It may correspondingly deserve some more attention.

The students planning further mathematical studies are advised to take this opportunity to look into countable and uncountable infinite sets (Appendix A.1), although this topic is not strictly necessary for the present course.

#### *Some perspectives on set theory before we start*

Mathematics, as it has been practiced especially in the 20th and 21st centuries, is (almost entirely) founded on the notions of *sets* and their *elements*. To just quickly illustrate the expressive power of set theory we mention that, e.g., the notion of a function can be reduced to that of sets<sup>1</sup>, and constructions of real numbers<sup>2</sup> and even of natural numbers<sup>3</sup> can be given based on sets. Such flexibility of set theory justifies taking it as the foundations of mathematics.

Historical reasons for looking for solid foundations of this type include the discovery of numerous apparent paradoxes, questions about the validity of certain arguments about numbers and functions, and even disagreements about what can be considered legitimate topics of mathematics in the first place. The late 19th century and early 20th century featured particularly notable controversies of this kind, and lead to the adoption of set theory as the (more or less) universally agreed on foundations. Later in this course, we will take a look at a few of those examples that had been

---

<sup>1</sup>A function *is* its “graph”, a subset of the Cartesian product of the domain and codomain of the function. For more details, we refer to more serious treatments of set theory, e.g. [Hal74].

<sup>2</sup>One noteworthy construction of the real numbers is as Dedekind cuts of the set of rational numbers, see for example Wikipedia [https://en.wikipedia.org/wiki/Dedekind\\_cut](https://en.wikipedia.org/wiki/Dedekind_cut).

<sup>3</sup>Indeed, in a foundational crisis, mathematicians felt the need to *construct* natural numbers 1, 2, 3, ..., instead of assuming their existence! Again we refer to serious treatments of set theory about the details.

considered controversial before the foundations were clarified: for example the Cantor set (Appendix B.3) and the Weierstrass function (Appendix ??), and countable and uncountable infinities (Appendix A.1).

From a general perspective, being precise about the foundations probably became necessary when the level of abstraction increased sufficiently so that one could no longer rely on too informal or intuitive reasoning. The historical rationale therefore has its parallel in mathematical studies: there comes a point at which the level of abstraction requires the student to make peace with foundational questions. Given that mathematics is founded on set theory, a student of mathematics must become familiar and comfortable with set theoretical notation and reasoning. To be fair, in this course we use set theory in a rather boring way as a language which is necessary for precise definitions and statements. Set theory and foundations of mathematics themselves would lead to fascinating questions, but we will not pursue such directions here.

As a pragmatic motivation for the mathematics student, set theory forms the common language in which virtually all standard topics in university level mathematics are phrased: measure and integration, probability theory, group theory and algebra, differential geometry and algebraic geometry, real analysis, complex analysis, functional analysis, etc. etc.

The present course, specifically, falls almost entirely within what is often labeled as *point set topology* (as opposed to *algebraic topology*). This terminology is best understood by noting that the expression *an element of a set* is used exactly parallel to the expression *a point in space*.<sup>4</sup> To call a “space” a “point set” is to emphasize that as a mathematical object a space is first and foremost the set consisting of its points, and only as such potentially equipped with some further structure.

## I.1. Basics of set theory

### Sets and elements

Sets consist of elements — a **set** is the collection of all **elements** that belong to it. It is customary to denote sets by upper case letters and their elements by lower case letters, as we mostly do also below. However, this is merely a typographical practice often adopted because it can serve as a notational cue to the reader; it will be discarded when appropriate.<sup>5</sup>

We denote  $a \in A$ , if  $a$  is an element of a set  $A$ . If  $a$  is not an element of  $A$ , we denote  $a \notin A$ . The *binary relation* denoted by the symbol  $\in$  (whose negation is denoted by the symbol  $\notin$ ) is at the heart of set theory. In natural language, “ $\in$ ” can

---

<sup>4</sup>We will in fact not precisely define the notions of *a point* and *a space* as such, but rather we use them as synonymous to *an element* and *a set* whenever we wish to draw attention to the fact that the set is also furnished with some additional structure. Various specific structures will be defined precisely later on: we will consider vector spaces, inner product spaces, normed spaces, metric spaces, topological spaces, ... — but the general term *space* itself will not be used in a definite mathematical sense.

<sup>5</sup>Indeed, for example for sets whose elements themselves are sets, it becomes impossible to strictly adhere to such a typographical “rule”.

be read as “to belong to”<sup>6</sup>, for example

$$\begin{aligned} a \in A & : && \text{(the element) } a \text{ belongs to (the set) } A, \\ a \notin A & : && \text{(the element) } a \text{ does not belong to (the set) } A. \end{aligned} \tag{I.1}$$

An often used informal (not exactly well-defined) example of a set is the set of all fruits, which has as elements apples, kiwis, mangos, etc. More mathematical examples of sets include

- the set  $\mathbb{Z}$  of all integers, having  $\dots, -2, -1, 0, 1, 2, \dots$  as its elements;
- the open interval  $(-\pi, \pi)$  is the set whose elements are the real numbers  $x$  satisfying  $-\pi < x < \pi$ ;
- the set of all continuous real-valued functions on the real line;
- the set of all subsets of the plane  $\mathbb{R}^2$ ;
- ....

In particular, sets may have numbers as their elements (as is probably very familiar already). But sets may also have much more general types of elements: functions, other sets, etc. etc.<sup>7</sup>

**Remark I.1** (What does it take to specify a set?).

A set is known when its elements are known — i.e., when for every possible object we are able to decide whether the object is an element of the set or not.

In particular two sets  $A$  and  $B$  are equal, denoted  $A = B$ , if they have exactly the same elements.

A few more examples of sets are:

- the **empty set**  $\emptyset$  is the set which does not contain any elements;
- a **singleton** is a set which contains a single element, the singleton consisting of an element  $a$  is denoted  $\{a\}$ .

**Remark I.2** (Do not confuse a singleton with its element!).

Note that

$$a \in \{a\}, \quad \text{but} \quad a \neq \{a\}.$$

**Exercise (✓) I.1** (Empty set vs. the set consisting of the empty set, etc.).

How many elements are in the following sets:

$$(a): \emptyset, \quad (b): \{\emptyset\}, \quad (c): \{\{\emptyset\}\} \quad ?$$

The simplest method of specifying a set is to list its elements:

- the notation  $\{a_1, a_2, \dots, a_n\}$  stands for the set consisting of  $a_1, a_2, \dots, a_n$ .

<sup>6</sup>Suomeksi “ $\in$ ” luetaan “kuuluu (joukkoon)”.

<sup>7</sup>As an example, problems in nonparametric statistics of real-valued data are basically optimization problems over the set of all those functions from the set of all measurable subsets of the set of real numbers to the unit interval, with satisfy the three axioms of a probability measure. In traditional (parametric) statistics the optimization problem is restricted to a subset of the above set. Don’t be fooled by apparent simplicity; the expressive power of set theory is vast, and actual mathematical applications make good use of that expressive power.

Note that we have, e.g.,  $\{9, 1, 1\} = \{1, 9\}$  (this set has two elements: 1 and 9). A set is only the collection of elements belonging to it; the order of elements has no significance, and repetition is redundant.

Another common method of specifying a set is to use a logical condition as a criterion for whether an element belongs to the set:

- if  $A$  is a set and  $P(x)$  is a logical proposition depending on a variable  $x$ , then the notation

$$\{x \in A \mid P(x)\} \quad (\text{I.2})$$

stands for the set consisting of those elements  $x$  of the set  $A$  for which  $P(x)$  is true.

**Example I.3** (Examples of sets defined by a condition).

In the following examples, various conditions are used to “extract” from the set  $\mathbb{R}$  of all real numbers a subset of those that satisfy the condition in question:

$$\begin{aligned} \{x \in \mathbb{R} \mid \sin(x) = 0\} &= \{\dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots\} \\ \{x \in \mathbb{R} \mid x^4 = \frac{1}{3}\} &= \left\{ \frac{-1}{\sqrt{\sqrt{3}}}, \frac{1}{\sqrt{\sqrt{3}}} \right\} \\ \{x \in \mathbb{R} \mid x^2 < 0\} &= \emptyset. \end{aligned}$$

We will also take some liberties to modify the above notational conventions in what we hope are reasonably self-explanatory ways, such as:

$$\begin{aligned} \{a_j \mid j \in \{1, 2, \dots, n\}\} &= \{a_1, a_2, \dots, a_n\}, \\ \{\sqrt{x} \mid x \in \mathbb{N}\} &= \{1, \sqrt{2}, \sqrt{3}, 2, \sqrt{5}, \dots\}, \quad \text{etc.} \end{aligned}$$

## Subsets

A set  $A$  is said to be a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ ; we then denote  $A \subset B$ . Being a subset is a binary relation among sets, denoted by the symbol  $\subset$ . Its negation is denoted by  $\not\subset$ , so  $A \not\subset B$  means that there exists at least one element in the set  $A$  which is not an element of  $B$ .

Note the difference between the relations  $\in$  and  $\subset$ .<sup>8</sup>

In natural language, the subset symbol  $\subset$  could be read as<sup>9</sup>

$$A \subset B \quad : \quad (\text{the set}) A \text{ is contained in } (\text{the set}) B, \quad (\text{I.3})$$

or simply as:  $A$  is a subset of  $B$ . For precise and unambiguous mathematical meaning, it is best to avoid mixing the natural language expressions “belongs to” and “is contained in”, which stand for the relations  $\in$  and  $\subset$ , respectively.

<sup>8</sup>If you have some experience in programming with *Python* or *Mathematica* (or other programming languages which encourage extensive use of nested lists), you are probably familiar with bugs which occur if you mix up a test of whether one list (perhaps containing just one element) is an element rather than a sublist of another list. A mathematical equivalent of such a programming error is mixing up the relations “ $\in$ ” and “ $\subset$ ”. Avoid that bug!

<sup>9</sup>Suomeksi “ $\subset$ ” voidaan lukea “sisältyy” — erotuksena relaatiosta “ $\in$ ”, joka luettiin “kuuluu”. Toisinaan selkeyden vuoksi on kuitenkin turvallisinta käyttää ilmaisua “on osajoukko”.

Evidently, the subset relation is **transitive**: if  $A \subset B$  and  $B \subset C$ , then  $A \subset C$ .

### Inverted relation symbols

When it is more appropriate to mention, e.g., a set before its element or a set before its subset, the inverted relation symbols  $\ni$  and  $\supset$  are used so that

$$\begin{array}{lll} A \ni a & \text{means} & a \in A, \\ B \supset A & \text{means} & A \subset B. \end{array}$$

The relation  $\supset$  can be read as “to contain”, so that

$$B \supset A \quad : \quad (\text{the set}) B \text{ contains } (\text{the set}) A,$$

**Remark I.4** (Equality of two sets).

Two sets  $A$  and  $B$  are equal if they contain exactly the same elements, which occurs if and only if both are subsets of the other:  $A \subset B$  and  $B \subset A$ .

While this may initially seem like a useless remark, it lends itself to a strategy of proof that is very commonly used. To *prove* that  $A = B$ , it is often practical to separately show  $A \subset B$  and  $A \supset B$ , by first arguing that any element of  $A$  must necessarily be also an element of  $B$ , and then vice versa.

### Familiar sets of numbers

The following examples of sets of numbers should be familiar:<sup>10</sup>

$$\text{the set of natural numbers} \quad \mathbb{N} = \{1, 2, 3, 4, \dots\} \quad (\text{I.4})$$

$$\text{the set of integers} \quad \mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{I.5})$$

$$\text{the set of rational numbers} \quad \mathbb{Q} = \left\{ \frac{n}{m} \mid n \in \mathbb{Z}, m \in \mathbb{N} \right\} \quad (\text{I.6})$$

$$\text{the set of real numbers}^{11} \quad \mathbb{R}. \quad (\text{I.7})$$

To give just a few examples, we have

$$\begin{array}{lll} 2021 \in \mathbb{N}, & -42 \in \mathbb{Z}, & \sqrt{2} \in \mathbb{R}, \\ 0 \notin \mathbb{N}, & -\frac{5}{7} \in \mathbb{Q}, & \sqrt{2} \notin \mathbb{Q}. \end{array}$$

---

<sup>10</sup>Note that zero is not considered a natural number — in this course. . . Unfortunately, mathematical literature is very divided regarding conventions about this issue; different areas of mathematics and different authors adopt different conventions. As a rule of thumb, in analysis it is typical to follow our present convention  $0 \notin \mathbb{N}$ , whereas in algebra it is typical to follow the convention that zero is a natural number. In fact, if you ever attend my algebra courses, then the symbol  $\mathbb{N}$  will be used in the different meaning. . . Sorry! In the present course, if we want to include zero, we use the notation  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$  instead of  $\mathbb{N}$ .

The last example is classical: it was known to the ancient Greeks.<sup>12</sup> We provide it here as one of our first examples of a careful mathematical proof. It is also a great illustration of the method of proof by contradiction.

**Proposition I.5** (Square root of two is irrational).

*We have  $\sqrt{2} \notin \mathbb{Q}$ .*

*Proof.* Suppose, by contrary, that  $\sqrt{2} \in \mathbb{Q}$ . In that case we can write  $\sqrt{2} = \frac{n}{m}$  with  $n, m \in \mathbb{N}$ . If the integers  $n$  and  $m$  have common factors, they may be cancelled, so we may assume that  $m$  and  $n$  here are chosen coprime (i.e., no integer greater than one divides both).

Multiplying by  $m$ , we obtain that  $\sqrt{2}m = n$ . Then squaring both sides we see that  $2m^2 = n^2$ . This shows that  $n^2$  is even, which is only possible if  $n$  is even, i.e.,  $n = 2k$  for some  $k \in \mathbb{N}$ . But in this case we find  $m^2 = \frac{1}{2}n^2 = \frac{1}{2}(2k)^2 = 2k^2$ , which similarly implies that  $m^2$  is even, and thus also  $m$  is even. Therefore 2 divides both  $n$  and  $m$ , which contradicts the choice of these coprime integers.

We conclude that  $\sqrt{2}$  could not have been rational in the first place, so  $\sqrt{2} \notin \mathbb{Q}$ .  $\square$

**Exercise (# · · ·) I.2** (Irrationality of  $\pi$ ).

Prove that  $\pi \notin \mathbb{Q}$ .

From the above sets of numbers we obtain examples of subsets, for example  $\mathbb{N} \subset \mathbb{Z}$ ,  $\mathbb{Z} \subset \mathbb{Q}$ , and  $\mathbb{Q} \subset \mathbb{R}$ ; or more concisely

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}.$$

By transitivity, further subset relations such as  $\mathbb{N} \subset \mathbb{Q}$ ,  $\mathbb{Z} \subset \mathbb{R}$ , and  $\mathbb{N} \subset \mathbb{R}$  are apparent from this.

## Intervals

Intervals of various types are extremely frequently used subsets of the real line  $\mathbb{R}$ . For intervals between points  $a, b \in \mathbb{R}$ ,  $a \leq b$ , we use the following notation and terminology:

$$\text{closed interval} \quad [a, b] = \{x \in \mathbb{R} \mid a \leq x \leq b\} \quad (\text{I.8})$$

$$(\text{bounded}) \text{ open interval} \quad (a, b) = \{x \in \mathbb{R} \mid a < x < b\} \quad (\text{I.9})$$

$$(a, b] = \{x \in \mathbb{R} \mid a < x \leq b\} \quad (\text{I.10})$$

$$[a, b) = \{x \in \mathbb{R} \mid a \leq x < b\}. \quad (\text{I.11})$$

We also use the following notation for unbounded intervals,

$$(-\infty, b] = \{x \in \mathbb{R} \mid x \leq b\}, \quad [a, +\infty) = \{x \in \mathbb{R} \mid a \leq x\}, \quad (\text{I.12})$$

$$(-\infty, b) = \{x \in \mathbb{R} \mid x < b\}, \quad (a, +\infty) = \{x \in \mathbb{R} \mid a < x\}, \quad (\text{I.13})$$

---

<sup>12</sup>The (cult of) Pythagoreans held it that only rational numbers were of divine origin, and a (probably dubious) legend has it that they killed Hippasus for revealing the irrationality of the length ( $\sqrt{2}$ ) of the diagonal of the unit square to the outside world. As for written records, a proof of this statement is contained in Euclid's "Elements" — almost surely the mathematical textbook that has endured the longest time in bestseller lists (few others are measured in millenia).

and sometimes  $(-\infty, +\infty) = \mathbb{R}$  is used the whole real axis. In addition to the bounded open intervals (I.9), also the unbounded open intervals (I.13) are considered **open intervals**. By contrast, there are no unbounded closed intervals.

## Operations with sets

New sets can be formed from old ones by set theoretic operations. For example, for two sets  $A$  and  $B$ , we define

$$\text{the \textbf{union} of } A \text{ and } B: \quad A \cup B = \{x \mid x \in A \text{ or } x \in B\} \quad (\text{I.14})$$

$$\text{the \textbf{intersection} of } A \text{ and } B: \quad A \cap B = \{x \mid x \in A \text{ and } x \in B\} \quad (\text{I.15})$$

$$\text{the \textbf{set difference} of } A \text{ and } B: \quad A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}. \quad (\text{I.16})$$

If  $B \subset A$ , then  $A \setminus B$  is also called the **complement** of  $B$  in  $A$ .

**Example I.6** (A few examples of set operations).

If we let

$$A = \{1, 2, 3, 4, 5, 6, \dots\} \quad (\text{natural numbers})$$

$$B = \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\} \quad (\text{even integers}),$$

then the intersection, union, and set differences are

$$\begin{aligned} A \cap B &= \{2, 4, 6, \dots\}, & A \setminus B &= \{1, 3, 5, \dots\}, \\ A \cup B &= \{\dots, -6, -4, -2, 0, 1, 2, 3, 4, \dots\}, & B \setminus A &= \{\dots, -6, -4, -2, 0\}. \end{aligned}$$

The set operations follow more or less obvious rules of calculation: for any sets  $A, B, C$  we have

$$\begin{cases} A \cup B = B \cup A \\ A \cap B = B \cap A \end{cases} \quad (\text{commutativity}) \quad (\text{I.17})$$

$$\begin{cases} A \cup (B \cup C) = (A \cup B) \cup C \\ A \cap (B \cap C) = (A \cap B) \cap C \end{cases} \quad (\text{associativity}) \quad (\text{I.18})$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C) \quad (\text{distributivity}) \quad (\text{I.19})$$

$$\begin{cases} A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C) \\ A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C). \end{cases} \quad (\text{De Morgan's laws}) \quad (\text{I.20})$$

The properties (I.17), at least, should be obvious from the definitions (I.14) and (I.15). We leave it as an exercise to verify most of the remaining properties, but as an example we prove the first De Morgan's law below.

*Proof of the first De Morgan's law.* The most straightforward proof strategy is the following: look at what condition characterizes an element of the set on the left hand side according to the definitions, and then successively rewrite equivalent versions of this condition until we reach the defining condition for elements on the set on the right hand side. The same proof strategy is convenient for all of the rules above. In the case of the first De Morgan's law,

the appropriate chain of equivalent conditions is

$$\begin{aligned}
 x \in A \setminus (B \cup C) &\iff x \in A \quad \text{and} \quad x \notin B \cup C \\
 &\iff x \in A \quad \text{and} \quad (x \notin B \text{ and } x \notin C) \\
 &\iff (x \in A \text{ and } x \notin B) \quad \text{and} \quad (x \in A \text{ and } x \notin C) \\
 &\iff x \in (A \setminus B) \cap (A \setminus C).
 \end{aligned}$$

Thus an element  $x$  belongs to the set  $A \setminus (B \cup C)$  if and only if it belongs to the set  $(A \setminus B) \cap (A \setminus C)$ . This is the asserted equality (I.20) of these two sets.  $\square$

**Exercise I.3** (Proofs of set theoretic identities).

Prove the remaining formulas among (I.17) – (I.20).

We can also form unions and intersections of more than two sets — in fact of arbitrary collections of sets. By an indexed collection  $(A_j)_{j \in J}$  of sets, we mean that a set  $A_j$  is given for each index  $j \in J$  (the set  $J$  is called the index set of the collection). The **union** of the collection is defined as

$$\bigcup_{j \in J} A_j = \left\{ x \mid x \in A_j \text{ for some } j \in J \right\}. \quad (\text{I.21})$$

The **intersection** is defined if the collection is non-empty, i.e. if  $J \neq \emptyset$ , and then it is

$$\bigcap_{j \in J} A_j = \left\{ x \mid x \in A_j \text{ for all } j \in J \right\}. \quad (\text{I.22})$$

Unions and intersections of two sets, (I.14) – (I.15), are recovered from (I.21) – (I.22) in the special case that the index set has two elements, for example  $J = \{1, 2\}$ .

When the index set is finite, in particular in the case  $J = \{1, 2, \dots, n\}$ , as alternative notations we often use

$$\begin{aligned}
 \bigcup_{j \in \{1, \dots, n\}} A_j &= \bigcup_{j=1}^n A_j = A_1 \cup \dots \cup A_n \\
 \bigcap_{j \in \{1, \dots, n\}} A_j &= \bigcap_{j=1}^n A_j = A_1 \cap \dots \cap A_n.
 \end{aligned}$$

When the index set is  $J = \mathbb{N}$ , it is conventional to use the alternative notations

$$\bigcup_{j \in \mathbb{N}} A_j = \bigcup_{j=1}^{\infty} A_j \quad \text{and} \quad \bigcap_{j \in \mathbb{N}} A_j = \bigcap_{j=1}^{\infty} A_j.$$

**Example I.7** (Union and intersection of nested intervals).

Consider the example  $A_n = \left[ \frac{-1}{n}, \frac{1}{n} \right]$  for  $n \in \mathbb{N}$ . Then the union and intersection of the collection  $(A_n)_{n \in \mathbb{N}}$  are

$$\begin{aligned}
 \bigcup_{n=1}^{\infty} \left[ \frac{-1}{n}, \frac{1}{n} \right] &= [-1, 1] && \text{(closed interval)} \\
 \bigcap_{n=1}^{\infty} \left[ \frac{-1}{n}, \frac{1}{n} \right] &= \{0\} && \text{(singleton).}
 \end{aligned}$$

The first of the above claims is easy, but let us justify the second claim in detail. Suppose that  $x \in \bigcap_{n=1}^{\infty} \left[ \frac{-1}{n}, \frac{1}{n} \right]$ . According to the definition of an intersection, this means that  $x \in \left[ \frac{-1}{n}, \frac{1}{n} \right]$  for all  $n \in \mathbb{N}$ . At the very least  $x$  must then be a real number. But note



that if  $x \neq 0$ , then by choosing a natural number  $n_0 > \frac{1}{|x|}$ , we have  $|x| > \frac{1}{n_0}$ , and so  $x \notin [\frac{-1}{n_0}, \frac{1}{n_0}]$ . This shows that a non-zero real number  $x \neq 0$  can not belong to all of the sets  $[\frac{-1}{n}, \frac{1}{n}]$ ,  $n \in \mathbb{N}$ , and thus does not belong to the intersection. On the other hand,  $x = 0$  belongs to each of the sets:  $0 \in [\frac{-1}{n}, \frac{1}{n}]$  for all  $n \in \mathbb{N}$ . By definition of the intersection, then,  $0 \in \bigcap_{n=1}^{\infty} [\frac{-1}{n}, \frac{1}{n}]$ . We have thus showed that the intersection consists of the single element 0, i.e.,  $\bigcap_{n=1}^{\infty} [\frac{-1}{n}, \frac{1}{n}] = \{0\}$  as claimed.

**Example I.8** (Another union and intersection of nested intervals).

Consider now the example  $B_n = (0, \frac{1}{n})$  for  $n \in \mathbb{N}$ . Then the union and intersection of the collection  $(B_n)_{n \in \mathbb{N}}$  are

$$\begin{aligned} \bigcup_{n=1}^{\infty} \left(0, \frac{1}{n}\right) &= (0, 1) && \text{(open interval)} \\ \bigcap_{n=1}^{\infty} \left(0, \frac{1}{n}\right) &= \emptyset && \text{(empty set).} \end{aligned}$$

The precise justifications of these claims are left as an exercise.

**Exercise I.4** (Details of Example I.8).

Provide careful reasoning to justify the claims made in Example I.8.

**Example I.9** (Yet another union and intersection of nested intervals).

Consider now the example  $C_n = [0, 1 - \frac{1}{2n}]$  for  $n \in \mathbb{N}$ . Then the union and intersection of the collection  $(C_n)_{n \in \mathbb{N}}$  are

$$\begin{aligned} \bigcup_{n=1}^{\infty} \left[0, 1 - \frac{1}{2n}\right] &= [0, 1) && \text{(half-open interval)} \\ \bigcap_{n=1}^{\infty} \left[0, 1 - \frac{1}{2n}\right] &= [0, \frac{1}{2}] && \text{(closed interval).} \end{aligned}$$

Note in particular that 1 does not belong to the union. The precise justifications are again left as an exercise.

**Exercise I.5** (Details of Example I.9).

Provide careful reasoning to justify the claims made in Example I.9.

**Exercise I.6** (De Morgan laws for general unions and intersections).

Let  $X$  be a set,  $J \neq \emptyset$  a nonempty index set, and  $A_j \subset X$  subsets for each  $j \in J$ . Prove De Morgan's laws for arbitrary unions and intersections:

$$X \setminus \bigcup_{j \in J} A_j = \bigcap_{j \in J} (X \setminus A_j) \quad \text{and} \quad X \setminus \bigcap_{j \in J} A_j = \bigcup_{j \in J} (X \setminus A_j).$$

## Cartesian products

Another operation of set theory is forming Cartesian products of sets.

If  $A$  and  $B$  are sets, then their **Cartesian product**  $A \times B$  is the set whose elements are **ordered pairs**  $(a, b)$  whose first member belongs to the former set,  $a \in A$ , and second member to the latter set,  $b \in B$ . In symbols, the Cartesian product is

$$A \times B = \left\{ (a, b) \mid a \in A, b \in B \right\}. \quad (\text{I.23})$$

A familiar example is the plane, which (as a set) is the Cartesian product of the real line  $\mathbb{R}$  with itself:

$$\mathbb{R}^2 = \mathbb{R} \times \mathbb{R} = \left\{ (x, y) \mid x \in \mathbb{R}, y \in \mathbb{R} \right\}.$$

Another familiar example is the rectangle, which (as a set) is the Cartesian product of two closed intervals

$$[\alpha, \beta] \times [\gamma, \delta] = \left\{ (x, y) \mid \alpha \leq x \leq \beta, \gamma \leq y \leq \delta \right\}.$$

Cartesian products of three sets are defined similarly as ordered triples  $A \times B \times C = \{(a, b, c) \mid a \in A, b \in B, c \in C\}$ , etc. The familiar  $n$ -dimensional space is an  $n$ -fold Cartesian product of the real line with itself

$$\mathbb{R}^n = \underbrace{\mathbb{R} \times \cdots \times \mathbb{R}}_{n \text{ times}} = \left\{ (x_1, \dots, x_n) \mid x_1, \dots, x_n \in \mathbb{R} \right\}.$$

The elements of  $\mathbb{R}^n$  are ordered  $n$ -tuples<sup>13</sup> of real numbers, consisting of the  $n$  coordinates of a point in this space.

One can consider Cartesian products of arbitrary collections of sets, but in this course the finite Cartesian products, such as the ones above, will be quite sufficient.

## I.2. Functions

What is a function, precisely? According to the familiar idea, a function  $f$  associates a (unique) value  $f(x)$  to each argument  $x$ . For a precise meaning, we must specify what is the set of acceptable arguments, and what is the set of possible values. The mathematical terms for these two sets are the **domain** of the function and the **codomain**<sup>14</sup> of the function, respectively.

The notations

$$f: X \rightarrow Y \quad \text{or} \quad X \xrightarrow{f} Y$$

indicate that  $f$  is a **function**, whose **domain** is a set  $X$ , and whose **codomain** is a set  $Y$ , i.e., that  $f$  assigns a unique value  $f(x) \in Y$  to each  $x \in X$ . The notation

$$x \mapsto f(x) \quad (\text{read: “}x \text{ maps to } f(x)\text{”})$$

may be used to emphasize this assignment (but the domain  $X$  and the codomain  $Y$  must also be specified or should be otherwise clear from the context). We may view this as a mapping from the points  $x$  of the domain  $X$  to points of the codomain  $Y$ , and the term **mapping** is considered synonymous to the term **function**. Functions

<sup>13</sup>One uses the terms **pair**, **triple**, **quadruple**, **quintuple**, **sextuple**, ... for ordered collections of two, three, four, five, six, ... elements. For general  $n$  it has become conventional to refer to an ordered collection of  $n$  elements as an  **$n$ -tuple**.

<sup>14</sup>Sometimes the term **range** is used instead of *codomain*. Beware, however, as it is also common to use the term *range* to mean the set of actually attained values of the function, which may be smaller than the set of possible values that we are a priori willing to consider. The Finnish terms ”määrittelyjoukko” and ”arvojoukko” are often used for the domain and codomain, and the latter unfortunately suffers from exactly the same ambiguity of usage as the term *range*.

The terms *domain* and *codomain* are in my opinion not very illuminating, but they are well-established mathematical terminology. Were they not, I would advocate for using *source* and *target* instead. The unambiguous and descriptive Finnish terms ”lähtö”/”lähtöjoukko” and ”maali”/”maali joukko” are used by [Väi99].

are often defined by giving a formula (but also any other way that ensures that a unique value is assigned to each argument is acceptable).

**Example I.10** (An example of a function).

We may define a function

$$r: \mathbb{R}^3 \rightarrow [0, +\infty)$$

by the formula

$$r((x, y, z)) = \sqrt{x^2 + y^2 + z^2} \quad \text{for } (x, y, z) \in \mathbb{R}^3.$$

The domain of this function  $r$  is (by definition)  $\mathbb{R}^3$ , and the codomain is (by definition)  $[0, +\infty)$ . We must only make sure that for each  $(x, y, z) \in \mathbb{R}^3$ , the formula above is meaningful (as it is), and gives a value in the set  $[0, +\infty)$  (as it does).

Often giving a name or a symbol for a function is purposeful: the letters  $f, g, \dots$  are most commonly used,  $r$  was chosen in Example I.10, and you have without a doubt seen also

$$\log: (0, \infty) \rightarrow \mathbb{R}, \quad \sin: [0, 2\pi) \rightarrow [-1, 1], \quad \text{etc.}$$

Occasionally, it is more meaningful to talk of functions  $X \rightarrow Y$  (i.e., functions with domain  $X$  and codomain  $Y$ ) without explicitly naming them. The “maps to” notation “ $\mapsto$ ” may then be particularly convenient.

**Example I.11** (A few more examples of functions).

- The exponential function  $\mathbb{R} \rightarrow (0, +\infty)$  is given by the formula  $x \mapsto e^x$ .
- The formula  $\theta \mapsto (\cos(\theta), \sin(\theta))$  defines a function  $(-\pi, +\pi] \rightarrow \mathbb{R}^2$ .

Sometimes we simply consider established symbols as the names of functions, and indicate by a “dot” at which slot the argument is to be inserted to get the value.

**Example I.12** (Yet a few more examples of functions).

- The square root function  $\sqrt{\cdot}: [0, +\infty) \rightarrow [0, +\infty)$  is given by  $x \mapsto \sqrt{x}$ .
- The absolute value function  $|\cdot|: \mathbb{R} \rightarrow [0, +\infty)$  is given by  $x \mapsto |x|$ .

**Example I.13** (The identity function).

Let  $X$  be a set. Then the function

$$\text{id}_X: X \rightarrow X \quad \text{id}_X(x) = x \quad \text{for } x \in X \quad (\text{I.24})$$

is known as the **identity function** on  $X$ .

Note that functions defined by the same formula are not the same functions unless also their domains and codomains are the same!

**Example I.14** (Different functions given by the same formula).

The following functions are given by the rule  $x \mapsto x^2$ :

- $f_1: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f_1(x) = x^2$ ,
- $f_2: \mathbb{R} \rightarrow [0, +\infty)$  given by  $f_2(x) = x^2$ ,
- $f_3: [0, +\infty) \rightarrow \mathbb{R}$  given by  $f_3(x) = x^2$ ,
- $f_4: (-\infty, 0] \rightarrow \mathbb{R}$  given by  $f_4(x) = x^2$ ,
- $f_5: (-\infty, 0] \rightarrow [0, +\infty)$  given by  $f_5(x) = x^2$ .

Nevertheless, each of the above is a different function:  $f_1 \neq f_2$ ,  $f_3 \neq f_5$ , etc.

The examples we have used so far should appear unintimidating. The concept of a function is nevertheless very general: the domain and range of a function may be arbitrary sets. We will in particular encounter functions whose arguments are functions (i.e., the domain is some set of functions), functions whose values are sets (i.e., the codomain is some set whose elements are sets), etc. Be forewarned.

## Surjective, injective, and bijective functions

A few properties of functions are relevant.

**Definition I.15** (Surjectivity, injectivity, and bijectivity).

Let  $f: X \rightarrow Y$  be a function.

We say that  $f$  is **surjective** if for every  $y \in Y$  there exists an  $x \in X$  such that  $y = f(x)$ .

We say that  $f$  is **injective** if for any  $x_1, x_2 \in X$  which are different,  $x_1 \neq x_2$ , also the corresponding function values are different,  $f(x_1) \neq f(x_2)$ .

We say that  $f$  is **bijective** if it is both surjective and injective.

Consider a function  $f: X \rightarrow Y$  and a given  $y \in Y$ . If  $f$  is injective, there can be at most one  $x \in X$  such that  $f(x) = y$  (since for any  $x' \neq x$  we have  $f(x') \neq f(x) = y$ ). For a surjective function there always exists at least one such  $x$ . For bijective functions, therefore, there exists exactly one such  $x$ , i.e., the condition  $y = f(x)$  gives rise to a well-defined mapping  $y \mapsto x$ , the **inverse function**

$$f^{-1}: Y \rightarrow X.$$

**Exercise I.7** (Examples of injectivity and surjectivity).

Consider the functions

$$\begin{aligned} f_1: \mathbb{R} \rightarrow \mathbb{R}, & \quad f_2: \mathbb{R} \rightarrow [0, +\infty), & \quad f_3: [0, +\infty) \rightarrow \mathbb{R}, \\ f_4: (-\infty, 0] \rightarrow \mathbb{R}, & \quad f_5: (-\infty, 0] \rightarrow [0, +\infty), \end{aligned}$$

from Example I.14, each given by the formula  $x \mapsto x^2$ . Check that:

- (a) The functions  $f_2$  and  $f_5$  are surjective, whereas  $f_1$ ,  $f_3$ , and  $f_4$  are not.
- (b) The functions  $f_3$ ,  $f_4$ , and  $f_5$  are injective, whereas  $f_1$  and  $f_2$  are not.
- (c) The function  $f_5$  is bijective, whereas  $f_1$ ,  $f_2$ ,  $f_3$ , and  $f_4$  are not.
- (d) The inverse function  $f_5^{-1}: [0, +\infty) \rightarrow (-\infty, 0]$  of  $f_5$  is given by  $f_5^{-1}(y) = -\sqrt{y}$ , whereas  $f_1$ ,  $f_2$ ,  $f_3$ , and  $f_4$  do not have inverse functions.

**Example I.16** (The identity function is bijective).

For any set  $X$ , the identity function  $\text{id}_X: X \rightarrow X$  of Example I.13 is bijective. It is its own inverse function<sup>15</sup>:  $\text{id}_X^{-1} = \text{id}_X$ .

---

<sup>15</sup>The property of the identity function that it is its own inverse may seem special, but there are also other functions satisfying this. Such functions are called *involutions*. We leave it to the reader to construct examples: the formulas  $x \mapsto -x$ ,  $y \mapsto \frac{1}{y}$ ,  $z \mapsto \frac{-1}{z}$ ,  $\beta \mapsto -\frac{1}{2} \log(\tanh(\beta))$  may serve as an inspiration, but remember to specify the domains and codomains!

## Compositions of functions

Suppose that

$$f: X \rightarrow Y \quad \text{and} \quad g: Y \rightarrow Z$$

are two functions such that the codomain of  $f$  is the same set as the domain of  $g$  (both are  $Y$ ). Then we may form a new function,

$$g \circ f: X \rightarrow Z$$

called the **composition** of  $f$  and  $g$  by the rule that at a point  $x \in X$  of its domain the value is

$$(g \circ f)(x) = g(f(x)) \in Z.$$

The composition is illustrated in the following diagram<sup>16</sup>:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ & \searrow & \text{\scriptsize $g \circ f$} & \nearrow & \\ x & \xrightarrow{f} & f(x) & \xrightarrow{g} & g(f(x)) \end{array}$$

**Example I.17** (A composition of functions).

Consider the functions

$$\begin{aligned} f: (0, 2\pi) &\rightarrow [-1, 1] & f(\theta) &= \cos(\theta) \quad \text{for } \theta \in (0, 2\pi) \\ g: [-1, 1] &\rightarrow \mathbb{R} & g(u) &= \frac{1}{(1-u)^2} \quad \text{for } u \in [-1, 1]. \end{aligned}$$

Their composition is

$$g \circ f: (0, 2\pi) \rightarrow \mathbb{R} \quad (g \circ f)(\theta) = \frac{1}{(1 - \cos(\theta))^2} \quad \text{for } \theta \in (0, 2\pi).$$

**Example I.18** (Compositions with an inverse function).

Suppose that

$$f: X \rightarrow Y$$

is bijective, and consider its inverse function

$$f^{-1}: Y \rightarrow X.$$

Then the composition of  $f$  and  $f^{-1}$  is (as you should verify from the definitions)

$$f^{-1} \circ f: X \rightarrow X \quad (f^{-1} \circ f)(x) = x \quad \text{for } x \in X.$$

The composition of  $f^{-1}$  and  $f$ , on the other hand, is (as you should also verify)

$$f \circ f^{-1}: Y \rightarrow Y \quad (f \circ f^{-1})(y) = y \quad \text{for } y \in Y.$$

So both compositions are identity functions (see Example exa: the identity function) but on different sets:

$$f^{-1} \circ f = \text{id}_X \quad f \circ f^{-1} = \text{id}_Y.$$

<sup>16</sup>This should also make it clear why the composition of  $f$  and  $g$  is denoted by  $g \circ f$  in this order. We take an element  $x \in X$ , first apply  $f$  to it to get  $f(x) \in Y$ , and then apply  $g$  to the result to get  $g(f(x)) \in Z$ . The “strange” order in the composition is dictated by our convention that the function is written to the left of its argument. If we would denote for example  $f(x) = x * f$  instead, then the composition in this order would be defined by the formula  $x \mapsto (x * f) * g$  in which  $f$  and  $g$  appear in the same order as they are applied (assuming you read from left to right! — Hebrew, Persian, or Arabic readers probably have less qualms about the order of  $g \circ f$  in the first place...).

**Exercise (✓) I.8** (Composition of injective functions is injective).

Show that if both  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  are injective, then  $g \circ f: X \rightarrow Z$  is also injective.

**Exercise (✓) I.9** (Composition of surjective functions is surjective).

Show that if both  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  are surjective, then  $g \circ f: X \rightarrow Z$  is also surjective.

**Exercise I.10** (Injectivity of composition).

Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be functions such that  $g \circ f: X \rightarrow Z$  is injective.

- (a) Show that  $f$  is also injective.
- (b) Construct an example which shows that  $g$  does not have to be injective.

**Exercise I.11** (Surjectivity of composition).

Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be functions such that  $g \circ f: X \rightarrow Z$  is surjective.

- (a) Show that  $g$  is also surjective.
- (b) Construct an example which shows that  $f$  does not have to be surjective.

The following exercise shows that the conditions in Example I.18 characterize the inverse function.

**Exercise I.12** (Characterization of the inverse function).

Suppose that  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  are two functions such that

$$g \circ f = \text{id}_X \qquad f \circ g = \text{id}_Y.$$

Prove that the both  $f$  and  $g$  are bijective, and we have  $g = f^{-1}$  and  $f = g^{-1}$ .

## Restricting the domain

Given a function  $f: X \rightarrow Y$  and a subset  $A \subset X$  of its domain, it is possible to consider a function defined only in the subset by otherwise the same formula. This function  $A \rightarrow Y$  is denoted by  $f|_A$  and is called the **restriction** of  $f$  to the subset  $A$ ; it is given by

$$\begin{aligned} f|_A: A &\rightarrow Y \\ f|_A(a) &= f(a) \quad \text{for } a \in A. \end{aligned} \tag{I.25}$$

**Example I.19** (Examples of restrictions of functions).

Consider again the functions

$$\begin{aligned} f_1: \mathbb{R} &\rightarrow \mathbb{R}, & f_2: \mathbb{R} &\rightarrow [0, +\infty), & f_3: [0, +\infty) &\rightarrow \mathbb{R}, \\ f_4: (-\infty, 0] &\rightarrow \mathbb{R}, & f_5: (-\infty, 0] &\rightarrow [0, +\infty), \end{aligned}$$

from Example I.14, each given by the formula  $x \mapsto x^2$ . Then we have

$$f_3 = f_1|_{[0, +\infty)}, \quad f_4 = f_1|_{(-\infty, 0]}, \quad f_5 = f_2|_{(-\infty, 0]}$$

## Images and preimages under functions

Let  $f: X \rightarrow Y$  be a function.

If  $A \subset X$  is a subset of the domain of  $f$ , then the **image** of  $A$  under  $f$  is the subset

$$f[A] = \{f(a) \mid a \in A\} \tag{I.26}$$

of the codomain of  $f$  consisting of those points  $y \in Y$  which are obtained as the value at some point  $a \in A \subset X$ , i.e.,  $y = f(a)$ .

**Example I.20** (Some images).

Consider the function

$$f: \mathbb{R} \rightarrow \mathbb{R} \quad \text{given by} \quad f(x) = x^2 \quad \text{for } x \in \mathbb{R}.$$

The image of the subset  $A = \{-7, -5, 5, 7\} \subset \mathbb{R}$  is

$$f[\{-7, -5, 5, 7\}] = \{25, 49\}.$$

In particular the image of a subset may contain fewer elements than the subset itself.

**Example I.21** (The image of a singleton).

The image of a singleton  $\{x\} \subset X$  is  $f[\{x\}] = \{f(x)\}$ .

**Exercise (✓) I.13** (Singleton image).

If the image of  $A \subset X$  is a singleton  $f[A] = \{y\}$ , then what can be said about the behavior of the function  $f$  on the subset  $A$ ?

**Example I.22** (Surjectivity by images).

A function  $f: X \rightarrow Y$  is surjective if and only if the image of the whole domain is the whole codomain,  $f[X] = Y$ .

If  $B \subset Y$  is a subset of the codomain of  $f$ , then the **preimage** of  $B$  under  $f$  is the subset

$$f^{-1}[B] = \{x \in X \mid f(x) \in B\} \quad (\text{I.27})$$

of the domain of  $f$  consisting of those points  $x \in X$  at which the value belongs to the subset  $B \subset Y$ . The notation  $f^{-1}$  is used for both the inverse function and the preimage, but we try to consistently use square brackets in the latter case.<sup>17</sup> The inverse function only exists if  $f$  is a bijection, whereas preimages are always defined.

**Example I.23** (Some preimages).

Consider the function

$$f: \mathbb{R} \rightarrow \mathbb{R} \quad \text{given by} \quad f(x) = x^2 \quad \text{for } x \in \mathbb{R}.$$

The preimage of the (singleton) subset  $A = \{5\} \subset \mathbb{R}$  is

$$f^{-1}[\{5\}] = \{-\sqrt{5}, \sqrt{5}\}.$$

In particular the preimage of a singleton can contain more than one element.

The preimage of the (singleton) subset  $A = \{-5\} \subset \mathbb{R}$  is the empty set,

$$f^{-1}[\{-5\}] = \emptyset.$$

In particular the preimage of a nonempty set can be empty.

---

<sup>17</sup>In most mathematical literature ordinary parentheses are used also for preimages, or parentheses are altogether omitted. You are in any case expected to figure out which one is which, since the inverse function is applied to *elements* of the codomain, whereas in the preimage is applied to *subsets* of the codomain.

Similarly, we try to consistently use square brackets for the images of subsets under a function. Elsewhere in literature ordinary parentheses or no parentheses at all are used. Again you are in any case expected to figure out which one is which, since the function is applied to *elements* of the domain, whereas in the image is applied to *subsets* of the domain.

**Example I.24** (Bijectivity by preimages).

A function  $f: X \rightarrow Y$  is bijective if and only if the preimage of every singleton  $\{y\} \subset Y$  is a singleton.

**Example I.25** (Injectivity by preimages).

A function  $f: X \rightarrow Y$  is injective if and only if the preimage of every singleton  $\{y\} \subset Y$  is either a singleton or the empty set.

**Exercise I.14** (Images and preimages of unions and intersections).

Let  $X$  and  $Y$  be sets and  $f: X \rightarrow Y$  a function.

- (a) Show that for any  $C, D \subset Y$ , the preimages satisfy

$$f^{-1}[C \cup D] = f^{-1}[C] \cup f^{-1}[D].$$

- (b) Show that for any  $C, D \subset Y$ , the preimages satisfy

$$f^{-1}[C \cap D] = f^{-1}[C] \cap f^{-1}[D].$$

- (c) Show that for any  $A, B \subset X$ , the images satisfy

$$f[A \cup B] = f[A] \cup f[B].$$

- (d) Give an example in which for the images of subsets  $A, B \subset X$  we have

$$f[A \cap B] \neq f[A] \cap f[B].$$

Hint: In parts (a)–(c) it is possible to argue by a chain of equivalent conditions

$$x \in \text{left hand side set} \iff \dots \iff x \in \text{right hand side set}.$$

A comparison of (b) and (d) above indicates that preimages behave better with set operations than images. Another example is the following.

**Exercise I.15** (Images and preimages of complements).

Let  $X$  and  $Y$  be sets and  $f: X \rightarrow Y$  a function.

- (a) Suppose that  $C \subset D \subset Y$ . Show that the preimages satisfy

$$f^{-1}[D \setminus C] = f^{-1}[D] \setminus f^{-1}[C].$$

- (b) Give an example in which for the images of subsets  $A \subset B \subset X$  we have

$$f[B \setminus A] \neq f[B] \setminus f[A].$$

The formulas of Exercise I.14(a)–(c) generalize to arbitrary unions and intersections (more than two sets), and it is easy to modify the proofs to account for this general case. Specifically, if  $f: X \rightarrow Y$  is a function and  $B_j \subset Y$  for  $j \in J$ , then the preimages satisfy

$$f^{-1}\left[\bigcup_{j \in J} B_j\right] = \bigcup_{j \in J} f^{-1}[B_j], \quad f^{-1}\left[\bigcap_{j \in J} B_j\right] = \bigcap_{j \in J} f^{-1}[B_j].$$

Likewise if  $A_j \subset X$  for  $j \in J$ , then the images satisfy

$$f\left[\bigcup_{j \in J} A_j\right] = \bigcup_{j \in J} f[A_j].$$



### I.3. Logic and related notation

Mathematics in general concerns with statements that can be logically proven to be true, starting from some specified assumptions. In other words, we want to deduce conclusions of interest from known premises. The rules for valid deduction are the subject of logic. Logic itself could be formalized, but for our purposes it suffices to introduce a little bit of notation and typical examples. The main message is that in this course and in all mathematics, we must have unambiguous definitions and statements, and we must provide valid arguments to justify the statements made.

#### Quantifiers

##### *Statements involving free variables*

Commonly, statements (predicates, logical formulas) involve one or more variables, and whether the statements are true or false depends on those variables. A statement may be thought of as a function of those variables, whose possible values are true and false.

**Example I.26** (An example statement depending on a variable).

The statement

“ $k$  is even”

is true for if the variable  $k$  has the value  $k = -8$ , but it is false if  $k = 33$ .

Those variables of which the truth value of a statement depends are called **free variables**; without specifying values for the free variables, the statement itself is not yet true nor false. A statement without free variables is called a **sentence** — it is then either true or false as it stands. Self-contained meaningful logical statements must be sentences!

In mathematical text we often indicate that we (temporarily) fix the values of (some) variables by expressions such as: “fix  $n \in \mathbb{N}$ ” (after which the variable  $n$  is no longer thought of as free, but having some a priori arbitrary but fixed natural number value) or “let  $\varepsilon > 0$ ” (after which the variable  $\varepsilon$  is no longer thought of as free, but fixed to some a priori arbitrary positive real value). Fixing a value for a variable in effect makes it a named constant.<sup>18</sup> Then statements involving the variables in question again have definite truth values.

##### *Statements involving quantifiers*

The following two example statements,

“for all real numbers  $x$  we have  $x^2 \geq 0$ ”

and

“there exists a real number  $x$  such that  $x^2 = -1$ ”,

---

<sup>18</sup>This is the way a computer treats variables when running a program: at any time they are used, they must have assigned values or otherwise errors arise. The computer scientist or a programmer, on the other hand, may think of variables much like a mathematician or a logician.

both involve a variable  $x$  which has not been fixed, but the statements are in fact sentences. In particular they have definite truth values: the former sentence is true and the latter is false. The variable  $x$  in the above formulas is not free, because a *quantifier* applies to it: each statement above makes a claim about the *quantity* of values of  $x$  for which another statement, which involves  $x$  as a free variable, is true.

There are two quantifiers in ordinary logic. The **universal quantifier** denoted by the symbol  $\forall$  means that the statement that follows it is true *for all values of the variable*; in natural language its symbol  $\forall$  is read as “for all”. The **existential quantifier** denoted by the symbol  $\exists$  means that the statement that follows it is true *for at least one value of the variable*; in natural language its symbol  $\exists$  is read as “there exists”. Both quantifiers must be followed first by the variable which they quantify, and they should (in careful usage) also be followed by a specification of the set in which the variable is allowed to take values, and then another statement (possibly) involving the variable. For clarity we usually also separate the other statement (in which the variable appears as free) by a colon (:). For example the two example statements that we started with should be written as

$$\forall x \in \mathbb{R} : \quad x^2 \geq 0$$

and

$$\exists x \in \mathbb{R} : \quad x^2 = -1.$$

**Example I.27** (Revisiting subsets).

The meaning of the subset relation  $A \subset B$  is, in concise logical notation with quantifiers,

$$\forall x \in A : \quad x \in B.$$

In other words, the statement above is true if  $A \subset B$ , and false if  $A \not\subset B$ .

**Example I.28** (Revisiting unions and intersections).

Let  $(A_j)_{j \in J}$  be a collection of sets. Then the meaning of the statement that an element  $x$  belongs to the union of this collection,  $x \in \bigcup_{j \in J} A_j$ , can be expressed in concise logical notation with quantifiers as

$$\exists j \in J : \quad x \in A_j.$$

Similarly, if the collection is nonempty ( $J \neq \emptyset$ ), then the meaning of the statement that an element  $x$  belongs to the intersection of this collection,  $x \in \bigcap_{j \in J} A_j$ , can be expressed as

$$\forall j \in J : \quad x \in A_j.$$

Note that the quantified variable appears as a free variable in the statement that follows, but that statement may itself have quantifiers for other variables. This is in fact very common in, e.g., definitions related to metric space topology (continuity, limits of sequences, etc.) — we will see plenty of examples later. For now, as just one example statement with more than one quantifier, consider

$$\forall \alpha \in \mathbb{R} : \quad \exists C > 0 : \quad \forall x \in \mathbb{R} : \quad |e^{3+5\alpha x-x^2}| \leq C$$

(this statement is true, by the way). Literally this statement reads “for all real numbers  $\alpha$  there exists a positive number  $C$  such that for all real numbers  $x$  we have  $|e^{3+5\alpha x-x^2}| \leq C$ ”, and allowing for a bit more liberties of expression and interpretation could also be read “whatever the value of the real parameter  $\alpha$ , the expression  $e^{3+5\alpha x-x^2}$  involving a real variable  $x$  is bounded in absolute value by some positive constant  $C$ ”. The use of logical notation makes the statements more concise

and unambiguous. By contrast, in natural language one has to be quite careful to achieve an unambiguous meaning.<sup>19</sup>

Note that the order of quantifiers matters! For instance the statement

$$\forall x \in \mathbb{R} : \exists C > 0 : \forall \alpha \in \mathbb{R} : |e^{3+5\alpha x-x^2}| \leq C$$

is false, although it “only” differs from the previous example (a true statement) by the order of quantifiers.

**Exercise I.16** (Check the truth values of the above examples).

Prove that the statement  $\forall \alpha \in \mathbb{R} : \exists C > 0 : \forall x \in \mathbb{R} : |e^{3+5\alpha x-x^2}| \leq C$  is true and that the statement  $\forall x \in \mathbb{R} : \exists C > 0 : \forall \alpha \in \mathbb{R} : |e^{3+5\alpha x-x^2}| \leq C$  is false.

## Implications and equivalences

### Implications

Suppose that  $P$  and  $Q$  are two statements (predicates) possibly involving free variables (the same variables in both). Then one can form a new statement out of them, called an **implication** and denoted by

$$P \implies Q,$$

which in natural language can be read in any of the following ways

$P$  implies  $Q$  ;

if  $P$ , then  $Q$  ;

$P$  only if  $Q$  .

The natural language descriptions (the most transparent of which is probably “if  $P$ , then  $Q$ ”) already should explain exactly how the truth value of an implication is determined:  $P \implies Q$  is true if either  $P$  and  $Q$  are both true, or if  $P$  is false.<sup>20</sup>

**Example I.29** (Revisiting subsets again).

In concise logical notation, the meaning of the subset relation  $A \subset B$  is the implication

$$x \in A \implies x \in B.$$

In other words, the above logical statement is true if  $A \subset B$ , and false if  $A \not\subset B$ .

A typical use of an implication is in theorem statements:  $P$  may represent the assumptions of a theorem (the premises) and  $Q$  the conclusion, so the theorem statement itself is the implication  $P \implies Q$  (“if the assumptions hold, then also the conclusion holds”). An implication in a theorem statement should of course be true (the role of the *proof* of the theorem is exactly to verify the validity of the asserted implication).

<sup>19</sup>This does not mean that natural language is strictly worse. Among advantages of natural language are that it allows to emphasize certain aspects of the statement and to draw attention to interpretations (for example interpreting  $\alpha$  as a parameter,  $C$  as a constant, and  $x$  as a variable in the above expression). None of this affects the validity of logical statements, but to claim that such things are irrelevant would be rather extreme...

<sup>20</sup>In particular, if  $P$  does not hold, the implication is claiming nothing whatsoever about the truth or falsity of  $Q$ . The only situation in which the implication is false is if  $P$  is true but  $Q$  is nevertheless false.

Note, however, that the implication  $P \Rightarrow Q$  is merely a statement about the relationship of the truth values of the two statements  $P$  and  $Q$  — it does not signify any *causal* relationship.

Instead of  $P \Rightarrow Q$  we also occasionally switch the places of  $P$  and  $Q$  and invert the arrow direction, i.e., we alternatively write

$$Q \Leftarrow P,$$

which in natural language can be read in any of the following ways

$$\begin{aligned} Q \text{ is implied by } P ; \\ Q \text{ if } P . \end{aligned}$$

So we have seen altogether five different phrases in natural language for the same implication — the logical meaning of each is exactly the same, but they may be used to express a slightly different emphasis.

### *Equivalences*

Two statements  $P$  and  $Q$ , possibly depending on some free variables (same in both), are said to be **logically equivalent** if they have the same truth value for all values of the free variables. This precisely amounts to requiring that both implications  $P \Rightarrow Q$  and  $P \Leftarrow Q$  hold ( $Q$  is true if  $P$  is true and vice versa). We denote **logical equivalence** by

$$P \Longleftrightarrow Q,$$

and in natural language read it as either

$$\begin{aligned} P \text{ and } Q \text{ are (logically) equivalent ;} \\ P \text{ if and only if } Q . \end{aligned}$$

**Example I.30** (Revisiting the equality of sets).

The meaning of the equality of two sets  $A = B$  is

$$x \in A \Longleftrightarrow x \in B.$$

**Remark I.31** (Warning: a convention used in definitions).

When a new term is defined, we usually say that the term is used if some conditions are satisfied. In this context we actually mean that the term is used *if and only if* the conditions are satisfied! But it is conventional in definitions to only state the “if” part, leaving it implicit that the newly introduced term is *not* used unless the conditions are satisfied.

For example our definition of a bijective function was: “we say that  $f$  is bijective if it is both injective and surjective”. We of course *also* meant that we *do not* call  $f$  bijective *unless* it is both injective and surjective.

So while the natural language word “if” in usual mathematical text signifies just one implication, in definitions it is conventionally used to signify equivalence.

### Negations

If  $P$  is a logical statement (predicate), its **negation**

$$\text{not } P$$

is the logical statement which always has the opposite truth value: if  $P$  is true, then **not**  $P$  is false and if  $P$  is false, then **not**  $P$  is true. In natural language, **not** is read as “not”.

Clearly the statement **not** (**not**  $P$ ) is logically equivalent to  $P$  (a double negation).

We have already introduced specific symbols for the negations of a few common statements, e.g.,

$$\begin{array}{lll} \text{not } (a = b) & \text{is denoted} & a \neq b \\ \text{not } (a \in A) & \text{is denoted} & a \notin A \\ \text{not } (A \subset B) & \text{is denoted} & A \not\subset B. \end{array}$$

### *Contrapositives*

The implications

$$\begin{array}{ll} P \implies Q \\ \text{not } Q \implies \text{not } P \end{array} \tag{I.28}$$

are logically equivalent — you should make sure that you understand why!<sup>21</sup> The second implication is known as the **contrapositive** of the first.

Let us take an example from Section I.2.

**Example I.32** (Revisiting injectivity).

Recall definition of an injective function  $f: X \rightarrow Y$  from Definition I.15. In logical symbols, injectivity means

$$\forall x_1, x_2 \in X : \quad x_1 \neq x_2 \implies f(x_1) \neq f(x_2). \tag{I.29}$$

An equivalent form of this condition can be obtained by taking a contrapositive of the implication above. This is particularly helpful, since the contrapositive will involve negations of the propositions  $x_1 \neq x_2$  and  $f(x_1) \neq f(x_2)$ , which are simply  $x_1 = x_2$  and  $f(x_1) = f(x_2)$ , respectively. Using the contrapositive form, we find that injectivity equivalently means

$$\forall x_1, x_2 \in X : \quad f(x_1) = f(x_2) \implies x_1 = x_2. \tag{I.30}$$

The original formulation (I.29) of injectivity can be read: “At any two different points, the function has different values”. The formulation (I.30) obtained by contrapositive can be read: “Whenever the values at two points are the same, the points themselves must be the same”. With a bith of thinking, one easily convinces oneself about the equivalence of these, but routine use of logical symbols clearly facilitates this.

The equivalence of the two implications in (I.28) in particular underlies the idea of a *proof by contrapositive* or an *indirect proof* — of which the *proof by contradiction* is essentially a special case.

To elaborate, suppose our goal is to prove that from assumption  $P$ , conclusion  $Q$  follows. Another way of doing that is to prove that if we assume that the desired conclusion  $Q$  does not hold, then it implies that the assumption  $P$  cannot hold, either. This is the proof by contrapositive.

---

<sup>21</sup>The most straightforward way to convince oneself about this is to write a truth table, i.e., to consider the truth value of both implications for all the four possibilities of the truth values of the propositions  $P$  and  $Q$  ( $P$  false and  $Q$  false,  $P$  false and  $Q$  true,  $P$  true and  $Q$  false,  $P$  true and  $Q$  true).

In particular, if there are no (explicit) assumptions, we can consider the assumptions represented by an identically true proposition  $P$ . Then this proof strategy is proof by contradiction. Indeed, by showing the contrapositive, we find that if the the desired conclusion  $Q$  would not hold, then the negation of the identically true assumption  $P$  would hold, which is absurd; a contradiction. The proof of Proposition I.5 is a classic example.

### *Negations and quantifiers*

Since typical definitions in metric space topology involve many logical quantifiers, it is essential to be comfortable working with them. In particular one has to be able to routinely form the negation of a statement involving a quantifier.

Consider a statement involving the universal quantifier, say

$$\forall x \in X : P(x).$$

Its negation, **not**  $(\forall x \in X : P(x))$ , is logically equivalent with

$$\exists x \in X : \text{ not } P(x);$$

you should make sure you understand why!

A common use of this observation is that to *disprove* the validity of a claim (for example a belief that we might have) that  $P(x)$  holds for all  $x \in X$ , it is sufficient to find just one example of an  $x \in X$  for which  $P(x)$  is false. Such an example is called a **counterexample**, because it is sufficient to invalidate the whole original claim (which started with the universal quantifier).

And even if we are seeking to prove rather than to disprove a claim of this form, in an indirect proof (contrapositive) we would first form the negation.

Symmetrically, consider a statement involving the existential quantifier, say

$$\exists x \in X : Q(x).$$

Its negation, **not**  $(\exists x \in X : Q(x))$ , is logically equivalent with (think about it)

$$\forall x \in X : \text{ not } Q(x).$$

On the surface this does not appear as practical as the negation of a statement with a universal quantifier. But in indirect proofs, or when disproving a claim (say a false belief) of this form, we would have to form the negation. Another typical use of this appears when a statement involves many quantifiers; we may want to consider the negation to get rid of some universal quantifiers, but the existential quantifiers are affected as well.

**Example I.33** (There are arbitrarily large natural numbers).

The logical statement

$$\forall q \in \mathbb{Q} : \exists n \in \mathbb{N} : n > q \tag{I.31}$$

has the interpretation that there are arbitrarily large natural numbers (“for any rational number  $q$  there exists a larger natural number  $n$ ”).

The statement (I.31) is true: it is the so called **Archimedean property** of the rational numbers. Let us examine a proof of this statement using a proof by contradiction.

For a proof by contradiction, one would suppose that (I.31) is false, and show that this leads to a contradiction. If (I.31) is false, then its negation

$$\text{not} \left( \forall q \in \mathbb{Q} : \exists n \in \mathbb{N} : n > q \right).$$

is true. The original statement (I.31) begins with a universal quantifier ( $\forall$ ), and its negation takes the form

$$\exists q \in \mathbb{Q} : \text{not} \left( \exists n \in \mathbb{N} : n > q \right).$$

In this form, after the first existential quantifier ( $\exists$ ) we have the negation of a statement which begins with an existential quantifier ( $\exists$ ), which we may further unravel. The negation of (I.31) is thus written as

$$\exists q \in \mathbb{Q} : \forall n \in \mathbb{N} : \text{not} (n > q).$$

The negation  $\text{not} (n > q)$  is obviously  $n \leq q$ , so the negation of (I.31) has been rewritten as

$$\exists q \in \mathbb{Q} : \forall n \in \mathbb{N} : n \leq q,$$

which in natural language claims that there exists some rational number  $q$  such that all natural numbers  $n$  are bounded from above by it. This clearly already sounds absurd, but we leave it to the reader to now derive the contradiction.<sup>22</sup> The point here was to illustrate how we typically end up considering negations of statements with quantifiers, and how they can be systematically unraveled.

## I.4. ♥ In depth topics about the set theory

The reader interested in deepening her understanding may wish to look into some of the following topics related to the subject of this lecture:

- The notion of cardinality as a way of comparing the sizes sets; in particular countable infinity as the smallest infinite cardinality and the uncountability of the set  $\mathbb{R}$  of real numbers, Appendix A.1.

The notions of countable and uncountable cardinalities will be crucial in various advanced topics of mathematics, notably in measure theory and probability theory. Familiarity with these notions will be assumed in the corresponding courses.

---

<sup>22</sup>Hint: If such a rational number  $q \in \mathbb{Q}$  exists, write it as  $q = \frac{m}{k}$  with  $m \in \mathbb{Z}$  and  $k \in \mathbb{N}$ . Then use for example the natural number  $n = |m| + 1$  as a counterexample to the statement  $\forall n \in \mathbb{N} : n \leq q$ .





## Lecture II

### Real numbers

There may not be much of a feeling of novelty to the real numbers, but we spend a lecture on this topic for two reasons: to appreciate some subtleties of real numbers themselves, and to get concrete examples of topological notions that will arise in more general context in the subsequent lectures.

We start by discussing the absolute value of real numbers and the related notion of distance on the real line, which is our first example of a metric, although the general definition will have to wait until Lecture V. This is the notion that underlies for example the convergence of sequences of real numbers. We look at basic properties of convergence as an instructive example of the more general theory to be developed later in Lecture VIII. We also show that both rational and irrational numbers are dense on the real line.

We then briefly turn to foundational questions again: what exactly are real numbers? The standard rigorous mathematical approach is to start from their axiomatic properties, most of which are entirely unsurprising. We only mention the other axioms in passing, but discuss the *completeness axiom* in more detail, since it is the most subtle and the most consequential for topology as well. It involves the notions of *supremum* (least upper bound) and *infimum* (greatest lower bound).

#### II.1. Absolute value, distances between numbers, and triangle inequality

At the core of topological considerations of numbers is the concept of distances on the real line. They are based on the absolute value — which itself is the distance of a given number to the origin.

The **absolute value** of a real number  $x \in \mathbb{R}$  is defined by

$$|x| = \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases} \quad (\text{II.1})$$

It is easy to see that another expression for it is

$$|x| = \sqrt{x^2}, \quad (\text{II.2})$$

where  $\sqrt{\cdot}$  denotes the non-negative square root of a non-negative number. Clearly the absolute value satisfies

$$|x| = |-x|, \quad |x| \geq 0, \quad x \leq |x|, \quad -x \leq |x|$$

for any  $x \in \mathbb{R}$ , and

$$|xy| = |x| |y|$$

for any  $x, y \in \mathbb{R}$ .

The **distance**<sup>1</sup> between two real numbers  $x, y \in \mathbb{R}$  is the absolute value of their difference,

$$|x - y|. \quad (\text{II.3})$$

The following **triangle inequality** is a simple and intuitive result, but as we will see later from Lecture V on, a rather straightforward generalization of it serves as the basis of a very fruitful general theory of metric spaces.

**Proposition II.1** (Triangle inequality on  $\mathbb{R}$ ).

*For all  $x, y \in \mathbb{R}$  we have*

$$\left| |x| - |y| \right| \leq |x + y| \leq |x| + |y|. \quad (\text{II.4})$$

*Proof.* Let  $x, y \in \mathbb{R}$ . There are two inequalities to prove in (II.4).

Let us start by proving the second inequality. Calculate the square of the absolute value of the sum, and estimate each term from above by its absolute value to get

$$\begin{aligned} |x + y|^2 &= (x + y)^2 = x^2 + 2xy + y^2 \\ &\leq |x|^2 + 2|x||y| + |y|^2 \\ &= (|x| + |y|)^2. \end{aligned}$$

Taking square roots ( $\sqrt{\cdot}$  is an increasing function and thus respects the inequality), we obtain that

$$|x + y| \leq |x| + |y|.$$

To prove the first inequality, we use the second one already proved above. Applying the second inequality to the numbers  $x + y$  and  $-y$ , on the other hand, gives

$$|x| = |(x + y) + (-y)| \leq |x + y| + |-y| = |x + y| + |y|,$$

and by subtracting  $|y|$  from both sides, we get

$$|x| - |y| \leq |x + y|.$$

A similar application of the second inequality to the numbers  $x + y$  and  $-x$  (or just interchanging the roles of  $x$  and  $y$  above) yields

$$|y| - |x| \leq |x + y|.$$

Combining these gives the first asserted inequality

$$\left| |x| - |y| \right| \leq |x + y|,$$

and the proof is complete.  $\square$

Another common form of the triangle inequality, which features the distance  $|x - y|$  in exactly this form is

$$|x - y| \leq |x| + |y|. \quad (\text{II.5})$$

It is obtained from the second inequality in (II.4) by plugging in  $x$  and  $-y$ , and noting  $|-y| = |y|$ .

There is a straightforward generalization of the triangle inequality to sums with finitely many many terms.

---

<sup>1</sup>Later, in Lecture V, we will see that this notion of a distance (II.3) is a special case of a *metric*, and it makes the real line a *metric space*. For now, however, we will work exclusively with the concrete case of real numbers, and we will not refer to the general notion.

**Proposition II.2** (Triangle inequality for finite sums).

Let  $x_1, \dots, x_n \in \mathbb{R}$ . Then we have

$$|x_1 + \dots + x_n| \leq |x_1| + \dots + |x_n|. \quad (\text{II.6})$$

**Exercise II.1** (Proof of Proposition II.2).

Prove Proposition II.2 by induction over  $n \in \mathbb{N}$ , using Proposition II.1.

**Remark II.3** (Triangle inequality for infinite series).

Let  $x_1, x_2, x_3, \dots \in \mathbb{R}$ . The natural generalization of (II.6) to infinite series would be

$$\left| \sum_{j=1}^{\infty} x_j \right| \stackrel{?}{\leq} \sum_{j=1}^{\infty} |x_j|.$$

With suitable interpretations, this generalization would indeed be valid. However, it immediately requires consideration of the convergence of the infinite series on both sides, and/or suitable interpretations in the case that one or both of the series diverge. For now, we will not try to use these types of generalizations, but we invite the reader to think about what can and cannot be said about it.

## II.2. Sequences on the real line

### Sequences of real numbers

A **sequence** (of real numbers) is an “infinite list”

$$(a_1, a_2, a_3, \dots)$$

of real numbers  $a_1, a_2, a_3, \dots \in \mathbb{R}$ . A precise definition of a sequence is a function

$$a: \mathbb{N} \rightarrow \mathbb{R},$$

and the “list” consists of the values of this function:  $a_n = a(n)$  for  $n \in \mathbb{N}$ . We usually use notations such as

$$(a_1, a_2, a_3, \dots) = (a_n)_{n=1}^{\infty} = (a_n)_{n \in \mathbb{N}}$$

for a sequence, and it is even common to use the lazy notation  $(a_n)$ , although this fails to explicitly indicate that the indexing of the members of the sequence uses natural numbers.

Occasionally we also let the indexing start from something other than  $n = 1$ . When we do so, this should be quite clear from the context. The notation is then modified in obvious ways: e.g.,  $(a_0, a_1, a_2, \dots) = (a_n)_{n=0}^{\infty}$  or  $(a_{25}, a_{26}, a_{27}, \dots) = (a_n)_{n=25}^{\infty}$ ; the precise interpretation is as functions  $\{0, 1, 2, \dots\} \rightarrow \mathbb{R}$  or  $\{25, 26, 27, \dots\} \rightarrow \mathbb{R}$ . The definitions in the following subsections must then be adapted in obvious ways. Some properties of interest (particularly limits) in fact do not even depend on any finitely many initial members of the sequence, in which case the choice of the starting index is largely irrelevant.

### Monotonicity properties of number sequences

The real axis has an order. Sequences which respect that order have some particularly nice properties.

**Definition II.4** (Monotonicity properties of sequences of real numbers).

A real number sequence  $(a_n)_{n \in \mathbb{N}}$  is

- **increasing**, if we have<sup>2</sup>  $a_{n+1} \geq a_n$  for all  $n \in \mathbb{N}$
- **strictly increasing**, if we have<sup>3</sup>  $a_{n+1} > a_n$  for all  $n \in \mathbb{N}$
- **decreasing**, if we have<sup>4</sup>  $a_{n+1} \leq a_n$  for all  $n \in \mathbb{N}$
- **strictly decreasing**, if we have<sup>5</sup>  $a_{n+1} < a_n$  for all  $n \in \mathbb{N}$
- **monotone**, if it is either increasing or decreasing.

**Example II.5** (An increasing sequence).

The sequence  $(1, 2, 4, 8, 16, 32, \dots)$  is (strictly) increasing: its  $n$ :th member is given by the formula  $a_n = 2^{n-1}$ , and we have  $a_{n+1} = 2^n > 2^{n-1} = a_n$  for all  $n \in \mathbb{N}$ .

**Example II.6** (Another increasing sequence).

Let  $(a_n)_{n \in \mathbb{N}}$  be defined by the formula  $a_n = \frac{n}{n+1}$  for  $n \in \mathbb{N}$ . We claim that this sequence is increasing.

One way to verify this is to calculate the difference of consecutive members,

$$a_{n+1} - a_n = \frac{n+1}{n+2} - \frac{n}{n+1} = \frac{(n+1)^2 - n(n+2)}{(n+2)(n+1)} = \frac{1}{(n+2)(n+1)} > 0,$$

which gives  $a_{n+1} > a_n$  (by adding  $a_n$  to both sides in the inequality).

Another (perhaps easier?) way is to calculate the ratio of consecutive terms

$$\frac{a_n}{a_{n+1}} = \frac{n/(n+1)}{(n+1)/(n+2)} = \frac{n(n+2)}{(n+1)^2} = \frac{n^2 + 2n}{n^2 + 2n + 1} < 1,$$

which gives  $a_n < a_{n+1}$  (by multiplying both sides by the positive number  $a_{n+1}$ ).

**Example II.7** (A decreasing sequence).

The sequence  $(1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \frac{1}{25}, \dots)$  is (strictly) decreasing: its  $n$ :th member is given by the formula  $a_n = \frac{1}{n^2}$ , and we have  $a_{n+1} = \frac{1}{(n+1)^2} < \frac{1}{n^2} = a_n$  for all  $n \in \mathbb{N}$ .

## Boundedness properties of number sequences

**Definition II.8** (Boundedness properties of sequences of real numbers).

A real number sequence  $(a_n)_{n \in \mathbb{N}}$  is

- **bounded from above**, if there exists some  $u \in \mathbb{R}$  (called an **upper bound** for the sequence) such that  $a_n \leq u$  for all  $n \in \mathbb{N}$ ;
- **bounded from below**, if there exists some  $\ell \in \mathbb{R}$  (called a **lower bound** for the sequence) such that  $a_n \geq \ell$  for all  $n \in \mathbb{N}$ ;
- **bounded**, if there exists some  $r \in \mathbb{R}$  such that  $|a_n| \leq r$  for all  $n \in \mathbb{N}$ .

**Exercise II.2** (Boundedness from both above and below).

Prove that a real number sequence  $(a_n)_{n \in \mathbb{N}}$  is bounded if and only if it is both bounded from above and bounded from below.

**Example II.9** (A sequence bounded from below).

The sequence  $(1, 2, 4, 8, 16, 32, \dots)$  is bounded from below by, for example, the constant

<sup>2</sup>In other words, a sequence is increasing if  $a_1 \leq a_2 \leq a_3 \leq \dots$ .

<sup>3</sup>In other words, a sequence is strictly increasing if  $a_1 < a_2 < a_3 < \dots$ .

<sup>4</sup>In other words, a sequence is decreasing if  $a_1 \geq a_2 \geq a_3 \geq \dots$ .

<sup>5</sup>In other words, a sequence is strictly decreasing if  $a_1 > a_2 > a_3 > \dots$ .

$\ell = 1$ .<sup>6</sup> To see this, note that the  $n$ :th member of the sequence is given by the formula  $a_n = 2^{n-1}$ , and we have  $a_n = 2^{n-1} \geq 1 = \ell$  for all  $n \in \mathbb{N}$ .

This sequence is not bounded from above. Indeed for any  $u \in \mathbb{R}$ , we can take a natural number  $n > \log_2(u) + 1$ , and then  $a_n = 2^{n-1} > 2^{\log_2(u)} = u$ , so  $u$  does not work as an upper bound. Since  $u \in \mathbb{R}$  was arbitrary, no upper bound exists for this sequence.

**Example II.10** (A bounded sequence).

Let  $(a_n)_{n \in \mathbb{N}}$  be defined by the formula  $a_n = \frac{n}{n+1}$  for  $n \in \mathbb{N}$ . We claim that this sequence is bounded by, for example, the constant  $r = 1$ .<sup>7</sup> To see this, note that for any  $n \in \mathbb{N}$  we have

$$|a_n| = \left| \frac{n}{n+1} \right| = \frac{n}{n+1} \leq 1 = r.$$

### II.3. Limits of sequences on the real line

#### The definition of the limit of a sequence

**Definition II.11** (Limit of a sequence of numbers).

A sequence  $(a_n)_{n \in \mathbb{N}}$  of real numbers **converges** to a **limit**  $\alpha \in \mathbb{R}$  if for every  $\varepsilon > 0$  there exists an index  $n_\varepsilon \in \mathbb{N}$  such that  $|a_n - \alpha| < \varepsilon$  whenever  $n \geq n_\varepsilon$ .

**Remark II.12** (Meaning of convergence as a logical statement).

Definition II.11 is written in plain English (a natural language), as usual. It nevertheless has a precise logical meaning, and it is instructive to unravel the definition using logical symbols (a formal language). The meaning of the statement  $a_n \rightarrow \alpha$  as  $n \rightarrow \infty$  is:

$$\forall \varepsilon > 0 : \quad \exists n_\varepsilon \in \mathbb{N} : \quad n \geq n_\varepsilon \implies |a_n - \alpha| < \varepsilon. \quad (\text{II.7})$$

A few advantages of the formal statement (II.7) are that it is succinct, unambiguous, and understandable to mathematicians irrespective of whether their mother tongue is English, Finnish, Swedish, or some other natural language (being familiar with the logical symbols, you can easily “read it” in your native language).

The logical definition is crucial, because it gives the precise meaning to the word *limit* (and *converge*). But of course it is still also important to have an intuitive idea about the notion as well! So how should you think about Definition II.11? In Figure II.1 and its caption we attempt to illustrate and describe this.

**Exercise II.3** (Non-zero limit implies members are eventually non-zero).

Suppose that  $(a_n)_{n \in \mathbb{N}}$  is a sequence of real numbers which tends to a non-zero limit

$$\alpha = \lim_{n \rightarrow \infty} a_n \neq 0.$$

Show that there exists some  $N \in \mathbb{N}$  such that  $a_n \neq 0$  for all  $n \geq N$ .

Hint: We have  $|a_n| = |\alpha - (\alpha - a_n)| \geq |\alpha| - |\alpha - a_n|$  by triangle inequality, Proposition II.1.

#### Basic properties of limits of sequences

Having given the precise definition of limit, we are ready to state and prove some basic properties of limits.

<sup>6</sup>As a lower bound here we could equally well use  $\ell = 0$  or  $\ell = -7$  or indeed any number  $\ell \leq 1$ .

<sup>7</sup>As a bound here we could equally well use  $r = 42$  or  $r = 10^{23}$  or indeed any number  $r \geq 1$ .



FIGURE II.1. An illustration of the limit of a number sequence. The limit of the sequence  $(a_n)_{n \in \mathbb{N}}$  is  $\alpha \in \mathbb{R}$  if for however small error  $\varepsilon > 0$  we are willing to tolerate, the entire tail of the sequence starting from some index  $n_\varepsilon \in \mathbb{N}$  lies within that error range from the value  $\alpha$ ; i.e., for all  $n \geq n_\varepsilon$  we have  $\alpha - \varepsilon < a_n < \alpha + \varepsilon$ .

The first one says that the property of convergence implies that of boundedness.

**Proposition II.13** (A convergent sequence of number is bounded).

*If a sequence  $(a_n)_{n \in \mathbb{N}}$  of real numbers converges, then it is bounded.*

**Exercise II.4** (Proof of Proposition II.13).

Prove Proposition II.13.

**Exercise II.5** (Boundedness does not imply convergence).

Show that there exists a bounded sequence which is not convergent. Conclude that the implication in the converse direction compared to Proposition II.13 does not hold.

**Lemma II.14** (Preservation of inequalities).

*Let  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  be two convergent sequences of real numbers, with respective limits  $\alpha = \lim_{n \rightarrow \infty} a_n$  and  $\beta = \lim_{n \rightarrow \infty} b_n$ . If we have*

$$a_n \leq b_n \quad \text{for all } n \in \mathbb{N},$$

*then we also have the corresponding inequality*

$$\alpha = \lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} b_n = \beta$$

*for the limits.*

*Proof.* Let  $\varepsilon > 0$ . Since  $\frac{\varepsilon}{2} > 0$ , it follows from the assumptions  $\lim_{n \rightarrow \infty} a_n = \alpha$  and  $\lim_{n \rightarrow \infty} b_n = \beta$  and the definition of limits that there exists numbers  $n', n'' \in \mathbb{N}$  such that

$$|a_n - \alpha| < \frac{\varepsilon}{2} \quad \text{for } n \geq n', \quad |b_n - \beta| < \frac{\varepsilon}{2} \quad \text{for } n \geq n''.$$

Consider now  $n = \max \{n', n''\} \in \mathbb{N}$ . Then  $n \geq n'$  and  $n \geq n''$ , so both of the above hold. Using furthermore the assumption  $a_n \leq b_n$ , we find

$$\begin{aligned} 0 \leq b_n - a_n &= b_n - \beta + \beta - \alpha + \alpha - a_n \\ &\leq |b_n - \beta| + \beta - \alpha + |\alpha - a_n| \\ &< \frac{\varepsilon}{2} + \beta - \alpha + \frac{\varepsilon}{2} \\ &= \beta - \alpha + \varepsilon. \end{aligned}$$

Rearranging this inequality, we get  $\alpha < \beta + \varepsilon$ . Since this was shown to hold for an arbitrary  $\varepsilon > 0$ , it must be that  $\alpha \leq \beta$ .<sup>8</sup>  $\square$

**Exercise II.6** (An improvement to Lemma II.14).

Show that in Lemma II.14 it suffices to assume that the inequality holds for all sufficiently large  $n$ ; more precisely that there exists  $N \in \mathbb{N}$  such that  $a_n \leq b_n$  for all  $n \geq N$ .

For sequences bounded from above or below, if a limit exists, it must also lie within the same bounds.

**Corollary II.15** (Preservation of bounds).

Let  $(a_n)_{n \in \mathbb{N}}$  be a convergent sequences of real numbers, with limit  $a = \lim_{n \rightarrow \infty} a_n$ . If for some  $u \in \mathbb{R}$  we have

$$a_n \leq u \quad \text{for all } n \in \mathbb{N},$$

then we also have the corresponding inequality for the limit,

$$\alpha = \lim_{n \rightarrow \infty} a_n \leq u.$$

Similarly, if for some  $\ell \in \mathbb{R}$  we have  $a_n \geq \ell$  for all  $n \in \mathbb{N}$ , then we also have  $\lim_{n \rightarrow \infty} a_n \geq \ell$ .

**Exercise II.7** (Proof of Corollary II.15).

Prove the first part of Corollary II.15, by using a constant sequence  $(b_n)_{n \in \mathbb{N}}$  in Lemma II.14. Prove the second part by a similar idea.

**Exercise II.8** (An improvement to Corollary II.15).

Formulate the following improvement precisely and prove it:

In Corollary II.15 it suffices to assume that the inequality holds for all sufficiently large  $n$ .

Hint: Compare with Exercise II.6.

Note that from Definition II.11 it is *not immediately obvious* that there could not exist several different numbers that satisfy what is required of a limit. In other words, if the condition (II.7) holds for  $\alpha \in \mathbb{R}$ , then could it also hold for some other

---

<sup>8</sup>This is a very common step in proofs, so let us look at it in detail now that we first use it. The claim is that

$$\forall \varepsilon > 0 : \alpha < \beta + \varepsilon \quad \implies \quad \alpha \leq \beta.$$

This is easy to prove by *contrapositive*: the claim is logically equivalent to (recall contrapositives and negations of statements with universal quantifier)

$$\alpha > \beta \quad \implies \quad \exists \varepsilon > 0 : \alpha \geq \beta + \varepsilon$$

In this formulation the claim is evident: if  $\alpha > \beta$ , then  $\alpha - \beta > 0$  and we can use  $\varepsilon = \alpha - \beta$  which indeed satisfies  $\alpha = \beta + \varepsilon \geq \beta + \varepsilon$ .

real number  $\alpha' \in \mathbb{R}$  in place of  $\alpha$ ?<sup>9</sup> Fortunately the following result lifts any concerns about the possibility of such ambiguity.<sup>10</sup>

**Corollary II.16** (Uniqueness of limits of real number sequences).

*Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence of real numbers. If both  $\alpha \in \mathbb{R}$  and  $\alpha' \in \mathbb{R}$  are limits of this sequence, then  $\alpha = \alpha'$ .*

*Proof.* Trivially, the inequality  $a_n \leq a_n$  holds for all  $n \in \mathbb{N}$ . If we now suppose that both  $\alpha$  and  $\alpha'$  are limits of the sequence  $(a_n)_{n \in \mathbb{N}}$ , and we use the former limit for the left hand side and the latter for the right hand side, from Lemma II.14 we get that  $\alpha \leq \alpha'$ . Similarly using the former for the right hand side and the latter for the left hand side, we get  $\alpha' \leq \alpha$ . The combination of these gives  $\alpha = \alpha'$ .  $\square$

We next state a very practical result, which is in some sense an improved version of the preservation of inequalities — in it we do not even need to assume the existence of the limit, but the existence is a part of the conclusion! It states that if a sequence is “squeezed” in between two convergent sequences with same limit, then the sequence itself has to converge to this limit, too. This is often called *the squeeze theorem* (though we regard it more as a lemma). It also has various other affectionate nicknames. Some call it *the sandwich principle* — since it talks about a sequence “sandwiched” in between two others. Also *the lemma of two policemen* is descriptive: the idea being that one sequence “guards” the sequence of interest from above, preventing its escape to the upwards direction, and another sequence “guards” the sequence of interest from below, preventing its escape to the downwards direction.<sup>11</sup>

**Lemma II.17** (Squeeze theorem).

*Let  $(a_n)_{n \in \mathbb{N}}$ ,  $(b_n)_{n \in \mathbb{N}}$ ,  $(c_n)_{n \in \mathbb{N}}$  be three sequences of real numbers. Suppose that for all  $n \in \mathbb{N}$  we have*

$$a_n \leq b_n \leq c_n.$$

*Suppose also that the sequences  $(a_n)_{n \in \mathbb{N}}$  and  $(c_n)_{n \in \mathbb{N}}$  are convergent and have the same limit*

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = \beta \in \mathbb{R}.$$

*Then the sequence  $(b_n)_{n \in \mathbb{N}}$  is also convergent and its limit is the same,*

$$\lim_{n \rightarrow \infty} b_n = \beta.$$

**Exercise II.9** (Proof of Lemma II.17).

Prove the squeeze theorem (Lemma II.17).

**Example II.18** (An example of the squeeze theorem).

Consider the sequence  $(b_n)_{n \in \mathbb{N}}$  given by

$$b_n = 3 + 4^{-n} \sin(5n) \quad \text{for } n \in \mathbb{N}.$$

<sup>9</sup>The notation  $\lim_{n \rightarrow \infty} a_n$  suggests that the limit is uniquely determined, but we should actually prove that this is so, in order to be sure that the notation is unambiguously defined!

<sup>10</sup>It would not have been difficult to prove the uniqueness of limits more directly, immediately after the definition. In fact, in Chapter VIII we treat sequences and limits in a more general setup, and we give a more direct (as well as more general) proof of the uniqueness. Here, however, we think it instructive to give a proof based on the preservation of inequalities.

<sup>11</sup>Clearly the result must be important, given how imaginative nomenclature it has inspired!



Since  $-1 \leq \sin(\theta) \leq +1$  for all  $\theta \in \mathbb{R}$ , we get

$$3 - 4^{-n} \leq b_n \leq 3 + 4^{-n} \quad \text{for all } n \in \mathbb{N}.$$

We have  $\lim_{n \rightarrow \infty} (3 - 4^{-n}) = 3$  and  $\lim_{n \rightarrow \infty} (3 + 4^{-n}) = 3$ , so we can use the sequences  $(a_n)_{n \in \mathbb{N}}$  and  $(c_n)_{n \in \mathbb{N}}$  defined by  $a_n = 3 - 4^{-n}$  and  $c_n = 3 + 4^{-n}$  in the squeeze theorem (Lemma II.17) to conclude that

$$\lim_{n \rightarrow \infty} b_n = 3.$$

### Rules of calculation with limits

Given two real number sequences  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$ , we often consider new sequences formed from these for example by<sup>12</sup>

$$(a_n + b_n)_{n \in \mathbb{N}}, \quad (a_n b_n)_{n \in \mathbb{N}}, \quad (a_n/b_n)_{n \in \mathbb{N}}.$$

**Theorem II.19** (Rules of calculation with limits).

Let  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  be two convergent sequences with respective limits  $\alpha = \lim_{n \rightarrow \infty} a_n$  and  $\beta = \lim_{n \rightarrow \infty} b_n$ . Then we have

$$\lim_{n \rightarrow \infty} (a_n + b_n) = \alpha + \beta, \quad (\text{II.8})$$

$$\lim_{n \rightarrow \infty} (a_n b_n) = \alpha \beta, \quad (\text{II.9})$$

and if moreover  $\beta \neq 0$ , then

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\alpha}{\beta}. \quad (\text{II.10})$$

Before addressing the proof, we note the following. From (II.8) and (II.9) above, we get the following special cases by choosing  $(b_n)_{n \in \mathbb{N}}$  to be a constant sequence  $b_n = c$  for all  $n \in \mathbb{N}$ .

**Corollary II.20** (Additive and multiplicative constants in limits).

Let  $(a_n)_{n \in \mathbb{N}}$  be a convergent sequence with limit  $\alpha = \lim_{n \rightarrow \infty} a_n$ , and let  $c \in \mathbb{R}$  be a constant. Then we have

$$\lim_{n \rightarrow \infty} (a_n + c) = \alpha + c, \quad \lim_{n \rightarrow \infty} (c a_n) = c \alpha.$$

*Proof of Theorem II.19.* The proofs of all cases are quite similar, so we only do (II.9) and leave the other two as exercises.

So suppose that  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  are two convergent sequences with limits  $\alpha = \lim_{n \rightarrow \infty} a_n$  and  $\beta = \lim_{n \rightarrow \infty} b_n$ . We must prove that the sequence  $(a_n b_n)_{n \in \mathbb{N}}$  converges to  $\alpha \beta$ , and we

---

<sup>12</sup>The last of these is only well defined if  $b_n \neq 0$  for all  $n \in \mathbb{N}$  — otherwise we encounter a division by zero in  $a_n/b_n$ . But in fact, since we will be concerned with calculating limits as  $n \rightarrow \infty$ , we do not necessarily need to care if such a problem occurs just finitely many times along the sequence. If there are only finitely many indices  $n \in \mathbb{N}$  for which  $b_n = 0$ , then we can still consider the sequence  $(a_n/b_n)_{n=n_0}^\infty$  starting from a sufficiently large index  $n_0$  such that  $b_n \neq 0$  for all  $n \geq n_0$ . It is meaningful to consider the limit of this sequence, which we still write as  $\lim_{n \rightarrow \infty} a_n/b_n$ .

will do this directly using the Definition II.11 of limits. For this purpose, we first estimate

$$\begin{aligned}
 & |a_n b_n - \alpha \beta| \\
 &= |a_n b_n - \alpha b_n + \alpha b_n - \alpha \beta| \quad (\text{added and subtracted the same term}) \\
 &= |(a_n - \alpha) b_n + \alpha (b_n - \beta)| \quad (\text{rearranged terms}) \\
 &\leq |(a_n - \alpha) b_n| + |\alpha (b_n - \beta)| \quad (\text{triangle inequality}) \\
 &= |b_n| |a_n - \alpha| + |\alpha| |b_n - \beta|.
 \end{aligned}$$

By Proposition II.13, the convergent sequence  $(b_n)_{n \in \mathbb{N}}$  is bounded, so there exists a constant  $M > 0$  such that  $|b_n| \leq M$  for all  $n \in \mathbb{N}$ . Plugging this in the above estimate, we obtain<sup>13</sup>

$$|a_n b_n - \alpha \beta| \leq M |a_n - \alpha| + |\alpha| |b_n - \beta|. \quad (\text{II.11})$$

Now let  $\varepsilon > 0$ . Because  $\lim_{n \rightarrow \infty} a_n = \alpha$  and  $\frac{\varepsilon}{3M}$  is a positive number, there exists an  $n' \in \mathbb{N}$  such that we have

$$|a_n - \alpha| < \frac{\varepsilon}{3M} \quad \text{for } n \geq n'.$$

Similarly, because  $\lim_{n \rightarrow \infty} b_n = \beta$  and  $\frac{\varepsilon}{3(|\alpha|+1)}$  is a positive number, there exists an  $n'' \in \mathbb{N}$  such that we have

$$|b_n - \beta| < \frac{\varepsilon}{3(|\alpha|+1)} \quad \text{for } n \geq n''.$$

Then if  $n \geq \max\{n', n''\}$ , we have both  $n \geq n'$  and  $n \geq n''$ , so both of the above estimates hold, and from (II.11) we get

$$\begin{aligned}
 |a_n b_n - \alpha \beta| &\leq M \underbrace{|a_n - \alpha|}_{\leq \varepsilon/3M} + |\alpha| \underbrace{|b_n - \beta|}_{\leq \varepsilon/3(|\alpha|+1)} \\
 &\leq \underbrace{\frac{M}{3M}}_{\leq 1/3} \varepsilon + \underbrace{\frac{|\alpha|}{3(|\alpha|+1)}}_{\leq 1/3} \varepsilon \\
 &\leq \frac{2}{3} \varepsilon < \varepsilon.
 \end{aligned}$$

If we set  $n_\varepsilon = \max\{n', n''\}$ , then from the above we got  $|a_n b_n - \alpha \beta| < \varepsilon$  for all  $n \geq n_\varepsilon$ . Since  $\varepsilon > 0$  was arbitrary, this by definition shows that  $\lim_{n \rightarrow \infty} (a_n b_n) = \alpha \beta$ , proving (II.9).

The proofs of (II.8) and (II.10) are left as exercises.  $\square$

**Exercise II.10** (Completing the proof of Theorem II.19).  
Prove (II.8) and (II.10).

**Exercise II.11** (Examine properties of some sequences).  
Let

$$a_n = \frac{n+1}{2n+1} \quad \text{and} \quad b_n = \sqrt[n]{3^n + 2^n} \quad \text{for } n \in \mathbb{N}.$$

Find out whether the sequences  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  are increasing, decreasing, bounded from above, bounded from below. Determine the limits of these sequence if they exist.

<sup>13</sup>After this, the idea is that assumed convergence of  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  implies that both  $|a_n - \alpha|$  and  $|b_n - \beta|$  become small for  $n$  large. This should allow us to show that the expression (II.11) becomes small for large  $n$ . The rest of the proof is about making this idea precise.

## II.4. Density of rational and irrational numbers

We will next look at an interesting property of the real axis: both the rational numbers and the irrational numbers are dense on  $\mathbb{R}$ . In particular, between any two rational numbers there is an irrational number, and between any two irrational number there is a rational number. The fact that the countably infinite (and thus set theoretically rather small) set of rational numbers is already dense in the (uncountably infinite) set of real numbers is a topological property of  $\mathbb{R}$  known as *separability*.

### Density of rational numbers

Let us show that between any two (different) real numbers, there exists a rational number.

**Theorem II.21** (Density of rational numbers).

*Let  $x, y \in \mathbb{R}$  with  $x < y$ . Then there exists a  $q \in \mathbb{Q}$  such that  $x < q < y$ .*

*Proof.* Since  $y - x > 0$ , we can choose an  $n \in \mathbb{N}$  such that  $n > \frac{1}{y-x}$ . Now in the set  $\{\frac{m}{n} \mid m \in \mathbb{Z}\} \subset \mathbb{Q}$  of all integer multiples of  $\frac{1}{n}$ , the distance between consecutive ones is  $\frac{1}{n}$ . In particular since the interval  $(x, y) \subset \mathbb{R}$  has length  $y - x > \frac{1}{n}$ , at least one such integer multiple  $\frac{m}{n}$  must lie in this interval. In other words, the rational number  $q = \frac{m}{n} \in \mathbb{Q}$  lies on the desired interval,  $q \in (x, y)$ .  $\square$

Although we only apparently showed that an open interval  $(x, y) \subset \mathbb{R}$  of the real axis contains one rational number, it almost immediately follows that the interval in fact contains infinitely many rational numbers.

**Corollary II.22** (An open interval contains infinitely many rational numbers).

*On any nonempty open interval  $(x, y) \subset \mathbb{R}$ , there are infinitely many rational numbers, i.e.,  $\#((x, y) \cap \mathbb{Q}) = \infty$ .*

*Proof.* By Theorem II.21, we may first choose one rational number  $q_1 \in \mathbb{Q}$  such that  $x < q_1 < y$ . Then applying the same result to the interval  $(x, q_1)$ , we can choose  $q_2 \in \mathbb{Q}$  such that  $x < q_2 < q_1 < y$ . Inductively, once  $q_1, \dots, q_n \in \mathbb{Q}$  have been chosen so that

$$x < q_n < q_{n-1} < \dots < q_2 < q_1 < y,$$

we choose a rational number  $q_{n+1} \in \mathbb{Q}$  from the interval  $(x, q_n)$ . This yields an infinite sequence  $(q_n)_{n \in \mathbb{N}}$  of rational numbers on the interval  $(x, y)$ . Finally note that by construction the sequence is strictly decreasing, so no two members of it are equal, and we have indeed found infinitely many different rational numbers on the interval.  $\square$

### Density of irrational numbers

Similarly, between any two (different) real numbers, there exists an irrational number.

**Theorem II.23** (Density of irrational numbers).

*Let  $x, y \in \mathbb{R}$  with  $x < y$ . Then there exists a  $z \in \mathbb{R} \setminus \mathbb{Q}$  such that  $x < z < y$ .*

*Proof.* Let us first use Corollary II.22 to choose two different rational numbers  $q, r \in \mathbb{Q}$  so that  $x < q < r < y$ . It suffices to show that the subinterval  $(q, r) \subset (x, y)$  contains an irrational number.

Let

$$z = q + \frac{r - q}{\sqrt{2}}.$$

Clearly we have  $z > q$ , since  $r - q > 0$  and  $\frac{1}{\sqrt{2}} > 0$ . On the other hand  $\frac{1}{\sqrt{2}} < 1$ , so we also have

$$z = q + \frac{r - q}{\sqrt{2}} < q + 1 \cdot (r - q) = r.$$

From these we get that the number  $z$  lies in the subinterval of interest,  $x < q < z < r < y$ . It remains to show that  $z$  is irrational. From the definition of  $z$ , we can solve

$$\sqrt{2} = \frac{r - q}{z - q}.$$

Now argue by contradiction: if  $z$  were rational, then the right hand side above would also be rational (recall that  $q, r$  were chosen rational), so  $\sqrt{2}$  would be rational, contradicting Proposition I.5. We conclude the desired irrationality  $z \in \mathbb{R} \setminus \mathbb{Q}$ .  $\square$

Again from knowing that an open interval  $(x, y) \subset \mathbb{R}$  of the real axis contains one irrational number, we easily get that the interval in fact contains infinitely many irrational numbers.

**Corollary II.24** (An open interval contains infinitely many irrational numbers).

*On any nonempty open interval  $(x, y) \subset \mathbb{R}$ , there are infinitely many irrational numbers, i.e.,  $\#((x, y) \setminus \mathbb{Q}) = \infty$ .*

**Exercise II.12** (Proof of Corollary II.24).

Prove Corollary II.24, using Theorem II.23 and an argument similar to Corollary II.22.

## II.5. Axioms of the real numbers

So far we have talked about real numbers, but we have not in fact precisely said what they are! If one looks closely, this actually turns out not to be an entirely trivial philosophical matter...

As with many other mathematical structures, the standard rigorous approach is to take certain properties of real numbers as axioms (statements that are simply accepted as true), and require that any other statement made about real numbers is logically deduced from these axioms.<sup>14 15</sup> We want the axioms to be as modest

<sup>14</sup>In practice, it would be very tedious to give full proofs of all properties of real numbers that we use, as this would include much of what you have learned about mathematics since kindergarten... We will therefore not ask you to provide proofs of totally commonplace statements about the real numbers — but instead to just realize that they must in principle be logical consequences of the axioms. In Appendix B.1 all of the axioms of  $\mathbb{R}$  are at least stated, and derivations of other properties from them are exemplified.

<sup>15</sup>In addition to giving the axioms, it would be desirable to provide a (set theoretic) construction of real numbers, for which the axioms can be shown to be true. Suffice to say that there are such constructions, but we will not discuss them in detail here. Any such construction of course will admit something else as a starting point. An apparently benign starting point is to assume “uncontroversial” axioms of natural numbers (but in view of Gödel’s incompleteness theorem this is in fact not without its own issues!).

as possible, so that the starting point is uncontroversial. And since everything else is deduced from the starting point by logical reasoning, also everything else will be uncontroversial — which of course is the point of mathematics!

The set of real numbers is denoted by  $\mathbb{R}$ , and it is equipped with two binary operations:

$$\begin{aligned} \text{addition “+”}: \quad & \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \\ & (x, y) \mapsto x + y, \\ \text{multiplication “.”}: \quad & \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \\ & (x, y) \mapsto x \cdot y \\ & (\text{omitted in usual notation, so } x \cdot y = xy), \end{aligned}$$

and a binary relation

less than “ $<$ ”.

The axioms of real numbers fall into three types: the *field axioms* concern only the operations of addition and multiplication, the *order axioms* concern the order relation  $<$  and how it behaves under addition and multiplication, and finally the *completeness axiom* which is in many ways the most subtle.

### On the field axioms and order axioms

The field axioms and order axioms concern properties of the real numbers that are without a doubt familiar already. We list these axioms and briefly discuss them in Appendix B.1. For a broader mathematical perspective, it is worthwhile to take a look and to appreciate also these familiar properties of addition, multiplication, and order.

Here we content ourselves to mentioning that there are various other *fields* regularly used in mathematics — for example:

- the field of rational numbers  $\mathbb{Q}$ ;
- the field of complex numbers  $\mathbb{C}$ ;
- finite fields, the simplest example of which is the field  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  of  $p$  elements, where  $p$  is a prime number;
- algebraic number fields, a simple example of which is the field  $\mathbb{Q}(\sqrt{33})$  of rational numbers adjoined with a square root of 33;
- the field  $\mathbb{Q}_p$  of  $p$ -adic numbers, where  $p$  is a prime number;
- the field  $\mathbb{K}(q)$  of rational functions in a single variable  $q$  over another field  $\mathbb{K}$ ;
- ...

All of these satisfy the same *field axioms* as  $\mathbb{R}$ , so the rules governing addition and multiplication with all their consequences are similar.

---

From the natural numbers, it is not difficult to construct the rational numbers. From the rational numbers, the real numbers can be constructed for example using what is known as Dedekind cuts. There are many other constructions as well. Importantly, however, a sort of uniqueness property of real numbers holds: any two constructions (in which the axioms hold) yield results that are in all essential ways the same (isomorphic). So we do not need to care about the chosen construction — the axioms of real numbers themselves are indeed a good starting point.

The field  $\mathbb{Q}$  of rational numbers (as well as some but not all algebraic number fields) moreover has an order relation compatible with the addition and multiplication, in that it satisfies also the same *order axioms* as  $\mathbb{R}$ . The “only” difference between the real numbers and the rational numbers must therefore stem from the *completeness axiom*, which we discuss in more detail below.

### Completeness axiom

We will next address to the one really subtle axiom of the real numbers: the *completeness axiom*. But before stating it, we first need to introduce a few notions that will make an appearance here.

#### *Supremum and infimum*

Let us start from notions of boundedness for subsets of the real line. If you compare the following with Definition II.8, a common theme becomes evident.

**Definition II.25** (Upper and lower bounds).

Let  $A \subset \mathbb{R}$  be a subset.

A real number  $u \in \mathbb{R}$  is an **upper bound** for  $A$  if for all  $a \in A$  we have  $a \leq u$ .

A real number  $\ell \in \mathbb{R}$  is a **lower bound** for  $A$  if for all  $a \in A$  we have  $a \geq \ell$ .

**Example II.26** (Examples of upper and lower bounds).

- Consider the open interval  $(-\pi, +\pi) \subset \mathbb{R}$ . The number  $\pi$  is an upper bound for the set  $(-\pi, +\pi)$ . But also any number  $u > \pi$  is an upper bound for  $(-\pi, +\pi)$ . Likewise, any number  $\ell \leq -\pi$  is a lower bound for the set  $(-\pi, +\pi)$ .
- Consider the set  $\mathbb{N} \subset \mathbb{R}$  of natural numbers. The number 1 is a lower bound for  $\mathbb{N}$ , and so is any number  $\ell < 1$ . The set  $\mathbb{N}$  does not have any upper bounds: for any  $u \in \mathbb{R}$  there exists a natural number  $n \in \mathbb{N}$  such that  $n > u$ .
- The sets  $\mathbb{Z}$ ,  $\mathbb{Q}$ , and  $\mathbb{R}$  do not have any upper or lower bounds.
- Consider the empty set  $\emptyset \subset \mathbb{R}$ . Any real number  $u \in \mathbb{R}$  is an upper bound for  $\emptyset$ .<sup>16</sup> Likewise, any real number  $\ell \in \mathbb{R}$  is a lower bound for  $\emptyset$ .

From these examples we learn that upper and lower bounds for arbitrary sets do not need to exist. Moreover, when they exist, they are not unique.

**Definition II.27** (Sets bounded from above and below).

A subset  $A \subset \mathbb{R}$  is **bounded from above** if it has an upper bound, and **bounded from below** if it has a lower bound.

The following definition introduces the key notion in the completeness axiom: the least upper bound or supremum. The supremum and its counterpart infimum also feature in a number of contexts in real analysis — some applications are discussed in Appendix B.4.

---

<sup>16</sup>Indeed, there are no elements  $a \in \emptyset$  for which we would get a nontrivial condition that  $u$  needs to satisfy to qualify as an upper bound for the empty set.

**Definition II.28** (Supremum and infimum).

Let  $A \subset \mathbb{R}$  be a nonempty subset ( $A \neq \emptyset$ ) which is bounded from above. A number  $u_0 \in \mathbb{R}$  is called the **least upper bound** or **supremum** of  $A$ , if  $u_0$  is an upper bound for  $A$  (i.e.  $\forall a \in A : a \leq u_0$ ) and for all upper bounds  $u$  of  $A$  we have  $u_0 \leq u$ . We then denote  $u_0 = \sup A$ .

Let  $B \subset \mathbb{R}$  be a nonempty subset ( $B \neq \emptyset$ ) which is bounded from below. A number  $\ell_0 \in \mathbb{R}$  is called the **greatest lower bound** or **infimum** of  $B$ , if  $\ell_0$  is a lower bound for  $B$  (i.e.  $\forall b \in B : b \geq \ell_0$ ) and for all lower bounds  $\ell$  of  $B$  we have  $\ell_0 \geq \ell$ . We then denote  $\ell_0 = \inf B$ .

Implicit in the notations  $\sup A$  and  $\inf A$  (and in our use of the definite article above) is that the least upper bound and the greatest lower bound are uniquely determined by the above properties. This is indeed not difficult to check, and we leave it as an exercise to the reader.

The notions of supremum and infimum are crucial in this course as well as mathematics more generally, so let us give a more practical characterization.

**Lemma II.29** (Characterization of supremum).

*A number  $u_0 \in \mathbb{R}$  is the supremum of a nonempty subset  $A \subset \mathbb{R}$  which is bounded from above, if and only if the following two conditions hold:*

- For all  $a \in A$  we have  $a \leq u_0$ .<sup>17</sup>
- For all  $\varepsilon > 0$  there exists some  $a \in A$  such that  $a > u_0 - \varepsilon$ .<sup>18</sup>

**Exercise II.13** (Proof of Lemma II.29).

Prove Lemma II.29.

Hint: Compare the defining properties of supremum in Definition II.28 with the interpretations of the two conditions in the lemma, given in the footnotes.

**Exercise II.14** (Characterization of infimum).

Formulate and prove a the counterpart of Lemma II.29 for infimum, i.e., two conditions characterizing the infimum of a non-empty set  $B \subset \mathbb{R}$  which is bounded from below.

Supremum and infimum are generalizations of the notions of maximum (i.e., the largest element) and minimum (i.e., the smallest element). For comparison, let us recall also these notions.

**Definition II.30** (Maximum and minimum).

Let  $A \subset \mathbb{R}$  be a subset.

An element  $a' \in A$  is the **maximum** (i.e. the **largest element**) of  $A$ , if for all  $a \in A$  we have  $a \leq a'$ . We then denote  $a' = \max A$ .

An element  $a'' \in A$  is the **minimum** (i.e. the **smallest element**) of  $A$ , if for all  $a \in A$  we have  $a \geq a''$ . We then denote  $a'' = \min A$ .

The crucial difference between the maximum and the supremum (or an upper bound more generally) is that the maximum of  $A$  is required to be an element of the set  $A$

<sup>17</sup>This condition simply says that  $u_0$  is an upper bound for  $A$ .

<sup>18</sup>This condition says that any smaller number  $u_0 - \varepsilon$  cannot be an upper bound for  $A$ .

itself. A similar remark applies to the difference between the minimum and the infimum.

Note, however, that not every subset  $A \subset \mathbb{R}$  has a maximum (resp. minimum) — even if we assume  $A$  to be non-empty and bounded from above (resp. from below).

**Example II.31** (Open intervals have no maximum and minimum).

The open interval  $(-\pi, \pi)$  has no largest element and no smallest element ( $\nexists \max(-\pi, \pi)$ ,  $\nexists \min(-\pi, \pi)$ ). In fact, open intervals never have a maximum or minimum.

**Example II.32** (A bounded nonempty set).

The set  $A = \{\frac{1}{n} \mid n \in \mathbb{N}\}$  is bounded and non-empty. The largest element of this set is  $\max A = 1$ . The set  $A$  has no smallest element ( $\nexists \min A$ ). The least upper bound of this set is  $\sup A = 1$  and the greatest lower bound of it is  $\inf A = 0$ . Note that in this case the infimum is not an element of the set,  $0 \notin A$ .

**Exercise II.15** (Examine properties of some sets).

Consider the sets

$$A_1 = [0, \sqrt{2}] \cap \mathbb{Q}, \quad A_2 = \{e^{-x} \mid x \in \mathbb{R}\}, \quad A_3 = \left\{ \frac{7n(-1)^n + 4}{6n} \mid n \in \mathbb{N} \right\}.$$

Answer the following questions about each of the sets  $A_j$ ,  $j \in \{1, 2, 3\}$ .

- Is the set  $A_j$  bounded from above?
- Is it bounded from below?
- Does it have a maximum, and if it does, what is  $\max A_j$ ?
- Does it have a supremum, and if it does, what is  $\sup A_j$ ?
- Does it have a minimum, and if it does, what is  $\min A_j$ ?
- Does it have an infimum, and if it does, what is  $\inf A_j$ ?

**Exercise II.16** (The relationship between supremum and maximum).

Let  $A \subset \mathbb{R}$ .

- (a) Prove that if  $\max A$  exists, then we have  $\sup A = \max A$ .
- (b) Prove that if  $\sup A \in A$  (so in particular the supremum exists), then the maximum exists and we have  $\max A = \sup A$ .

**Exercise II.17** (The relationship between infimum and minimum).

Formulate and prove an analogous result for the infimum and minimum.

### *Formulations of the completeness axiom*

There are various equivalent ways of formulating the completeness axiom of  $\mathbb{R}$ . We first state three formulations, and later discuss their equivalence.

The **completeness axiom** of  $\mathbb{R}$  refers to any of the following (equivalent) statements:

- (C1): Every non-empty subset  $A \subset \mathbb{R}$  which is bounded from above has a least upper bound  $\sup A \in \mathbb{R}$ .
- (C2): Every increasing real number sequence  $(a_n)_{n \in \mathbb{N}}$  which is bounded from above has a limit  $\lim_{n \rightarrow \infty} a_n \in \mathbb{R}$ .
- (C3): Every collection  $(I_n)_{n \in \mathbb{N}}$  of closed intervals  $I_n \subset \mathbb{R}$ , which is nested in the sense that  $I_{n+1} \subset I_n$  for every  $n \in \mathbb{N}$ , has a nonempty intersection

$$\bigcap_{n \in \mathbb{N}} I_n \neq \emptyset.$$



Regarding the last formulation (C3), note that in Example I.8 we saw that the intersection of nested *open intervals* can be empty. It is therefore important that in (C3) the intervals are required to be *closed*.

Furthermore, it is easy to see that the first two formulations above are equivalent with the following two, respectively:

**(C1'):** Every non-empty subset  $A \subset \mathbb{R}$  which is bounded from below has a greatest lower bound  $\inf A \in \mathbb{R}$ .

**(C2'):** Every decreasing real number sequence  $(a_n)_{n \in \mathbb{N}}$  which is bounded from below has a limit  $\lim_{n \rightarrow \infty} a_n \in \mathbb{R}$ .

**Exercise II.18** (Equivalence of (C1) and (C1')).

Prove that (C1) and (C1') are logically equivalent.

**Exercise II.19** (Equivalence of (C2) and (C2')).

Prove that (C2) and (C2') are logically equivalent.

The equivalence of the three formulations of the completeness axiom is proven in Appendix B.2.

As an example application of the formulation (C2) of the completeness axiom, we discuss the existence of real numbers given by decimal expansions.

**Example II.33** (Decimal expansions).

Let  $(d_1, d_2, d_3, \dots) = (d_k)_{k \in \mathbb{N}}$  be a sequence of digits  $d_k \in \{0, 1, 2, \dots, 8, 9\}$ . The precise definition of the **decimal number**

$$0.d_1d_2d_3\dots$$

is the sum of the series

$$\sum_{k=1}^{\infty} d_k 10^{-k}. \quad (\text{II.12})$$

In particular, the existence of this decimal number requires the series to be convergent.

The partial sums of the series

$$a_n = \sum_{k=1}^n d_k 10^{-k}$$

clearly form an increasing sequence  $(a_n)_{n \in \mathbb{N}}$ , because

$$a_{n+1} - a_n = \sum_{k=1}^{n+1} d_k 10^{-k} - \sum_{k=1}^n d_k 10^{-k} = d_{n+1} 10^{-n-1} \geq 0.$$

The sequence of the partial sums is also bounded from above, because

$$\begin{aligned}
 a_n &= \sum_{k=1}^n d_k 10^{-k} \\
 &\leq \sum_{k=1}^n 9 \cdot 10^{-k} && (\text{since } d_k \leq 9) \\
 &= \frac{9}{10} \sum_{\ell=0}^{n-1} 10^{-\ell} && (\text{change summation index to } \ell = k - 1) \\
 &= \frac{9}{10} \frac{1 - 10^{-n}}{1 - \frac{1}{10}} && (\text{finite geometric sum}) \\
 &\leq \frac{9}{10} \frac{1}{9/10} = 1.
 \end{aligned}$$

According to the formulation (C2) of the completeness axiom, the sequence  $(a_n)_{n \in \mathbb{N}}$  of partial sums converges, because it is increasing and bounded from above. This means that the series (II.12) converges.

We conclude that every decimal expansion indeed represents some real number.

### *Conventions to extend the notion of supremum and infimum*

In Definition II.28 we considered the supremum only for subsets  $A$  which are nonempty and bounded from above. If  $A$  fails one of these properties, we will use the following conventions:

- If a set  $A \subset \mathbb{R}$  is not bounded from above (and thus has no upper bounds), we interpret the least upper bound as the symbol  $+\infty$  (not a number),

$$\sup A = +\infty.$$

- For the empty set  $\emptyset \subset \mathbb{R}$  (for which any real number is an upper bound), we interpret the least upper bound as the symbol  $-\infty$  (not a number),

$$\sup \emptyset = -\infty.$$

Symmetrically for the infimum we use the conventions:

- If a set  $A \subset \mathbb{R}$  is not bounded from below (and thus has no lower bounds), we interpret the greatest lower bound as the symbol  $-\infty$  (not a number),

$$\inf A = -\infty.$$

- For the empty set  $\emptyset \subset \mathbb{R}$  (for which any real number is a lower bound), we interpret the greatest lower bound as the symbol  $+\infty$  (not a number),

$$\inf \emptyset = +\infty.$$

## **II.6. ♥ In depth topics about the real numbers**

The reader interested in deepening her understanding may wish to look into some of the following topics related to the subject of this lecture:

- The more familiar axioms of the real numbers, Appendix B.1.
- The proof of equivalence of the different formulations of the completeness axiom, Appendix B.2.

- The Cantor set — a fractal subset of the unit interval with interesting set theoretic, topological, and measure theoretic properties, Appendix B.3.
- Applications of supremum and infimum, Appendix B.4.



## Lecture III

### Sequences and functions on the real line

The main theme of this section is continuous functions of a real variable — certainly to some extent a familiar topic already. We will in particular verify that polynomials and rational functions are continuous. We will examine continuous functions on closed intervals, and show that they always have a maximum and a minimum — a crucial existence result for optimization tasks<sup>1</sup>, for example — and especially that they are bounded. We also look into the intermediate value theorem (Bolzano's theorem), according to which continuous functions on intervals "can not skip values".

One of the main objectives is to precisely define the notion of continuity, and to rigorously prove the above (probably familiar) facts. Another goal is to start drawing attention to the underlying topological reasons behind such important results: later in the course it is possible to appreciate the role of compactness (of closed intervals) and connectedness (of intervals) in the results.

As a tool we will still use real number sequences, and in particular their judiciously chosen subsequences. A key role is played by the fundamental fact that from a bounded sequence it is always possible to pick some convergent subsequence.<sup>2</sup>

#### III.1. Real number sequences

##### Subsequences

Consider a sequence of real numbers,

$$(x_1, x_2, x_3, x_4, x_5, \dots).$$

The idea of a subsequence is that we leave out some members from the original sequence. For example if we leave out the first member of the above sequence, we obtain the (sub)sequence

$$(x_2, x_3, x_4, x_5, x_6, \dots),$$

and if we leave out all members of even indices, we obtain the (sub)sequence

$$(x_1, x_3, x_5, x_7, x_9, \dots).$$

We anyway require that infinitely many members remain after some leaving out some. Also the convention is that the remaining members are relabeled by the

---

<sup>1</sup>Mathematically, optimization is formulated as the task of finding the maximum (or minimum) of some objective function: e.g., "profit", "accuracy", "efficiency", ... (or "cost", "error", "risk", ...). If the maximum (or minimum) does not exist, the problem is not even well posed and it is meaningless to look for the optimum!

<sup>2</sup>This fact that makes use of the completeness axiom of the real number field, and will itself in later chapters be recognized as a compactness statement.

original index set  $\mathbb{N}$  again<sup>3</sup> — this ensures that the subsequence itself will in fact be a sequence (in the sense of Section II.2).

The precise definition of a subsequence is the following.

**Definition III.1** (Subsequence).

Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence, and  $\varphi: \mathbb{N} \rightarrow \mathbb{N}$  a strictly increasing function. Then the sequence

$$(x_{\varphi(n)})_{n \in \mathbb{N}} = (x_{\varphi(1)}, x_{\varphi(2)}, x_{\varphi(3)}, \dots)$$

is called a **subsequence** of the sequence  $(x_n)_{n \in \mathbb{N}}$ .

Another common notation for a subsequence is

$$(x_{n_k})_{k \in \mathbb{N}} = (x_{n_1}, x_{n_2}, x_{n_3}, \dots),$$

where  $k$  is used for the new indices, and from the original sequence we have picked members with indices  $n_1 < n_2 < n_3 < \dots$ , corresponding to  $n_k = \varphi(k)$  in the above definition.

It is easy to show (by induction) that for a strictly increasing function  $\varphi: \mathbb{N} \rightarrow \mathbb{N}$  we have

$$\varphi(n) \geq n \quad \text{for all } n \in \mathbb{N},$$

which implies that a member in a subsequence corresponds to a member in the original sequence which is at least as far in the tail of the original sequence (its index in the original sequence is at least as large as its index in the subsequence). This observation makes the proof of the following lemma very easy.

**Lemma III.2** (All subsequences of a convergent sequence have the same limit).

*Suppose that  $(x_n)_{n \in \mathbb{N}}$  is a real number sequence which converges to a limit  $\lim_{n \rightarrow \infty} x_n = x$ . Then any subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  also converges to the same limit,  $\lim_{n \rightarrow \infty} x_{\varphi(n)} = x$ .*

**Exercise III.1** (Proof of Lemma III.2).

Prove Lemma III.2.

**Example III.3** (The alternating sign sequence).

Consider the sequence  $(a_n)_{n \in \mathbb{N}}$  where  $a_n = (-1)^n$ , i.e.,

$$(a_n)_{n \in \mathbb{N}} = (-1, +1, -1, +1, -1, \dots).$$

The subsequence of even members, obtained with  $\varphi_{\text{even}}(n) = 2n$ , is

$$(a_{2n})_{n \in \mathbb{N}} = (+1, +1, +1, +1, \dots),$$

and it obviously converges to  $+1$ .

The subsequence of odd members, obtained with  $\varphi_{\text{odd}}(n) = 2n - 1$ , is

$$(a_{2n-1})_{n \in \mathbb{N}} = (-1, -1, -1, -1, \dots),$$

and it obviously converges to  $-1$ .

The fact that these two subsequences have different limits implies (by Lemma III.2) that the original sequence  $(a_n)_{n \in \mathbb{N}}$  does not converge. (Of course this could also be proven directly!)

---

<sup>3</sup>It is possible to consider index sets other than  $\mathbb{N}$ , but even in such cases we require that the indices are consecutive integers. For simplicity, let us focus on the case when the index set is  $\mathbb{N}$ .

**Example III.4** (A sequence without convergent subsequences).

Consider the sequence  $(a_n)_{n \in \mathbb{N}}$  where  $a_n = -n$ , i.e.,

$$(a_n)_{n \in \mathbb{N}} = (-1, -2, -3, -4, \dots).$$

It is easy to see that this sequence has no convergent subsequences (in fact it has no bounded subsequences, so this follows from Proposition II.13).

### *Existence of monotone subsequences*

It turns out that from any real number sequence, it is possible to pick a subsequence which is monotone (either increasing or decreasing). This observation is somewhat interesting on its own right, but its real value is as a lemma towards the next very important result, Theorem III.6.

**Lemma III.5** (Any real number sequence has a monotone subsequence).

*Let  $(x_n)_{n \in \mathbb{N}}$  be a real number sequence. Then there exists a subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  of  $(x_n)_{n \in \mathbb{N}}$  which is monotone.*

*Proof.* Define the set

$$J := \left\{ n \in \mathbb{N} \mid x_m \leq x_n \text{ for all } m > n \right\}$$

of those indices, whose corresponding value is never exceeded later in the sequence. We consider two cases: the set  $J$  is either finite or infinite.

Suppose first that the set  $J$  is infinite. In this case, by enumerating the elements of  $J$  in an increasing order

$$J = \{n_1, n_2, n_3, \dots\} \quad \text{with} \quad n_1 < n_2 < n_3 < \dots,$$

we obtain indices such that the corresponding subsequence

$$(x_{n_1}, x_{n_2}, x_{n_3}, \dots)$$

is decreasing: indeed we have  $x_{n_{k+1}} \leq x_{n_k}$  for all  $k \in \mathbb{N}$  because  $n_k \in J$  and  $n_{k+1} > n_k$ . In this case we have thus found a monotone (decreasing) subsequence.

Suppose then that the set  $J$  is finite. In this case there exists<sup>4</sup> some  $n_1 \in \mathbb{N}$  such that all of the elements of  $J$  are smaller than  $n_1$ . By the definition of the set  $J$ , this implies that starting from the index  $n_1$ , every member of the sequence is later exceeded; or more formally

$$\forall n \geq n_1 : \exists m > n : x_m > x_n.$$

We can use this property to recursively find members which form a good subsequence. When the first  $k$  indices  $n_1 < \dots < n_k$  have been chosen, the property above implies that there exists some  $n_{k+1} > n_k$  such that  $x_{n_{k+1}} > x_{n_k}$ , which we will use as the  $k+1$ :st index. The resulting subsequence  $(x_{n_1}, x_{n_2}, x_{n_3}, \dots)$  will be strictly increasing by construction:  $x_{n_1} < x_{n_2} < x_{n_3} < \dots$ . In this case, too, we have found a monotone (increasing) subsequence. The proof is thus complete.  $\square$

### *Bounded sequences have convergent subsequences*

The following result is crucially important in real analysis. We will see some of its familiar but powerful consequences already in this lecture.

**Theorem III.6** (Bounded real number sequences have convergent subsequences).

*Let  $(x_n)_{n \in \mathbb{N}}$  be a real number sequence which is bounded, i.e., for some  $r \in \mathbb{R}$  we have  $|x_n| \leq r$  for all  $n \in \mathbb{N}$ . Then there exists a subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  of  $(x_n)_{n \in \mathbb{N}}$  which is convergent.*

<sup>4</sup>If  $J \neq \emptyset$ , then  $n_1 = \max J + 1$  works, and if  $J = \emptyset$  then  $n_1 = 1$  works.

*Proof.* By Lemma III.5, the sequence  $(x_n)_{n \in \mathbb{N}}$  has a monotone subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$ . But this subsequence is clearly also bounded (by the same constant  $r$  as the original sequence). As a bounded monotone sequence of real numbers,  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  converges by the completeness axiom — formulation (C2) or (C2'). We conclude that  $(x_n)_{n \in \mathbb{N}}$  has a convergent subsequence.  $\square$

### III.2. Functions of real variable

We will now consider real-valued functions of a real variable. Being a function of a real variable means that we take the domain of the function to be a subset  $A \subset \mathbb{R}$  of the real axis. Real-valuedness means that we (can) take the codomain of the functions to be  $\mathbb{R}$ . Therefore we are interested in functions

$$f: A \rightarrow \mathbb{R} \quad \text{where } A \subset \mathbb{R}.$$

Important special cases of the choice of the domain include, e.g., closed intervals  $A = [a, b] \subset \mathbb{R}$  and the whole real axis  $A = \mathbb{R}$ .

#### Continuity of a function of real variable

The intuitive idea of continuous functions should be familiar from calculus courses. As a precise definition, we take the following.<sup>5</sup>

**Definition III.7** (Continuity of a real-valued function of a real variable).

Let  $A \subset \mathbb{R}$ , and let  $f: A \rightarrow \mathbb{R}$  be a function. We say that  $f$  is **continuous** if the following holds: whenever  $(a_n)_{n \in \mathbb{N}}$  is a sequence in  $A$  which is convergent and its limit is in the domain,  $\lim_{n \rightarrow \infty} a_n \in A$ , then we have

$$\lim_{n \rightarrow \infty} f(a_n) = f\left(\lim_{n \rightarrow \infty} a_n\right).$$

To elaborate — given a sequence  $(a_n)_{n \in \mathbb{N}}$  in the domain  $A$  of the function  $f: A \rightarrow \mathbb{R}$ , we map the members  $a_n \in A$  of the sequence to real numbers  $f(a_n) \in \mathbb{R}$ , and form a real number sequence  $(f(a_n))_{n \in \mathbb{N}}$  of these images under  $f$ , i.e., of the corresponding function values. Continuity of  $f$  (Definition III.7) is the requirement that if the original sequence is convergent and its limit is in the domain  $A$ , then also the image sequence is convergent and its limit is the image of the limit of the original sequence. In somewhat imprecise terms, “we are allowed to interchange the order of (i) taking the limit and (ii) applying the function”.<sup>6</sup>

Certainly a function which abruptly jumps from one value to another should be the simplest example of a function that is not continuous; see Figure III.1. In the next exercise you check the discontinuity of such a step function directly from the definition.

<sup>5</sup>Later, in Chapter VI, we define continuity in a more general context. The general definition we take will be shown to be equivalent to the definition used here, in the special case of real-valued functions of a real variable.

<sup>6</sup>Note, however, that we still have to assume convergence of the original sequence to a limit in the domain.





FIGURE III.1. The step function of Exercise III.2 is a basic example of a function that is not continuous.

**Exercise III.2** (Step function is discontinuous).

Consider the function

$$f: \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) := \begin{cases} -1 & \text{for } x < 0 \\ +1 & \text{for } x \geq 0. \end{cases}$$

Show that it is not a continuous function  $\mathbb{R} \rightarrow \mathbb{R}$ .

*Hint:* By Definition III.7 it suffices to find some convergent sequence  $(x_n)_{n \in \mathbb{N}}$  on  $\mathbb{R}$  such that  $\lim_n f(x_n) \neq f(\lim_n x_n)$ .

**Example III.8** (Another discontinuous function).

Consider the function

$$f: [0, +\infty) \rightarrow \mathbb{R}, \quad f(x) := \begin{cases} -1 & \text{for } x = 0 \\ \cos(\pi + 2\pi/x) & \text{for } x > 0. \end{cases}$$

Is this function continuous or not? (For a plot, see Figure III.2.)

Intuitively it seems that any potential problem should occur at  $x = 0$ , if at all. So let us try out some sequences  $(x_n)_{n \in \mathbb{N}}$  which tend to 0.

First try the sequence with members  $x_n = \frac{1}{n}$ . This sequence has limit

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

The values of  $f$  at members of the sequence are

$$f(x_n) = \cos\left(\pi + \frac{2\pi}{1/n}\right) = \cos(\pi + 2\pi n) = -1 \quad \text{for } n \in \mathbb{N}.$$

Therefore we find

$$\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} (-1) = -1 = f(0) = f\left(\lim_{n \rightarrow \infty} x_n\right).$$

This is as required in the definition of continuity. (Note that  $x_n = \frac{1}{n} \in [0, +\infty)$  for all  $n \in \mathbb{N}$ , so this is a sequence in the domain of definition of  $f$ , and also the limit  $\lim_n x_n = 0$  remains in the domain,  $0 \in [0, +\infty)$ .)

If you were to try for example the sequence with members  $x_n = 2^{-n}$ , which also tends to zero, you would again find  $\lim f(x_n) = f(\lim x_n)$ .

But Definition III.7 requires a similar conclusion for *all* sequences (which are convergent in the domain of  $f$ ). So a single counterexample will be sufficient to show discontinuity! For




FIGURE III.2. The function of Example III.8.

such a counterexample, take the sequence with members  $x_n = \frac{4}{4n-1}$ . One again notes that  $x_n \in [0, +\infty)$  for all  $n \in \mathbb{N}$ , and  $\lim x_n = 0$ . Now the function values along this sequence are

$$f(x_n) = \cos\left(\pi + \frac{2\pi}{4/(4n-1)}\right) = \cos\left(\pi + \frac{\pi}{2}(4n-1)\right) = \cos\left(\frac{\pi}{2} + 2\pi n\right) = 0$$

for  $n \in \mathbb{N}$ . Therefore

$$\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0 \neq -1 = f(0) = f\left(\lim_{n \rightarrow \infty} x_n\right).$$

This shows that  $f$  is not continuous.

**Exercise III.3** (Indicator function of rational numbers).

Consider the function

$$f: \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) := \begin{cases} 1 & \text{for } x \in \mathbb{Q} \\ 0 & \text{for } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

Is it continuous?

Note that continuity depends on the domain of definition of the function: a discontinuous function restricted to a smaller set can become continuous, as you will see in the next example.

**Exercise III.4** (The restriction of a discontinuous function can be continuous).

Consider the step function  $f: \mathbb{R} \rightarrow \mathbb{R}$  of Exercise III.2, and let  $\tilde{f} = f|_{\mathbb{R} \setminus \{0\}}$  be its restriction to the set  $\mathbb{R} \setminus \{0\}$  non-zero real numbers. Show that  $\tilde{f}: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$  is continuous.

In the converse direction things are stable: a function never loses its continuity when restricting the domain.

**Exercise III.5** (Restrictions of continuous functions are continuous).

Suppose  $f: A \rightarrow \mathbb{R}$  is a continuous function and  $\tilde{A} \subset A$  is a subset. Show that the restriction  $\tilde{f} = f|_{\tilde{A}}$  is a continuous function  $\tilde{f}: \tilde{A} \rightarrow \mathbb{R}$ .

## Operations on continuous functions

Because the definition of continuity relies on limits of real number sequences, our earlier results about limits easily yield some fundamental results about continuous functions. Particularly important is the fact that the following pointwise operations preserve continuity.

**Definition III.9** (Pointwise operations on real-valued functions of real variable).

Let  $A \subset \mathbb{R}$ , and let  $f, g: A \rightarrow \mathbb{R}$  be two real-valued functions on  $A$ .

The **pointwise sum** of  $f$  and  $g$  is the function

$$f + g: A \rightarrow \mathbb{R} \quad (f + g)(x) := f(x) + g(x) \quad \text{for } x \in A.$$

The **pointwise product** of  $f$  and  $g$  is the function

$$fg: A \rightarrow \mathbb{R} \quad (fg)(x) := f(x)g(x) \quad \text{for } x \in A.$$

The **pointwise quotient** of  $f$  and  $g$  is the function

$$f/g: A' \rightarrow \mathbb{R} \quad (f/g)(x) := \frac{f(x)}{g(x)} \quad \text{for } x \in A',$$

where  $A' := \{x \in A \mid g(x) \neq 0\}$ .

Note that the pointwise quotient  $f/g$  is only defined on the subset  $A' \subset A$  of the domain, where the function  $g$  is non-vanishing — in order to avoid ill-defined division by zero.

Scalar multiplication of a function  $f: A \rightarrow \mathbb{R}$  is a special case of the pointwise product: it is natural to interpret a real number  $c \in \mathbb{R}$  also as the constant function  $x \mapsto c$  on  $A$ , so  $cf: A \rightarrow \mathbb{R}$  is defined by  $(cf)(x) := cf(x)$  for  $x \in A$ .

**Proposition III.10** (Continuity preserving operations).

Let  $A \subset \mathbb{R}$ , and let  $f, g: A \rightarrow \mathbb{R}$  be two continuous real-valued functions on  $A$ . Then also

- (i) the pointwise sum function  $f + g: A \rightarrow \mathbb{R}$  is continuous,
- (ii) the pointwise product function  $fg: A \rightarrow \mathbb{R}$  is continuous,
- (iii) the pointwise quotient function  $f/g: A' \rightarrow \mathbb{R}$  is continuous on the subset  $A' = \{x \in A \mid g(x) \neq 0\}$ .

*Proof.* The proofs of all cases are essentially similar, so let us provide the details only for (ii), and leave (i) and (iii) as exercises.

*proof of (ii):* Assume  $f, g: A \rightarrow \mathbb{R}$  are continuous. We must prove that  $fg: A \rightarrow \mathbb{R}$  is continuous, and we will do this directly using Definition III.7 of continuity. So suppose  $(x_n)_{n \in \mathbb{N}}$  is a sequence such that  $x_n \in A$  for all  $n \in \mathbb{N}$ , and  $(x_n)_{n \in \mathbb{N}}$  converges to a limit in  $A$ . Let us denote the limit by  $x = \lim_{n \rightarrow \infty} x_n \in A$ . By definition of continuity, we must consider the sequence  $((fg)(x_n))_{n \in \mathbb{N}}$  of values of the pointwise product function  $fg$  and show that it converges to the limit  $(fg)(x)$ . By definition of pointwise products, the values are  $(fg)(x_n) = f(x_n)g(x_n)$ . By assumptions of continuity of  $f$  and  $g$  we know that  $\lim_{n \rightarrow \infty} f(x_n) = f(x)$  and  $\lim_{n \rightarrow \infty} g(x_n) = g(x)$ , so by (II.9) of Theorem II.19 we have

$$\lim_{n \rightarrow \infty} (f(x_n)g(x_n)) = \left( \lim_{n \rightarrow \infty} f(x_n) \right) \left( \lim_{n \rightarrow \infty} g(x_n) \right) = f(x)g(x).$$

By definition of the pointwise products the right hand side above is  $(fg)(x)$ , so we have shown

$$\lim_{n \rightarrow \infty} ((fg)(x_n)) = (fg)(x),$$

which is what was needed to prove continuity.  $\square$

**Exercise III.6** (Proof of Proposition III.10 (i) and (iii)).

Prove parts (i) and (iii) of the above proposition.

**Exercise III.7** (Continuity of the tangent trigonometric function).

Is the function

$$\tan: \mathbb{R} \setminus \left\{ \frac{\pi}{2} + n\pi \mid n \in \mathbb{Z} \right\} \rightarrow \mathbb{R}$$

continuous?

Hint: The continuity of the trigonometric functions  $\sin: \mathbb{R} \rightarrow \mathbb{R}$  and  $\cos: \mathbb{R} \rightarrow \mathbb{R}$  can be considered known here (a precise justification will be given in Chapter IX). Recall that  $\tan(x) = \frac{\sin(x)}{\cos(x)}$ .

**Corollary III.11** (Polynomials and rational functions are continuous).

Let  $a_0, a_1, a_2, \dots, a_n \in \mathbb{R}$  and consider the polynomial function

$$P(x) := a_0 + a_1x + a_2x^2 + \dots + a_nx^n \quad \text{for } x \in \mathbb{R}.$$

Then  $P: \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function. Let  $a_0, a_1, a_2, \dots, a_n \in \mathbb{R}$  and consider the polynomial function

Let also  $b_0, b_1, b_2, \dots, b_m \in \mathbb{R}$  and

$$Q(x) := b_0 + b_1x + b_2x^2 + \dots + b_mx^m \quad \text{for } x \in \mathbb{R}.$$

Consider the rational function

$$R(x) = \frac{P(x)}{Q(x)} \quad \text{for } x \in A,$$

where  $A = \{x \in \mathbb{R} \mid Q(x) \neq 0\}$ . Then  $R: A \rightarrow \mathbb{R}$  is a continuous function.

*Proof.* Let us first observe that the continuity of the identity function  $x \mapsto x$  (as a function  $\mathbb{R} \rightarrow \mathbb{R}$ ) is trivial from the definition. The function  $x \mapsto x^2$  is a pointwise product of identity functions, and therefore also continuous by Proposition III.10(ii). Continuing to take pointwise products with the identity function, we find that the monomial functions  $x \mapsto x^k$  are continuous for all  $k \in \mathbb{N}$  (easy induction).

Then note that for any  $c \in \mathbb{R}$  the continuity of the constant function  $x \mapsto c$  (as a function  $\mathbb{R} \rightarrow \mathbb{R}$ ) is also obvious from the definition. Taking  $c = a_k \in \mathbb{R}$  and further taking pointwise product with a monomial function, we find that  $x \mapsto a_kx^k$  is continuous, for any  $k \in \mathbb{Z}_{\geq 0}$  (for the case  $k = 0$  one does not even need a product with monomial function). The polynomial  $P(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$  is a finite sum of terms of this type, so continuous by Proposition III.10(i), inductively applied.

Since polynomial functions given by  $P(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$  and  $Q(x) = b_0 + b_1x + b_2x^2 + \dots + b_mx^m$  are continuous, we get from Proposition III.10(iii) that the rational function given by the pointwise quotient  $R(x) = \frac{P(x)}{Q(x)}$  is continuous on the subset  $A = \{x \in \mathbb{R} \mid Q(x) \neq 0\} \subset \mathbb{R}$  where the denominator is non-vanishing.  $\square$


Later in the course we will develop more powerful tools for verifying the continuity of functions. But knowing the continuity of polynomial and rational functions at least assures us that some nontrivial continuous functions exist; see Figure III.3 for a few concrete examples.



(a) The polynomial  $x \mapsto x^3 - 2x$  defines a function  $\mathbb{R} \rightarrow \mathbb{R}$ .



(b) The function  $x \mapsto \frac{1}{1+5x^2}$  is a rational function  $\mathbb{R} \rightarrow \mathbb{R}$ .



(c) The rational function  $x \mapsto \frac{1}{x(x^2-1)^2}$  has poles at  $x = 0$  and  $x = \pm 1$ , and defines a (continuous) function  $\mathbb{R} \setminus \{-1, 0, +1\} \rightarrow \mathbb{R}$ .

FIGURE III.3. Polynomials and rational functions are continuous.

### III.3. Continuous functions on a closed interval

Let us now address specifically the case when the domain of the function is closed interval on the real axis,  $A = [a, b] \subset \mathbb{R}$ . This case appears very frequently in applications. We address some key properties of continuous functions

$$f: [a, b] \rightarrow \mathbb{R}$$

on a closed interval; later in the course we will understand them from more general perspectives.

Let us recall a few key concepts: a function  $f: A \rightarrow \mathbb{R}$  is **bounded** if there exists a  $M \geq 0$  such that  $|f(x)| \leq M$  for all  $x \in A$ . A function  $f: A \rightarrow \mathbb{R}$  has a **maximum** if there exists a point  $x_{\max} \in A$  such that  $f(x) \leq f(x_{\max})$  for all  $x \in A$ , and it has a **minimum** if there exists a point  $x_{\min} \in A$  such that  $f(x) \geq f(x_{\min})$  for all  $x \in A$ .

**Remark III.12** (Reinterpretation in terms of the range of the function).

Consider the image  $f[A] = \{f(x) \mid x \in A\} \subset \mathbb{R}$  of the whole domain  $A$  (this is often called the range of  $f$ ). It is straightforward to verify that the function  $f$  is bounded if and only if the set  $f[A] \subset \mathbb{R}$  is bounded, and the function  $f$  has a maximum (resp. minimum) if and only if the set  $f[A]$  has a maximum (resp. minimum).

The following important properties are related to compactness (see Chapter XI) of the closed interval.

**Theorem III.13** (Extrema of continuous functions on a closed interval).

*Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function. Then,*

- (i)  *$f$  is bounded,*
- (ii)  *$f$  has a maximum and a minimum.*


Before giving the proof, let us note that the conclusions of Theorem III.13 rely crucially on the assumption that the interval is closed. The examples in Figure III.4 illustrate what can go wrong on non-closed intervals.

**Exercise III.8** (Continuous functions on non-closed intervals).


- (a) Prove that the function of Figure III.4(a) is continuous but not bounded.
- (b) Prove that the function of Figure III.4(b) is continuous but has no minimum.
- (c) Find an example of a continuous function on an interval, which has no maximum.

*Proof of Theorem III.13* Let us prove the two assertions separately.

*proof of (i):* Suppose, by way of contradiction, that  $f$  is not bounded. Then for every  $n \in \mathbb{N}$ , it is possible to choose some  $x_n \in [a, b]$  such that  $|f(x_n)| \geq n$ . Let us form a sequence  $(x_n)_{n \in \mathbb{N}}$  using such choices. This sequence is bounded, since  $a \leq x_n \leq b$  for all  $n \in \mathbb{N}$ . Therefore, by Theorem III.6, it has a convergent subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$ . If we denote the limit of such a convergent subsequence by  $x = \lim_{n \rightarrow \infty} x_{\varphi(n)}$ , then by the preservation of bounds we have  $a \leq x \leq b$  (see Corollary II.15). By continuity, then,  $(f(x_{\varphi(n)}))_{n \in \mathbb{N}}$  converges to  $f(x)$ . But by the choice of  $x_n$  we have  $|f(x_{\varphi(n)})| \geq \varphi(n) \geq n$  for all  $n \in \mathbb{N}$ , which means that the sequence  $(f(x_{\varphi(n)}))_{n \in \mathbb{N}}$  is not bounded and so cannot be convergent (see Proposition II.13). This is a contradiction. We conclude that  $f$  had to be bounded.



(a) The function  $f: (0, 1] \rightarrow \mathbb{R}$  given by  $f(x) = 1/x$  is continuous (a rational function) but not bounded.



(b) The function  $f: (1, 3] \rightarrow \mathbb{R}$  given by  $f(x) = x$  is continuous (a polynomial function) but has no minimum.

FIGURE III.4. Continuous functions on general intervals do not need to be bounded and do not need to have minima and maxima.

*proof of (ii):* Let us only prove that  $f$  has a maximum — the existence of minimum can be concluded similarly (or by considering the maximum of the continuous function  $-f$ ).

By (i)  $f$  is bounded, so the supremum of its values is finite,

$$C := \sup \{f(x) \mid x \in [a, b]\} \in \mathbb{R}.$$

For all  $n \in \mathbb{N}$  there then exists some  $x_n \in [a, b]$  such that  $f(x_n) \geq C - \frac{1}{n}$ . Let us form a sequence  $(x_n)_{n \in \mathbb{N}}$  using such choices. As in part (i), this sequence is bounded and therefore has a convergent subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  whose limit  $x = \lim_{n \rightarrow \infty} x_{\varphi(n)}$  is also on the interval  $[a, b]$ . By the choice of  $C$  (supremum of the values) and of  $x_n$  we have

$$C \geq f(x_{\varphi(n)}) \geq C - \frac{1}{\varphi(n)} \geq C - \frac{1}{n}.$$

The squeeze theorem (Lemma II.17) thus gives  $\lim_{n \rightarrow \infty} f(x_{\varphi(n)}) = C$ . On the other hand, by continuity we have

$$f(x) = f\left(\lim_{n \rightarrow \infty} x_{\varphi(n)}\right) = \lim_{n \rightarrow \infty} f(x_{\varphi(n)}) = C.$$

We conclude that

$$f(x) = \sup \{f(x) \mid x \in [a, b]\},$$

which implies that the maximum of  $f: [a, b] \rightarrow \mathbb{R}$  is attained at  $x \in [a, b]$ .  $\square$

The following property is related to connectedness (see Chapter XII) of the closed interval.

**Theorem III.14** (Intermediate value theorem / Bolzano's theorem).

*Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function. Assume that  $f(a)f(b) < 0$ .<sup>7</sup> Then there exists a point  $z \in (a, b)$  such that  $f(z) = 0$ .*

*Proof.* We may assume that  $f(a) < 0$  and  $f(b) > 0$ ; the other case is similar (or is obtained from this one by considering the continuous function  $-f$  with the same zeroes as  $f$ ).

Consider the subset

$$N = \{x \in [a, b] \mid f(x) < 0\}$$

of the interval  $[a, b]$  where the function  $f$  is negative. Since  $f(a) < 0$ , we have at least  $a \in N$ , so this subset is non-empty,  $N \neq \emptyset$ . The subset is also bounded from above, because the right endpoint  $b$  of the interval is an upper bound for  $N$ . So let

$$z = \sup N.$$

The earlier observations yield  $z \geq a$  and  $z \leq b$ , so we have  $z \in [a, b]$ . By a characterization of the supremum (Lemma II.29), there exists a sequence  $(x_n^-)_{n \in \mathbb{N}}$  in the set  $N$  such that  $\lim_{n \rightarrow \infty} x_n^- = z$ . By continuity of  $f$  and by preservation of inequalities, we have

$$f(z) = f\left(\lim_{n \rightarrow \infty} x_n^-\right) = \lim_{n \rightarrow \infty} \underbrace{f(x_n^-)}_{<0} \leq 0.$$

In particular we have  $z \neq b$ , since  $f(b) > 0$ .

Then consider (for example) the sequence  $(x_n^+)_{n \in \mathbb{N}}$  with  $x_n^+ = z + \frac{1}{n}$ , which also has the property  $\lim_{n \rightarrow \infty} x_n^+ = z$  and  $f(x_n^+) \geq 0$  since  $x_n^+ \notin N$  (note that for large enough  $n$  we have  $x_n^+ = z + \frac{1}{n} \leq b$ , so the members of the sequence are in the domain  $[a, b]$  of  $f$ ). By continuity of  $f$  and by preservation of bounds, we have

$$f(z) = f\left(\lim_{n \rightarrow \infty} x_n^+\right) = \lim_{n \rightarrow \infty} \underbrace{f(x_n^+)}_{\geq 0} \geq 0.$$

In particular we have  $z \neq a$ , since  $f(a) < 0$ .

Therefore we have  $z \in (a, b)$ , and the above two inequalities combined yield  $f(z) = 0$ .  $\square$

**Corollary III.15** (Continuous images of intervals).

*If  $I \subset \mathbb{R}$  is an interval and  $f: I \rightarrow \mathbb{R}$  is continuous, then the image  $f[I]$  is also an interval.*

*Proof.* Let  $f: I \rightarrow \mathbb{R}$  be a continuous function defined on an interval  $I \subset \mathbb{R}$ . To show that the image  $f[I]$  is an interval, it suffices to show that for any two values  $\alpha, \beta \in f[I]$  in the image with  $\alpha < \beta$ , any value  $\gamma \in (\alpha, \beta)$  in between them also belongs to the image,  $\gamma \in f[I]$ .

So suppose  $\alpha, \beta \in f[I]$  with  $\alpha < \beta$ , and let  $\gamma \in (\alpha, \beta)$ . By definition of images, there exists some  $a, b \in I$  such that  $f(a) = \alpha$  and  $f(b) = \beta$ . We have either  $a < b$  or  $a > b$ . Both

<sup>7</sup>This is just a concise way of saying that  $f(a)$  and  $f(b)$  are non-zero and have opposite signs: either  $f(a) < 0$  and  $f(b) > 0$ , or  $f(a) > 0$  and  $f(b) < 0$ .



cases are handled similarly, so let us only present the details in the case  $a < b$ . Then by Exercise III.5 and Proposition III.10, the rule  $x \mapsto f(x) - \gamma$  defines a continuous function on the closed subinterval  $[a, b] \subset I$ . Moreover, its values at the endpoints of  $[a, b]$  have opposite signs:  $f(a) - \gamma = \alpha - \gamma < 0$  and  $f(b) - \gamma = \beta - \gamma > 0$ . By Theorem III.14, then, there exists a point  $z \in (a, b) \subset I$  such that  $f(z) - \gamma = 0$ , i.e.,  $f(z) = \gamma$ . This shows  $\gamma \in f[I]$ .  $\square$

**Example III.16** (The image of an interval under the sine function).

Consider the sine function

$$f_1: (0, \pi) \rightarrow \mathbb{R} \qquad f_1(x) = \sin(x) \quad \text{for } x \in (0, \pi).$$

This function is continuous.<sup>8</sup> The domain of definition is the open interval  $(0, \pi)$ , and its image under  $f_1$  is the half-open interval  $f_1[(0, \pi)] = (0, 1]$ .

If instead we considered the sine function on the longer interval

$$f_2: (0, 2\pi) \rightarrow \mathbb{R} \qquad f_2(x) = \sin(x) \quad \text{for } x \in (0, 2\pi),$$

then the image would be the closed interval  $f_2[(0, 2\pi)] = [-1, 1]$ .

The examples above show in particular that under a continuous function, the image of an open interval can be a closed or a half-open interval. Of course the image of an open interval can also be an open interval.<sup>9</sup>

However, by the next exercise the image of a closed interval under a continuous function is necessarily always a closed interval!<sup>10</sup>

**Exercise III.9** (The continuous image of a closed interval is a closed interval).

Using the combination of Corollary III.15 and Theorem III.13, prove that if  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f[[a, b]] \subset \mathbb{R}$  is a closed interval.

The following property is often used in non-degeneracy statements of formulas given by integrals; especially certain function space norms in the next chapter.

**Exercise III.10** (Integrals of non-zero nonnegative functions are positive).

Let  $[a, b]$  be a closed interval which is nondegenerate,  $a < b$ . Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function.

- Show that if there exists an  $x_0 \in [a, b]$  such that  $f(x_0) > 0$ , then there exists some  $\delta > 0$  such that  $f(x) > 0$  for all  $x \in (x_0 - \delta, x_0 + \delta) \cap [a, b]$ .  
*Hint: Argue by contradiction: if this property is not true, you can construct a sequence which violates the assumed continuity of  $f$ .*
- Improve the conclusion of part (a) as follows. Show that if there exists an  $x_0 \in [a, b]$  such that  $f(x_0) > 0$ , then there exists some  $c > 0$  and some  $\delta' > 0$  such that  $f(x) > c$  for all  $x \in (x_0 - \delta', x_0 + \delta') \cap [a, b]$ .  
*Hint: You can let  $c = \frac{1}{2}f(x_0)$ , and apply part (a) to the function defined by  $\tilde{f}(x) = f(x) - c$ .*
- Show that if  $f$  is non-negative, i.e.,  $f(x) \geq 0$  for all  $x \in [a, b]$ , and is not the constant function 0, then its integral is positive

$$\int_a^b f(x) \, dx > 0. \tag{III.1}$$

<sup>8</sup>The appropriate tools for the precise justification of the continuity of trigonometric functions are developed in Chapter IX; for now we just accept the continuity statement.

<sup>9</sup>Consider for example the identity function  $\text{id}_{(a,b)}: (a, b) \rightarrow (a, b)$ .

<sup>10</sup>The attentive reader may worry that constant functions are continuous and their images are singletons. This is indeed correct, but it does not contradict the assertion — a singleton is a special type of a closed interval, corresponding to the degenerate case when the interval endpoints coincide. Namely, when  $a = b$ , we have  $[a, b] := \{x \in \mathbb{R} \mid a \leq x \leq b\} = \{a\}$ .

Hint: Use part (b) and a lower bound for the integral based on what you know of the values of  $f$ .

## Lecture IV

### Normed spaces and inner product spaces

In this section we start to study more general spaces than just the familiar real axis  $\mathbb{R}$ . We will consider vector spaces equipped with some structure that in particular suffices to induce a topology. Specifically, we consider vector spaces equipped with an inner product or at least a norm.

The familiar  $d$ -dimensional Euclidean space  $\mathbb{R}^d$  has an inner product and a norm induced by it, but even this finite-dimensional vector space  $\mathbb{R}^d$  could be equipped with some slightly different norms — and for some applications it is meaningful to do so. There are also rather straightforward infinite-dimensional counterparts of these finite-dimensional normed spaces. Perhaps the most important applications of the structures introduced here are, however, various spaces of functions. We will already introduce a few function space examples that we keep returning to throughout the rest of this course, and that you will be very likely to encounter in whatever field of mathematics you end up using in the future. But the number of relevant examples of such function spaces is so vast that we are in fact only scratching the surface of a deep topic: in courses on Hilbert spaces, Banach spaces, Measure theory, Probability theory, Partial differential equations, etc. you will encounter many more. The basic structures introduced here will nevertheless be present in almost all of those!<sup>1</sup>

#### IV.1. Vector spaces

The topic of vector spaces by itself belongs to algebra or linear algebra. But because vector spaces with some additional structure feature so crucially in analysis as well, we review the notion here. However, for concreteness, we only define *real vector spaces*, i.e., vector spaces over the field  $\mathbb{R}$  of real numbers. Algebraists would typically study vector spaces over arbitrary fields.<sup>2</sup> The term **vector** will be used for elements of the vector space, and the term **scalar** for elements of the field over which the vector space is defined — in our case for real numbers.

---

<sup>1</sup>If one deliberately tries to find vector spaces in analysis that are *not* either inner product spaces or at least normed spaces, then among the easiest relevant examples are certain spaces of distributions. Even they have a topology, and while the topology does not come from a norm, it is not too far from that — in typical cases it comes from a collection of seminorms.

<sup>2</sup>Actually also *complex vector spaces*, i.e., vector spaces over the field  $\mathbb{C}$  of complex numbers, appear frequently in analysis, probability, etc. — and the most important additional structures they can be equipped with are again (complex) inner products and norms. Complex inner product spaces and complex normed spaces would require only relatively minor modifications to the treatment in this chapter. We choose to focus on just the real case for the sake of clarity. We trust that when you will need to work with the complex case, you will be well prepared for that as well.

### Axioms of real vector spaces

For brevity, the term vector space will from here on always mean a real vector space — defined precisely by the following axiomatic properties.

**Definition IV.1** (Vector space).

A (real) vector space is a set  $V$  equipped with two operations,

vector addition	scalar multiplication
$V \times V \rightarrow V$	$\mathbb{R} \times V \rightarrow V$
$(\vec{u}, \vec{v}) \mapsto \vec{u} + \vec{v}$	$(c, \vec{v}) \mapsto c\vec{v}$ ,

such that the following properties hold:

Commutativity of vector addition:

$$\forall \vec{u}, \vec{v} \in V : \quad \vec{u} + \vec{v} = \vec{v} + \vec{u} \quad (\text{IV.1})$$

Associativity of vector addition:

$$\forall \vec{u}, \vec{v}, \vec{w} \in V : \quad \vec{u} + (\vec{v} + \vec{w}) = (\vec{u} + \vec{v}) + \vec{w} \quad (\text{IV.2})$$

Neutral element for vector addition:

$$\exists \vec{0} \in V : \quad \forall \vec{v} \in V : \quad \vec{v} + \vec{0} = \vec{v} \quad (\text{IV.3})$$

Opposite vectors:

$$\forall \vec{v} \in V : \quad \exists (-\vec{v}) \in V : \quad \vec{v} + (-\vec{v}) = \vec{0} \quad (\text{IV.4})$$

Compatibility of scalar multiplication:

$$\forall c, d \in \mathbb{R} : \quad \forall \vec{v} \in V : \quad c(d\vec{v}) = (cd)\vec{v} \quad (\text{IV.5})$$

Unit scalar multiplication:

$$\forall \vec{v} \in V : \quad 1\vec{v} = \vec{v} \quad (\text{IV.6})$$

Distributivity of vector addition:

$$\forall c \in \mathbb{R} : \quad \forall \vec{u}, \vec{v} \in V : \quad c(\vec{u} + \vec{v}) = c\vec{u} + c\vec{v} \quad (\text{IV.7})$$

Distributivity of scalar addition:

$$\forall c, d \in \mathbb{R} : \quad \forall \vec{v} \in V : \quad (c + d)\vec{v} = c\vec{v} + d\vec{v}. \quad (\text{IV.8})$$

These axioms are probably every bit as unsurprising as the field axioms of the real numbers given in Appendix B.1. But again, from such a simple starting point, a very fruitful theory emerges.

In the above definition, we used arrows on top of vectors, e.g.  $\vec{u}, \vec{v}, \vec{w} \in V$ . In the long run, that becomes mainly an additional nuisance, so we will soon adopt the convention to label elements of the vector space without such decorations<sup>3</sup>, e.g. by  $u, v, w \in V$ .

### First examples of real vector spaces

The most obvious example of a real vector space is the following.

---

<sup>3</sup>Also the zero vector  $\vec{0}$  is commonly denoted by simply 0, and the reader is assumed to figure out from the context whether 0 stands for the scalar zero  $0 \in \mathbb{R}$  or the vector zero  $0 \in V$  (two very different objects).

**Example IV.2** (The standard  $d$ -dimensional space  $\mathbb{R}^d$ ).

The set

$$\mathbb{R}^d = \left\{ (x_1, \dots, x_d) \mid x_1, \dots, x_d \in \mathbb{R} \right\} \quad (\text{IV.9})$$

is a real vector space, when equipped with the coordinatewise vector addition

$$(x_1, \dots, x_d) + (y_1, \dots, y_d) = (x_1 + y_1, \dots, x_d + y_d) \\ \text{for } (x_1, \dots, x_d), (y_1, \dots, y_d) \in \mathbb{R}^d$$

and coordinatewise scalar multiplication

$$c(x_1, \dots, x_d) = (cx_1, \dots, cx_d) \\ \text{for } c \in \mathbb{R} \text{ and } (x_1, \dots, x_d) \in \mathbb{R}^d$$

The zero vector is  $\vec{0} = (0, 0, \dots, 0) \in \mathbb{R}^d$ , and the opposite vector to  $(x_1, \dots, x_d) \in \mathbb{R}^d$  is  $(-x_1, \dots, -x_d)$ . Verifying the properties (IV.1) – (IV.8) for this case is straightforward.

The previous example has a straightforward infinite-dimensional counterpart as well.

**Example IV.3** (The space of real number sequences).

The set<sup>4</sup>

$$\mathbb{R}^{\mathbb{N}} = \left\{ (x_1, x_2, x_3, \dots) \mid x_1, x_2, x_3, \dots \in \mathbb{R} \right\} \quad (\text{IV.10})$$

of all real number sequences is a real vector space, when equipped with the coordinatewise vector addition

$$(x_1, x_2, x_3, \dots) + (y_1, y_2, y_3, \dots) = (x_1 + y_1, x_2 + y_2, x_3 + y_3, \dots) \\ \text{for } (x_1, x_2, x_3, \dots), (y_1, y_2, y_3, \dots) \in \mathbb{R}^{\mathbb{N}}$$

and coordinatewise scalar multiplication

$$c(x_1, x_2, x_3, \dots) = (cx_1, cx_2, cx_3, \dots) \\ \text{for } c \in \mathbb{R} \text{ and } (x_1, x_2, x_3, \dots) \in \mathbb{R}^{\mathbb{N}}.$$

The zero vector is the sequence of zeroes  $\vec{0} = (0, 0, 0, \dots)$ , and the opposite vector to a sequence  $(x_1, x_2, x_3, \dots)$  is the sequence  $(-x_1, -x_2, -x_3, \dots)$ . Verifying that the properties (IV.1) – (IV.8) is again straightforward.

**Example IV.4** (Space of polynomials).

Consider the space

$$\mathbb{R}[x] = \left\{ \sum_{k=0}^d c_k x^k \mid d \in \mathbb{N}_0, c_0, c_1, \dots, c_d \in \mathbb{R} \right\} \quad (\text{IV.11})$$

of all polynomials in variable  $x$ , with real coefficients. The addition and scalar multiplication are defined coefficientwise, as usual for polynomials. It is easy to verify properties (IV.1) – (IV.8), so the space  $\mathbb{R}[x]$  of real polynomials in one variable is a real vector space.<sup>5</sup>

**Example IV.5** (Spaces of real valued functions).

Let  $X$  be any set. Consider the space

$$\mathbb{R}^X = \left\{ \text{functions } f: X \rightarrow \mathbb{R} \right\} \quad (\text{IV.12})$$

<sup>4</sup>If  $X$  and  $Y$  are sets, the notation  $Y^X$  is often used for the set of all functions  $X \rightarrow Y$ . The notation  $\mathbb{R}^{\mathbb{N}}$  correspondingly stands for all functions  $\mathbb{N} \rightarrow \mathbb{R}$  — which in view of the precise definition of sequences is exactly the set of all real number sequences.

<sup>5</sup>In fact, besides just addition and scalar multiplication of polynomials, we can also define multiplication of polynomials as usual. So the space  $\mathbb{R}[x]$  has even further algebraic structure than just that of a vector space over  $\mathbb{R}$  — it is what is known as an algebra over  $\mathbb{R}$ . We refer to courses on algebra for details.

of all functions whose domain is  $X$  and codomain is  $\mathbb{R}$ . We equip it with the *pointwise addition* and *pointwise scalar multiplication* as follows. If  $f, g \in \mathbb{R}^X$ , i.e., if  $f, g: X \rightarrow \mathbb{R}$  are two functions, then we define the new function  $f + g: X \rightarrow \mathbb{R}$  by the formula

$$(f + g)(x) = f(x) + g(x) \quad \text{for } x \in X. \quad (\text{IV.13})$$

Likewise, if  $c \in \mathbb{R}$  is a scalar and  $f \in \mathbb{R}^X$ , i.e.,  $f: X \rightarrow \mathbb{R}$  is a function, then we define the new function  $cf: X \rightarrow \mathbb{R}$  by the formula

$$(cf)(x) = cf(x) \quad \text{for } x \in X. \quad (\text{IV.14})$$

It is straightforward to check that the properties (IV.1) – (IV.8) hold, so the space  $\mathbb{R}^X$  of real valued functions on an arbitrary set  $X$  is a real vector space.<sup>6</sup>

In fact, Example IV.3 is readily a special case of the above, corresponding to the choice of domain  $X = \mathbb{N}$ : functions  $\mathbb{N} \rightarrow \mathbb{R}$  are exactly real number sequences.

Also Example IV.2 can be seen as a special case of this:  $\mathbb{R}^d$  can be interpreted as the set of real-valued functions on the finite set  $\{1, 2, \dots, d\}$ . A vector  $(x_1, \dots, x_d) \in \mathbb{R}^d$  corresponds to the function  $f: \{1, 2, \dots, d\} \rightarrow \mathbb{R}$  such that  $f(j) = x_j$  for  $j \in \{1, 2, \dots, d\}$ . This way we can identify the spaces  $\mathbb{R}^d$  and  $\mathbb{R}^{\{1, 2, \dots, d\}}$ .

## Vector subspaces

We saw a number of examples of vector spaces above, but other important examples are often constructed as vector subspaces of other vector spaces. The precise definition is the following.

### Definition IV.6 (Vector subspace).

Let  $V$  be a vector space. A subset  $W \subset V$  is called a **vector subspace** of  $V$ , if the following conditions hold:

- We have  $\vec{0} \in W$ ;
- If  $\vec{u}, \vec{v} \in W$ , then we also have  $\vec{u} + \vec{v} \in W$ .
- If  $c \in \mathbb{R}$  and  $\vec{v} \in W$ , then we also have  $c\vec{v} \in W$ .

A vector subspace is therefore stable under both vector addition and scalar multiplication, and in particular naturally inherits these operations. The properties in Definition IV.1 remain valid on a subspace  $W \subset V$ , so in particular  $W$  is itself a vector space. This is an often practical way of constructing vector spaces: instead of checking the eight properties in Definition IV.1, it suffices to verify the three (usually easy) properties of Definition IV.6, if the space has been realized as a subset of some known vector space.

### Example IV.7 (The subspace spanned by finitely many vectors).

If  $V$  is a vector space and  $\vec{v}_1, \dots, \vec{v}_n \in V$  are vectors in it, then the set

$$W = \left\{ c_1 \vec{v}_1 + \dots + c_n \vec{v}_n \mid c_1, \dots, c_n \in \mathbb{R} \right\} \quad (\text{IV.15})$$

of all linear combinations of them satisfies the properties in Definition IV.6, i.e. it is a vector subspace  $W \subset V$ . It is called the linear span of the vectors  $\vec{v}_1, \dots, \vec{v}_n$ .<sup>7</sup>

<sup>6</sup>In fact, besides just pointwise addition and scalar multiplication of real valued functions, it is possible to also define pointwise multiplication of such functions. So the space  $\mathbb{R}^X$  also has even more algebraic structure than just that of a vector space over  $\mathbb{R}$  — it is also an algebra over  $\mathbb{R}$ .

<sup>7</sup>More generally, if  $S \subset V$  is any subset, then the subset consisting of all *finite* linear combinations of elements of  $S$  is still a vector subspace. This is denoted  $\text{span}(S) \subset V$ , and called the linear

Function spaces are often defined as subspaces of the space of all functions with a given domain and a codomain: instead of considering the space of arbitrary functions, we usually care about sufficiently well-behaved functions. The following typical example is very important.

**Example IV.8** (The space of continuous functions on an interval).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval. Consider the set

$$\mathcal{C}([a, b]) = \left\{ f: [a, b] \rightarrow \mathbb{R} \text{ continuous function} \right\}. \quad (\text{IV.16})$$

This set of continuous real-valued functions on  $[a, b]$  is obviously a subset of the space  $\mathbb{R}^{[a, b]}$  of all real valued functions on  $[a, b]$  from Example IV.5. Since the zero-function (constant) is continuous, and since pointwise sums and scalar multiples of continuous functions are continuous (by Proposition III.10), we see that  $\mathcal{C}([a, b])$  is a vector subspace in  $\mathbb{R}^{[a, b]}$ .

Let us finally give a few examples of subspaces of the sequence space of Example IV.3.

**Example IV.9** (The space of absolutely summable sequences).

Consider the subset

$$\ell^1 = \left\{ (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \sum_{n=1}^{\infty} |x_n| < \infty \right\} \subset \mathbb{R}^{\mathbb{N}}$$

of absolutely summable real sequences in the vector space  $\mathbb{R}^{\mathbb{N}}$  of all real number sequences (Example IV.3). The zero sequence  $(0, 0, 0, \dots)$  is of course absolutely summable, since  $\sum_{n=1}^{\infty} |0| = \sum_{n=1}^{\infty} 0 = 0 < \infty$ . Also if  $(x_n)_{n \in \mathbb{N}} \in \ell^1$  is an absolutely summable sequence, then multiplying it by a scalar  $c \in \mathbb{R}$  (coordinatewise), we get the sequence  $c(x_n)_{n \in \mathbb{N}} = (cx_n)_{n \in \mathbb{N}}$  which is absolutely summable since

$$\sum_{n=1}^{\infty} |cx_n| = \sum_{n=1}^{\infty} |c| |x_n| = |c| \underbrace{\sum_{n=1}^{\infty} |x_n|}_{< \infty} < \infty.$$

Finally, if  $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in \ell^1$  are two absolutely summable sequences, then their sum (coordinatewise) is the sequence  $(x_n)_{n \in \mathbb{N}} + (y_n)_{n \in \mathbb{N}} = (x_n + y_n)_{n \in \mathbb{N}}$  which is also absolutely summable since

$$\sum_{n=1}^{\infty} \underbrace{|x_n + y_n|}_{\leq |x_n| + |y_n|} \leq \sum_{n=1}^{\infty} (|x_n| + |y_n|) = \underbrace{\sum_{n=1}^{\infty} |x_n|}_{< \infty} + \underbrace{\sum_{n=1}^{\infty} |y_n|}_{< \infty} < \infty.$$

These three observations show that  $\ell^1 \subset \mathbb{R}^{\mathbb{N}}$  is a vector subspace.

**Exercise IV.1** (The space of bounded sequences).

Consider the set

$$\ell^\infty = \left\{ (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \exists r \geq 0 : \forall n \in \mathbb{N} : |x_n| \leq r \right\} \subset \mathbb{R}^{\mathbb{N}}$$

of all bounded real number sequences.

- Show that  $\ell^\infty \subset \mathbb{R}^{\mathbb{N}}$  is a vector subspace.
- Let  $\ell^1$  be as in Example IV.9. Show the inclusion  $\ell^1 \subset \ell^\infty$ , and conclude that the space of all absolutely summable real number sequences is a vector subspace of the space of all bounded real number sequences.

---

span of the subset  $S \subset \mathbb{V}$  — it is the smallest vector subspace of  $\mathbb{V}$ , which contains all vectors of  $S$ . Checking this is not difficult, but it is better suited for a course of linear algebra.

**Exercise IV.2** (The space of convergent sequences).

Show that the set of those real number sequences which are convergent is a vector subspace of the space  $\ell^\infty$  of all bounded real number sequences.

**Example IV.10** (The space of sequences tending to zero).

Show that the set of those real number sequences which converge to 0 is a vector subspace of the space of all convergent real number sequences.

**Exercise (✓) IV.3** (The set of sequences tending to a given value).

Show that the set of those real number sequences which converge to  $-\sqrt{\pi}$  is not a vector subspace of the space of all real number sequences.

*Hint:* Instead of  $-\sqrt{\pi}$ , you could use any non-zero value as the limit, and the conclusion would remain the same. This hints that even the easiest vector subspace property fails.

## IV.2. Inner product spaces

A vector space, by itself, is a purely algebraic notion. Applying the vector addition inductively, one can define finite sums of vectors, but it would be meaningless to form infinite series, for example. Infinite series, as well as for example continuity of functions, require some additional structure — in fact something that gives rise to a topology on the vector space.

Perhaps the most familiar of such additional structures is the notion of an inner product, from which a notion of length of vectors as well as a notion of angle between vectors can be obtained.<sup>8</sup>

### Axioms of inner product spaces

**Definition IV.11** (Inner product space).

Let  $V$  be a real vector space. A binary operation

$$\begin{aligned} V \times V &\rightarrow \mathbb{R} \\ (\vec{x}, \vec{y}) &\mapsto \langle \vec{x}, \vec{y} \rangle \end{aligned} \tag{IV.17}$$

is called an **inner product**, if it satisfies the following conditions

- (IP1)  $\langle \vec{x}, \vec{y} \rangle = \langle \vec{y}, \vec{x} \rangle$  for all  $\vec{x}, \vec{y} \in V$ ; (symmetricity)
- (IP2)  $\langle c\vec{x}, \vec{y} \rangle = c \langle \vec{x}, \vec{y} \rangle$  for all  $c \in \mathbb{R}$  and  $\vec{x}, \vec{y} \in V$ ; (linearity)<sup>9</sup>
- (IP3)  $\langle \vec{x} + \vec{y}, \vec{z} \rangle = \langle \vec{x}, \vec{z} \rangle + \langle \vec{y}, \vec{z} \rangle$  for all  $\vec{x}, \vec{y}, \vec{z} \in V$ ; (linearity)
- (IP4)  $\langle \vec{x}, \vec{x} \rangle \geq 0$  for all  $\vec{x} \in V$ ; (positive semi-definiteness)
- (IP5)  $\langle \vec{x}, \vec{x} \rangle = 0$  only if  $\vec{x} = \vec{0} \in V$ . (non-degeneracy)<sup>10</sup>

<sup>8</sup>Arguably the notion of angles is more geometric than topological, so inner products provide us with geometric as well as topological structure.

<sup>9</sup>The linearity of an inner product with respect to its first argument includes two conditions: that it respects both scalar multiplication (IP2) and vector addition (IP3). These can be expressed concisely together in the property that  $\langle c_1u_1 + c_2u_2, v \rangle = c_1 \langle u_1, v \rangle + c_2 \langle u_2, v \rangle$  for all  $u_1, u_2, v \in V$  and  $c_1, c_2 \in \mathbb{R}$ . Moreover, if we combine this with symmetricity (IP1), we also get  $\langle u, d_1v_1 + d_2v_2 \rangle = d_1 \langle u, v_1 \rangle + d_2 \langle u, v_2 \rangle$  for all  $u, v_1, v_2 \in V$  and  $d_1, d_2 \in \mathbb{R}$ . This yields **bilinearity** of the inner product: an inner product is linear in both of its arguments.

<sup>10</sup>In fact we have  $\langle \vec{x}, \vec{x} \rangle = 0$  if and only if  $\vec{x} = \vec{0}$ . In (IP5) we only require the “only if” implication in order to minimize what needs to be checked in applications. The other direction



A vector space  $V$  equipped with an inner product is called an **inner product space**.

To concisely and unambiguously mention both the vector space  $V$  and the inner product  $\langle \cdot, \cdot \rangle$  on it, we may refer to the *pair*  $(V, \langle \cdot, \cdot \rangle)$  as an inner product space.

### Examples of inner product spaces

**Example IV.12** (Euclidean spaces).

In the vector space  $\mathbb{R}^d$  of Example IV.2, the formula

$$\langle \vec{x}, \vec{y} \rangle = \sum_{j=1}^d x_j y_j \quad \text{for } \vec{x} = (x_1, \dots, x_d), \vec{y} = (y_1, \dots, y_d) \in \mathbb{R}^d \quad (\text{IV.18})$$

defines the usual inner product — verifying the properties is straightforward. The corresponding inner product space is called the  $d$ -dimensional **Euclidean space**.

Let us verify the properties (IP1) – (IP5) in this familiar case.

(IP1): For any  $\vec{x} = (x_1, \dots, x_d), \vec{y} = (y_1, \dots, y_d) \in \mathbb{R}^d$  we have

$$\langle \vec{x}, \vec{y} \rangle = \sum_{j=1}^d x_j y_j = \sum_{j=1}^d y_j x_j = \langle \vec{y}, \vec{x} \rangle.$$

(IP2): For any  $\vec{x} = (x_1, \dots, x_d), \vec{y} = (y_1, \dots, y_d) \in \mathbb{R}^d$  and  $c \in \mathbb{R}$ , we have first of all  $c\vec{x} = (cx_1, \dots, cx_d)$  and then

$$\langle c\vec{x}, \vec{y} \rangle = \sum_{j=1}^d (cx_j) y_j = c \sum_{j=1}^d x_j y_j = c \langle \vec{x}, \vec{y} \rangle.$$

(IP3): For any  $\vec{x} = (x_1, \dots, x_d), \vec{y} = (y_1, \dots, y_d), \vec{z} = (z_1, \dots, z_d) \in \mathbb{R}^d$ , we have first of all  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_d + y_d)$  and then

$$\begin{aligned} \langle \vec{x} + \vec{y}, \vec{z} \rangle &= \sum_{j=1}^d (x_j + y_j) z_j = \sum_{j=1}^d (x_j z_j + y_j z_j) \\ &= \sum_{j=1}^d x_j z_j + \sum_{j=1}^d y_j z_j = \langle \vec{x}, \vec{z} \rangle + \langle \vec{y}, \vec{z} \rangle. \end{aligned}$$

(IP4): For any  $\vec{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$  we have, by nonnegativity of squares of real numbers,

$$\langle \vec{x}, \vec{x} \rangle = \sum_{j=1}^d x_j x_j = \sum_{j=1}^d x_j^2 \geq 0,$$

(IP5): From the formula above, we see that  $\langle \vec{x}, \vec{x} \rangle$  can only be zero for  $\vec{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$ , if for each  $j \in \{1, \dots, d\}$  we have  $x_j^2 = 0$ . This only occurs if each component  $x_j$  is zero and the vector is  $\vec{x} = (0, \dots, 0) = \vec{0}$ .

---

follows from already required properties. Namely, in any vector space we have  $\vec{0} = 0\vec{0}$  (you can derive this from the properties of Definition IV.1 as an algebraic exercise), so using (IP2) we get that all inner products with the zero vector  $\vec{0}$  vanish:

$$\langle \vec{0}, \vec{v} \rangle = \langle 0\vec{0}, \vec{v} \rangle = 0 \langle \vec{0}, \vec{v} \rangle = 0.$$

In particular, the zero vector has zero inner product with itself, giving the “if” implication.

**Example IV.13** (An inner product in a function space).

Consider the vector space  $\mathcal{C}([a, b])$  of continuous real-valued functions on a closed interval  $[a, b] \subset \mathbb{R}$ , which is nondegenerate,  $a < b$ . Define  $\langle \cdot, \cdot \rangle: \mathcal{C}([a, b]) \times \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  by

$$\langle f, g \rangle = \int_a^b f(t)g(t) dt \quad \text{for } f, g \in \mathcal{C}([a, b]). \quad (\text{IV.19})$$

This is called the  $L^2$ -inner product of functions. Let us verify that this is indeed an inner product on  $\mathcal{C}([a, b])$ . Note that the verifications are very similar to those in Example IV.12; only in (IP5) an essentially new observation is used.

(IP1): For any  $f, g \in \mathcal{C}([a, b])$  we have

$$\langle f, g \rangle = \int_a^b f(t)g(t) dt = \int_a^b g(t)f(t) dt = \langle g, f \rangle.$$

(IP2): For any  $f, g \in \mathcal{C}([a, b])$  and  $c \in \mathbb{R}$ , recalling the definition  $(cf)(t) = cf(t)$  of pointwise scalar multiplication of functions, we get

$$\langle cf, g \rangle = \int_a^b (cf)(t)g(t) dt = \int_a^b cf(t)g(t) dt = c \int_a^b f(t)g(t) dt = c \langle f, g \rangle.$$

(IP3): For any  $f, g, h \in \mathcal{C}([a, b])$ , recalling the definition  $(f+g)(t) = f(t) + g(t)$  of pointwise addition of functions, we get

$$\begin{aligned} \langle f+g, h \rangle &= \int_a^b (f+g)(t)h(t) dt \\ &= \int_a^b (f(t) + g(t))h(t) dt \\ &= \int_a^b f(t)h(t) dt + \int_a^b g(t)h(t) dt = \langle f, h \rangle + \langle g, h \rangle. \end{aligned}$$

(IP4): For any  $f \in \mathcal{C}([a, b])$ , we have

$$\langle f, f \rangle = \int_a^b \underbrace{f(t)f(t)}_{=f(t)^2 \geq 0} dt \geq 0.$$

(IP5): From the formula above, we see that  $\langle f, f \rangle$  can only be zero for  $f \in \mathcal{C}([a, b])$  if  $\int_a^b f(t)^2 dt = 0$ . Since  $t \mapsto f(t)^2$  is a continuous nonnegative function on  $[a, b]$ , by the contrapositive of Exercise III.10, the vanishing of this integral implies that  $f(t)^2 = 0$  for all  $t \in [a, b]$ . But this implies that  $f(t) = 0$  for all  $t \in [a, b]$ , i.e., that  $f$  is the zero function.

The combination of the following two exercises amount to a complete characterization of inner products in finite-dimensional vector spaces.

**Exercise IV.4** (Constructing inner products on  $\mathbb{R}^d$ ).

Let  $A \in \mathbb{R}^{d \times d}$  be a symmetric<sup>11</sup> positive definite<sup>12</sup>  $d \times d$  matrix. Show that the formula  $\langle x, y \rangle = x^\top A y$  for  $x, y \in \mathbb{R}^d$  defines an inner product on  $\mathbb{R}^d$ .

**Exercise IV.5** (Characterizing inner products on  $\mathbb{R}^d$ ).

Let  $\langle \cdot, \cdot \rangle$  be an inner product on  $\mathbb{R}^d$  (not necessarily the standard Euclidean inner product, but any inner product that satisfies the requirements in Definition IV.11). Let  $\mathbf{e}_1, \dots, \mathbf{e}_d$  be the standard basis vectors of  $\mathbb{R}^d$ . For all  $i, j \in \{1, \dots, d\}$ , define  $a_{i,j} = \langle \mathbf{e}_i, \mathbf{e}_j \rangle$ , and let  $A \in \mathbb{R}^{d \times d}$  denote the matrix with these entries  $a_{i,j}$ .

<sup>11</sup>A square matrix  $A$  is symmetric if it is its own transpose,  $A^\top = A$ .

<sup>12</sup>A square matrix  $A$  is said to be **positive semi-definite** if  $x^\top A x \geq 0$  for all vectors  $x$ , and **positive definite** if  $x^\top A x > 0$  for all nonzero vectors  $x$ .

- (a) Show that with this definition, the inner product of any two vectors  $x, y \in \mathbb{R}^d$  can be written as  $\langle x, y \rangle = x^\top A y$ .
- (b) Prove that the matrix  $A$  is symmetric and positive definite.

### The norm induced by an inner product

An inner product in particular yields a notion of length of vectors, i.e., a norm. In an inner product space  $V$ , we define the **norm**  $\|v\|$  of  $v \in V$  as

$$\|v\| = \sqrt{\langle v, v \rangle}. \quad (\text{IV.20})$$

**Example IV.14** (The Euclidean norm).

The  $d$ -dimensional Euclidean space is the vector space  $\mathbb{R}^d$  equipped with the inner product of Example IV.12. The norm of  $\vec{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$  in this space takes the form

$$\|\vec{x}\| = \sqrt{\langle \vec{x}, \vec{x} \rangle} = \sqrt{x_1^2 + \dots + x_d^2}. \quad (\text{IV.21})$$

**Example IV.15** (The  $L^2$ -norm on the space of continuous functions on a closed interval).

Recall from Example IV.13 that the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on a nondegenerate closed interval  $[a, b] \subset \mathbb{R}$  is an inner product space with inner product (IV.19). We denote the norm induced by it by  $\|\cdot\|_2$ . Explicitly for a continuous function  $f: [a, b] \rightarrow \mathbb{R}$ , this norm reads

$$\|f\|_2 = \sqrt{\langle f, f \rangle} = \left( \int_a^b f(t)^2 dt \right)^{1/2}.$$

Note, however, that we will soon give a more general meaning to the term *norm* — formula (IV.20) is a special instance, which is only meaningful in inner product spaces (indeed, the right hand side features the inner product explicitly). To emphasize that (IV.20) is not just any norm, we also call it *the norm induced by the inner product*  $\langle \cdot, \cdot \rangle$ .

For perspective, we mention already that the norm, by giving the lengths  $\|v\|$  of vectors  $v \in V$ , can be used also to define distances  $\|u - v\|$  between vectors  $u, v \in V$ . It thus equips the space  $V$  with topological structure. We return to this in more detail in Lecture V.

A crucial property of inner products and the norms induced by them is the following inequality.

**Theorem IV.16** (Cauchy-Schwarz inequality in inner product spaces).

Let  $V$  be an inner product space. Then for any  $u, v \in V$  we have

$$|\langle u, v \rangle| \leq \|u\| \|v\|. \quad (\text{IV.22})$$

*Proof.* If either  $u = 0$  or  $v = 0$ , then both sides of (IV.22) are zero and the inequality holds trivially. We may therefore assume that  $u, v \in V \setminus \{0\}$ .

Observe that for any  $t \in \mathbb{R}$ , by (IP4) for the vector  $u + tv \in V$  we have  $0 \leq \langle u + tv, u + tv \rangle$ . Using bilinearity of the inner product, we can write this as

$$\begin{aligned} 0 &\leq \langle u + tv, u + tv \rangle \\ &= \langle u, u \rangle + \langle u, tv \rangle + \langle tv, u \rangle + \langle tv, tv \rangle \\ &= \|u\|^2 + 2t \langle u, v \rangle + t^2 \|v\|^2. \end{aligned}$$

This last expression is a polynomial  $at^2 + bt + c$ , with coefficients  $a = \|v\|^2$ ,  $b = 2\langle u, v \rangle$ , and  $c = \|u\|^2$ . The fact that polynomial is non-negative for all  $t \in \mathbb{R}$  means that it has either no real roots or just one real root, so its discriminant is non-positive,  $b^2 - 4ac \leq 0$ . By plugging in the coefficients  $a, b, c$ , this reads

$$4(\langle u, v \rangle)^2 - 4\|u\|^2\|v\|^2 \leq 0.$$

Dividing by 4 and rearranging we get  $(\langle u, v \rangle)^2 \leq \|u\|^2\|v\|^2$ . Then by taking square roots we get (IV.22).  $\square$

**Exercise IV.6** (When does equality occur in the Cauchy-Schwarz inequality?).

- (a) Show that if  $u, v \in V$  are such that  $u = cv$  for some  $c \in \mathbb{R}$ , then equality occurs in the Cauchy-Schwarz inequality (IV.22) for these vectors.
- (b) By carefully examining the steps of the proof of Theorem IV.16, show that for two vectors  $u, v \in V$ , equality can occur in the Cauchy-Schwarz inequality only if one of them is a real scalar multiple of the other.

The Cauchy-Schwarz inequality is a tool used in numerous circumstances, but already the following analogue of Proposition II.1 indicates its importance.

**Corollary IV.17** (Triangle inequality in inner product spaces).

*Let  $V$  be an inner product space. Then for any  $u, v \in V$  we have*

$$\|u + v\| \leq \|u\| + \|v\|. \quad (\text{IV.23})$$

*Proof.* The proof is very similar to that of Proposition II.1. We first expand the square of  $\|u + v\|$  using bilinearity, and then estimate it using (IV.22) as follows

$$\begin{aligned} \|u + v\|^2 &= \langle u + v, u + v \rangle && (\text{definition}) \\ &= \langle u, u \rangle + \langle u, v \rangle + \langle v, u \rangle + \langle v, v \rangle && (\text{bilinearity}) \\ &= \|u\|^2 + 2\langle u, v \rangle + \|v\|^2 && (\text{symmetricity}) \\ &\leq \|u\|^2 + 2|\langle u, v \rangle| + \|v\|^2 && (\text{taking absolute value}) \\ &\leq \|u\|^2 + 2\|u\|\|v\| + \|v\|^2 && (\text{Cauchy-Schwarz}) \\ &= (\|u\| + \|v\|)^2. && (\text{binomial formula}) \end{aligned}$$

Taking the square roots, we get (IV.23).  $\square$

**Exercise IV.7** (Parallelogram rule).

Let  $V$  be an inner product space. Show that for any two vectors  $u, v \in V$  we have the following parallelogram rule

$$\|u + v\|^2 + \|u - v\|^2 = 2\|u\|^2 + 2\|v\|^2. \quad (\text{IV.24})$$

Let us also indicate some direct applications of the Cauchy-Schwarz inequality.

**Exercise IV.8** (A sum bound based on Cauchy-Schwarz inequality).

Let  $x_1, \dots, x_d \in \mathbb{R}$ . Prove the following inequality

$$|x_1| + \dots + |x_d| \leq \sqrt{d} \sqrt{x_1^2 + \dots + x_d^2}.$$

Hint: Consider the Euclidean space  $\mathbb{R}^d$ , and apply the Cauchy-Schwarz inequality with  $u = (1, 1, \dots, 1)$  and a suitably chosen  $v$ .

**Example IV.18** (Cauchy-Schwarz inequality in a function space).

Recall from Examples IV.13 and IV.15 that the space  $\mathcal{C}([a, b])$  of continuous real-valued

functions on a nondegenerate closed interval  $[a, b] \subset \mathbb{R}$  is an inner product space. The Cauchy-Schwarz inequality in the inner product space  $\mathcal{C}([a, b])$  amounts to the following integral inequality

$$\left| \int_a^b f(t) g(t) dt \right| \leq \left( \int_a^b f(t)^2 dt \right)^{1/2} \left( \int_a^b g(t)^2 dt \right)^{1/2}$$

for all continuous functions  $f, g: [a, b] \rightarrow \mathbb{R}$ .

As a special case, if  $h \in \mathcal{C}([a, b])$  and we take  $f = |h|$  and  $g = 1$ , we get

$$\int_a^b |h(t)| dt \leq \sqrt{b-a} \left( \int_a^b h(t)^2 dt \right)^{1/2}.$$

**Example IV.19** (The sequence space  $\ell^2$ ).

Consider the set

$$\ell^2 = \left\{ x = (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \sum_{n=1}^{\infty} x_n^2 < \infty \right\}$$

of square-summable real sequences. We can use Cauchy-Schwarz inequality in  $\mathbb{R}^d$  to show that first of all  $\ell^2 \subset \mathbb{R}^{\mathbb{N}}$  is a vector subspace and moreover it is itself an inner product space.

**UNDER CONSTRUCTION!**

## Angles and orthogonality

Let  $V$  be an inner product space.

Suppose that  $u, v \in V \setminus \{0\}$  are two non-zero vectors. Then by the Cauchy-Schwarz inequality (Theorem IV.16) we have

$$-1 \leq \frac{\langle u, v \rangle}{\|u\| \|v\|} \leq 1.$$

By bijectivity of  $\cos: [0, \pi] \rightarrow [-1, 1]$ , there therefore exists a unique  $\theta \in [0, \pi]$  such that  $\frac{\langle u, v \rangle}{\|u\| \|v\|} = \cos(\theta)$ , i.e., that

$$\langle u, v \rangle = \cos(\theta) \|u\| \|v\|. \quad (\text{IV.25})$$

This  $\theta \in [0, \pi]$  is called the **angle** between the vectors  $u$  and  $v$ . In the Euclidean spaces  $\mathbb{R}^2$  and  $\mathbb{R}^3$  this of course coincides with the familiar notion of (undirected<sup>13</sup>) angle.

The case of the right angle  $\theta = \frac{\pi}{2}$  is special. Since  $\cos(\frac{\pi}{2}) = 0$ , it occurs when the inner product between two vectors vanishes. Correspondingly, we make the following definitions.

**Definition IV.20** (Orthogonality and orthonormality).

Two vectors  $u, v \in V$  are said to be **orthogonal** if their inner product vanishes,  $\langle u, v \rangle = 0$ .

A collection  $S \subset V$  is said to be **orthogonal** if any two different vectors  $u, v \in S$ ,  $u \neq v$ , are orthogonal.

<sup>13</sup>In the plane  $\mathbb{R}^2$  there is a convention about what is the positive orientation, and it is natural to define also the directed angle from  $u \in \mathbb{R}^2 \setminus \{0\}$  to  $v \in \mathbb{R}^2 \setminus \{0\}$ , which is a number  $\phi \in (-\pi, \pi]$ . But the sign of this directed angle changes if we interchange  $u$  and  $v$ . In a general inner product space, two linearly independent vectors  $u, v$  span a plane, but this plane may be given either one of two possible orientations. The undirected angle does not depend on the choice of the orientation.

A collection  $S \subset V$  is said to be **orthonormal** if it is an orthogonal collection and all  $v \in S$  have unit norm,  $\|v\| = 1$ .

The cases of angles  $\theta = 0$  and  $\theta = \pi$  are also special: they correspond to vectors that lie on the same line through the origin.

**Lemma IV.21** (Collinear vectors).

Let  $u, v \in V \setminus \{0\}$  be two non-zero vectors in an inner product space  $V$ , and let  $\theta$  be the angle between them. Then we have

- $\theta = 0$  if and only if  $u = cv$  for some  $c > 0$ ;
- $\theta = \pi$  if and only if  $u = cv$  for some  $c < 0$ .

**Exercise IV.9** (Proof of Lemma IV.21).

Prove the two statements made in Lemma IV.21.

Hint: The “if” directions of both statements are easy. For the “only if” statements, use Exercise IV.6.

**Exercise IV.10** (The dihedral angle of a regular tetrahedron).

Calculate the dihedral angle (the angle between two adjacent faces) of a regular tetrahedron using the following idea:

Place the vertices of the tetrahedron in the 4-dimensional space  $\mathbb{R}^4$  at points  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  ja  $\mathbf{e}_4$  (standard basis vectors).

- (i) Verify that the lengths of the edges  $\|\mathbf{e}_i - \mathbf{e}_j\|$ , for  $i \neq j$ , are all equal.
- (ii) By symmetry, the angle in question is the same as (for example) the angle between the vectors

$$\mathbf{u} = \mathbf{e}_4 - \frac{1}{2}(\mathbf{e}_1 + \mathbf{e}_2) \quad \text{and} \quad \mathbf{v} = \mathbf{e}_3 - \frac{1}{2}(\mathbf{e}_1 + \mathbf{e}_2),$$

so it is easily calculated.

Hint: Draw a figure! Answer:  $\arccos(1/3)$ .

There are various function spaces that have inner products, and in many cases appropriate orthogonal polynomials in such spaces play an important role. To get some flavor of this, the following exercise deals with the first few<sup>14</sup> Legendre polynomials.

**Exercise IV.11** (Some Legendre polynomials).

Consider the three first Legendre polynomials

$$\begin{aligned} P_0(x) &= 1, \\ P_1(x) &= x, \\ P_2(x) &= (3x^2 - 1)/2. \end{aligned}$$

- (a) Show that these polynomials are pairwise *orthogonal* with respect to the inner product

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x) \, dx,$$

i.e., that we have  $\langle P_i, P_j \rangle = 0$  for  $i \neq j$ .

- (b) Determine coefficients  $c_0, c_1, c_2 \in \mathbb{R}$  such that  $\|c_k P_k\| = 1$  for all  $k \in \{0, 1, 2\}$ , where as norm we use the one arising from the inner product in part (a).

<sup>14</sup>If you are more ambitious, you can try to continue and find polynomials of higher degree which satisfy similar orthogonality.

Also the idea of Fourier series is based on orthogonality of trigonometric functions in a suitable function space.

**Exercise IV.12** (Ideas behind Fourier series).

- (a) Consider a countable orthonormal collection  $(e_n)_{n \in \mathbb{N}}$  in an inner product space  $\mathbf{V}$ . Assume that a vector  $v \in \mathbf{V}$  can be expressed, for some  $m \in \mathbb{N}$ , as a (finite) linear combination

$$v = \alpha_1 e_1 + \cdots + \alpha_m e_m \quad \text{with some coefficients } \alpha_n \in \mathbb{R}.$$

Calculate the inner products between the vector  $v$  and the vectors  $e_n$  from the orthonormal collection, and deduce a formula for the coefficients  $\alpha_n$ .

- (b) Define functions  $c_0, s_1, c_1, s_2, c_2, s_3, c_3, \dots$  of a real variable  $x$  by the formulas

$$\begin{aligned} c_0(x) &= \frac{1}{\sqrt{2}}, \\ s_n(x) &= \sin(nx) \quad (\text{for } n \in \mathbb{N}), \\ c_n(x) &= \cos(nx) \quad (\text{for } n \in \mathbb{N}). \end{aligned}$$

Prove that the (countably infinite) collection of these functions is orthonormal in the space  $C([-\pi, \pi])$  of continuous functions on  $[-\pi, \pi]$ , with respect to the rescaled inner product

$$\langle f, g \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x)g(x) \, dx.$$

- (c) The idea of Fourier series is founded on the following result: Every continuously differentiable  $2\pi$ -periodic function  $f: \mathbb{R} \rightarrow \mathbb{R}$  can be represented as

$$f(x) = \alpha_0 + \sum_{n=1}^{\infty} (\alpha_n \cos(nx) + \beta_n \sin(nx)).$$

Consider for simplicity the case of a function  $f$ , for which the above series contains only finitely many terms, i.e., for some  $m \in \mathbb{N}$  we have

$$f(x) = \alpha_0 + \sum_{n=1}^m (\alpha_n \cos(nx) + \beta_n \sin(nx)).$$

Using parts (a) and (b), derive the following formula for the coefficients

$$\alpha_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) \, dx \quad (n \in \mathbb{N}).$$

### IV.3. Normed spaces

#### Axioms of normed spaces

**Definition IV.22** (Normed space).

Let  $\mathbf{V}$  be a real vector space. A function

$$\begin{aligned} \mathbf{V} &\rightarrow [0, \infty) \\ \vec{x} &\mapsto \|\vec{x}\| \end{aligned} \tag{IV.26}$$

is called a **norm**, if it satisfies the following conditions

- (N1)  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$  for all  $\vec{x}, \vec{y} \in \mathbf{V}$ ; (triangle inequality)  
 (N2)  $\|c \vec{x}\| = |c| \|\vec{x}\|$  for all  $c \in \mathbb{R}$  and  $\vec{x} \in \mathbf{V}$ ; (homogeneity)

(N3)  $\|\vec{x}\| = 0$  only if  $\vec{x} = \vec{0} \in V$ . (non-degeneracy)<sup>15</sup>

A vector space  $V$  equipped with a norm is called a **normed space**.

To concisely and unambiguously mention both the vector space  $V$  and the norm  $\|\cdot\|$  on it, we may refer to the *pair*  $(V, \|\cdot\|)$  as a normed space.

By the following two remarks, we in fact immediately get some examples of norms and normed spaces from Section IV.2.

**Remark IV.23** (The norm induced by an inner product is a norm).

In an inner product space, we used (IV.20) to define a norm using the inner product, and this gives rise to a norm also in the sense of Definition IV.22. Indeed, for the norm  $\|v\| = \sqrt{\langle v, v \rangle}$  induced by the inner product, we already saw in Corollary IV.17 that (N1) holds. The other two properties are easy. From (IP2) and (IP1) we get  $\langle cv, cv \rangle = c^2 \langle v, v \rangle$ , so

$$\|cv\| = \sqrt{c^2 \langle v, v \rangle} = |c| \sqrt{\langle v, v \rangle} = |c| \|v\|,$$

establishing (N2). The property (N3) follows directly from (IP5).

**Remark IV.24** (All inner product spaces are normed spaces).

It follows from the remark above that any inner product space becomes a normed space, when it is equipped with the norm induced by the inner product as in (IV.20).

In particular, the Euclidean space  $\mathbb{R}^d$  of Example IV.12 and the space  $\mathcal{C}([a, b])$  of continuous functions on a closed interval with the  $L^2$ -inner product of Example IV.13 can both be viewed as normed spaces as well.

## Examples of normed spaces

By Remark IV.24, any inner product space is an example of a normed space. However, the converse is not true: there are normed spaces that are not inner product spaces. The concept of a normed space is genuinely more general. Let us now look at some such examples.

One can in fact even equip the standard  $d$ -dimensional vector space  $\mathbb{R}^d$  with other norms besides the Euclidean norm on the inner product space of Example IV.12. First of all, we can use other inner products as indicated in Exercises IV.4 and IV.5, and they induced other norms on  $\mathbb{R}^d$ . But to focus on norms that are not induced by any inner product — here are some examples.

**Example IV.25** (The  $\ell^\infty$ -norm on  $\mathbb{R}^d$ ).

Define, for  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ ,

$$\|x\|_\infty = \max\{|x_1|, \dots, |x_d|\} = \max_{j=1, \dots, d} |x_j|. \quad (\text{IV.27})$$

We call this the  $\ell^\infty$ -**norm** on  $\mathbb{R}^d$  — but let us verify that this indeed is a norm!

(N1): Let  $x = (x_1, \dots, x_d)$  and  $y = (y_1, \dots, y_d)$  are two vectors in  $\mathbb{R}^d$ . By definition, their vector sum is  $x+y = (x_1+y_1, \dots, x_d+y_d)$ . For any  $j \in \{1, \dots, d\}$  we have  $|x_j+y_j| \leq |x_j|+|y_j|$

<sup>15</sup>We in fact have  $\|\vec{x}\| = 0$  if and only if  $\vec{x} = \vec{0}$ . In (N3) we only require the “only if” implication to minimize work to check these properties in examples. The other direction follows from already required properties. Namely, we have  $\vec{0} = 0\vec{0}$ , so by (N2) we get

$$\|\vec{0}\| = \|0\vec{0}\| = |0| \|\vec{0}\| = 0.$$



by the triangle inequality of real numbers (Proposition II.1), and we may further estimate

$$|x_j + y_j| \leq |x_j| + |y_j| \leq \max_{j=1,\dots,d} |x_j| + \max_{j=1,\dots,d} |y_j|.$$

Since this estimate holds for each  $j$ , we get

$$\max_{j=1,\dots,d} |x_j + y_j| \leq \max_{j=1,\dots,d} |x_j| + \max_{j=1,\dots,d} |y_j|,$$

which is exactly property (N1) for the  $\ell^\infty$  norm,  $\|x + y\|_\infty \leq \|x\|_\infty + \|y\|_\infty$ .

(N2): Let  $c \in \mathbb{R}$  and  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ . By definition of scalar multiplication, we have  $cx = (cx_1, \dots, cx_d)$ . For any  $j \in \{1, \dots, d\}$  we have

$$|cx_j| = |c| |x_j| \leq |c| \max_{j=1,\dots,d} |x_j|.$$

Since this estimate holds for each  $j$ , we get

$$\max_{j=1,\dots,d} |cx_j| \leq |c| \max_{j=1,\dots,d} |x_j|,$$

and by considering the index  $j$  for which the maximal  $x_j$  is attained, we see that equality in fact holds here. The equality is exactly property (N2) for the  $\ell^\infty$  norm,  $\|cx\|_\infty = |c| \|x\|_\infty$ .

(N3): If for  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$  we have  $\|x\|_\infty = 0$ , i.e.,  $\max\{|x_1|, \dots, |x_d|\} = 0$ , then for each  $j$  we must have  $x_j = 0$ , which implies that  $x = (0, \dots, 0)$ . This establishes (N3).

**Exercise IV.13** (The  $\ell^1$ -norm on  $\mathbb{R}^d$ ).

Define, for  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ ,

$$\|x\|_1 = |x_1| + \dots + |x_d| = \sum_{j=1}^d |x_j|. \quad (\text{IV.28})$$

We call this the  $\ell^1$ -**norm** on  $\mathbb{R}^d$ . Prove that it satisfies (N1), (N2), (N3).

**Exercise IV.14** (The “unit circles” in  $\mathbb{R}^2$  with different norms).

Consider the two-dimensional vector space  $\mathbb{R}^2$  equipped with the norms of Example IV.25 and Exercise IV.13.

(a) Draw the “unit circle” with respect to the  $\ell^\infty$ -norm,

$$\left\{ (x, y) \in \mathbb{R}^2 \mid \|(x, y)\|_\infty = 1 \right\}.$$

(b) Draw the “unit circle” with respect to the  $\ell^1$ -norm,

$$\left\{ (x, y) \in \mathbb{R}^2 \mid \|(x, y)\|_1 = 1 \right\}.$$

**Exercise IV.15** (The  $\ell^\infty$ -norm is not induced by an inner product).

Consider the norm  $\|\cdot\|_\infty$  of Example IV.25 on  $\mathbb{R}^d$  with  $d \geq 2$ . Find two vectors  $u, v \in \mathbb{R}^d$  such that  $\|u + v\|_\infty^2 + \|u - v\|_\infty^2 \neq 2\|u\|_\infty^2 + 2\|v\|_\infty^2$ . Use Exercise IV.7 to conclude that the norm  $\|\cdot\|_\infty$  is not induced by any inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{R}^d$ .

**Exercise IV.16** (The  $\ell^1$ -norm is not induced by an inner product).

Consider the norm  $\|\cdot\|_1$  of Exercise IV.13 on  $\mathbb{R}^d$  with  $d \geq 2$ . Show that the norm  $\|\cdot\|_1$  is not induced by any inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{R}^d$ .

So neither of the norms  $\|\cdot\|_\infty$  and  $\|\cdot\|_1$  on  $\mathbb{R}^d$  comes from an inner product (when  $d \geq 2$ ). By the following exercises, both are comparable up to constant multiples depending on the dimension  $d$ , and moreover both are similarly comparable to the Euclidean norm induced by the standard inner product.

**Exercise IV.17** (A comparison of the  $\ell^1$ -norm and the  $\ell^\infty$ -norm).

Let  $\|\cdot\|_\infty$  and  $\|\cdot\|_1$  denote the norms of Example IV.25 and Exercise IV.13 on  $\mathbb{R}^d$ . Prove that for any  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ , we have

$$\|x\|_\infty \leq \|x\|_1 \leq d \|x\|_\infty. \quad (\text{IV.29})$$

**Exercise IV.18** (A comparison of the  $\ell^1$ -norm and the Euclidean norm).

Let  $\|\cdot\|$  denote the ordinary Euclidean norm on  $\mathbb{R}^d$ , induced by the standard inner product, and let  $\|\cdot\|_1$  denote the  $\ell^1$ -norm of Exercise IV.13. Prove that for any  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ , we have

$$\|x\| \leq \|x\|_1 \leq \sqrt{d} \|x\|.$$

*Hint: Recall Exercise IV.8.*

**Exercise IV.19** (A comparison of the  $\ell^\infty$ -norm and the Euclidean norm).

Show that there exists a constant  $C_d > 0$  such that for any  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ , we have

$$\frac{1}{C_d} \|x\| \leq \|x\|_\infty \leq C_d \|x\|.$$

Both of the above are examples of  $\ell^p$ -norms.

**Example IV.26** (The  $\ell^p$ -norm on  $\mathbb{R}^d$ ).

Let  $p \in [1, \infty)$ . Define, for  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ ,

$$\|x\|_p = \left( \sum_{j=1}^d |x_j|^p \right)^{1/p}. \quad (\text{IV.30})$$

We call this the  $\ell^p$ -**norm** on  $\mathbb{R}^d$ . It satisfies (N1), (N2), (N3), but verifying (N1) requires slightly more work than in the cases above. This is a generalization which includes, at specific values of  $p \geq 1$ , some cases we have seen already:

- In the case  $p = 1$ , from (IV.30) we clearly recover the  $\ell^1$ -norm (IV.28) of Exercise IV.13.
- In the case  $p = 2$ , from (IV.30) we recover the Euclidean norm of Example IV.14,

$$\|x\|_2 = \left( \sum_{j=1}^d x_j^2 \right)^{1/2} = \sqrt{\langle x, x \rangle}.$$

- Exercise IV.20 will show that in the limit  $p \rightarrow \infty$ , we recover the  $\ell^\infty$ -norm of Example IV.25.

The case  $p = 2$  yields a norm induced by the inner product of Example IV.12, but for  $d \geq 2$  no other  $\ell^p$ -norm comes from an inner product.

**Exercise IV.20** (The  $\ell^p$ -norm as  $p \rightarrow \infty$ ).

On  $\mathbb{R}^d$ , consider the  $\ell^\infty$ -norm  $\|\cdot\|_\infty$  of Example IV.25, and the  $\ell^p$ -norms  $\|\cdot\|_p$  of Example IV.26, for  $p \geq 1$ .

- Prove that for any  $\vec{x} \in \mathbb{R}^d$  and any  $p \geq 1$ , we have  $\|\vec{x}\|_\infty \leq \|\vec{x}\|_p$ .
- Prove that for any  $\vec{x} \in \mathbb{R}^d$  and any  $p \geq 1$ , we have  $\|\vec{x}\|_p \leq d^{1/p} \|\vec{x}\|_\infty$ .
- Use (a), (b), and the calculation of  $\lim_{p \rightarrow \infty} d^{1/p}$  to prove that for any  $\vec{x} \in \mathbb{R}^d$  we have

$$\lim_{p \rightarrow \infty} \|\vec{x}\|_p = \|\vec{x}\|_\infty.$$

Also many important function spaces are equipped with norms that are not induced by any inner product. Let us give some such examples. The next one, in particular, is very frequently used in applications.

**Example IV.27** (The supremum norm on the space of continuous functions).

Consider again the vector space  $\mathcal{C}([a, b])$  of continuous real-valued functions on a closed interval  $[a, b] \subset \mathbb{R}$ . For  $f \in \mathcal{C}([a, b])$ , let us define<sup>16</sup>

$$\begin{aligned}\|f\|_\infty &= \sup \left\{ |f(t)| \mid t \in [a, b] \right\} \\ &= \sup_{t \in [a, b]} |f(t)|.\end{aligned}\tag{IV.31}$$

Note that since the continuous function  $f$  on a closed interval  $[a, b]$  is bounded (Theorem III.13(i)), the supremum here is finite,  $\|f\|_\infty \in [0, \infty)$ . In fact, since the continuous function  $t \mapsto |f(t)|$  on the closed interval  $[a, b]$  attains its maximum (Theorem III.13(ii)), we in fact have  $\|f\|_\infty = \max \{ |f(t)| \mid t \in [a, b] \}$ .

We call  $\|\cdot\|_\infty$  the **supremum norm**, the **sup-norm**, or the **uniform norm** — but let us verify that it indeed is a norm.

(N1): Let  $f, g \in \mathcal{C}([a, b])$ . Recall that addition  $f + g$  in  $\mathcal{C}([a, b])$  is defined pointwise by  $(f + g)(t) = f(t) + g(t)$ . For any  $t \in [a, b]$ , using the triangle inequality of real numbers (Proposition II.1), we get

$$\begin{aligned}|(f + g)(t)| &= |f(t) + g(t)| \\ &\leq |f(t)| + |g(t)| \leq \sup_{t \in [a, b]} |f(t)| + \sup_{t \in [a, b]} |g(t)| = \|f\|_\infty + \|g\|_\infty.\end{aligned}$$

Since this holds for any  $t \in [a, b]$ , we also get

$$\|f + g\|_\infty = \sup_{t \in [a, b]} |(f + g)(t)| \leq \|f\|_\infty + \|g\|_\infty,$$

which is exactly the triangle inequality (N1) for the norm  $\|\cdot\|_\infty$ .

(N2): Let  $f \in \mathcal{C}([a, b])$  and  $c \in \mathbb{R}$ . Recall that scalar multiplication  $cf$  in  $\mathcal{C}([a, b])$  is defined pointwise by  $(cf)(t) = c f(t)$ . For  $t \in [a, b]$ , we have

$$|(cf)(t)| = |c f(t)| = |c| |f(t)| \leq |c| \sup_{t \in [a, b]} |f(t)| = |c| \|f\|_\infty,$$

and since this holds for all  $t \in [a, b]$ , we get

$$\|cf\|_\infty = \sup_{t \in [a, b]} |(cf)(t)| \leq |c| \|f\|_\infty.$$

This inequality is only a half of the desired equality (N2), but we can use it as follows. If  $c = 0$ , then  $cf = 0$  is the constant function zero, so clearly  $\|cf\|_\infty = \|0\|_\infty = 0 = |0| \|f\|_\infty$  as desired. Assume therefore that  $c \neq 0$ . Then by using the already established inequality with scalar  $\frac{1}{c}$  and function  $cf$ , we get have

$$\|f\|_\infty = \left\| \frac{1}{c} (cf) \right\|_\infty \leq \frac{1}{|c|} \|cf\|_\infty,$$

which upon rearranging yields the inequality  $\|cf\|_\infty \geq |c| \|f\|_\infty$  in the other direction. The equality (N2) follows.

(N3): If  $f \in \mathcal{C}([a, b])$  satisfies  $\|f\|_\infty = 0$ , then  $\sup_{t \in [a, b]} |f(t)| = 0$  so we cannot have  $|f(t)| > 0$  for any  $t$ , i.e., we have  $|f(t)| = 0$  for every  $t$ , and  $f$  is the constant function zero.

**Exercise (#) IV.21** (Yet another norm on the space of continuous functions).

Consider again the vector space  $\mathcal{C}([a, b])$  of continuous real-valued functions on a nondegenerate closed interval  $[a, b] \subset \mathbb{R}$ . For  $f \in \mathcal{C}([a, b])$ , let us define

$$\|f\|_1 = \int_a^b |f(t)| dt.\tag{IV.32}$$

Verify that  $\|\cdot\|_1$  is a norm on  $\mathcal{C}([a, b])$ . We call it the  **$L^1$ -norm**.

<sup>16</sup>The second line of the formula here means exactly the same as the first line, but it is a more concise notation and as such often very convenient.

Contrary to the  $L^2$ -norm  $\|\cdot\|_2$  of Example IV.15 on  $\mathcal{C}([a, b])$ , neither of the above norms is induced by an inner product.

**Exercise IV.22** (The sup-norm does not come from an inner product).

Prove that if the interval  $[a, b]$  is non-degenerate,  $a < b$ , then the sup-norm  $\|\cdot\|_\infty$  on  $\mathcal{C}([a, b])$  is not induced by any inner product.

**Exercise IV.23** (The  $L^1$ -norm does not come from an inner product).

Prove that if the interval  $[a, b]$  is non-degenerate,  $a < b$ , then the  $L^1$ -norm  $\|\cdot\|_1$  on  $\mathcal{C}([a, b])$  is not induced by any inner product.

**Exercise IV.24** (A non-norm).

UNDER CONSTRUCTION!

**Exercise IV.25** (Another non-norm).

UNDER CONSTRUCTION!

**Exercise IV.26** (The sequence space  $\ell^1$ ).

Consider the vector space

$$\ell^1 = \left\{ x = (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \sum_{n=1}^{\infty} |x_n| < \infty \right\}$$

of absolutely summable real sequences (Example IV.9). For  $x = (x_n)_{n \in \mathbb{N}} \in \ell^1$ , define

$$\|x\|_1 = \sum_{n=1}^{\infty} |x_n|. \quad (\text{IV.33})$$

Show that  $\|\cdot\|_1$  is a norm on  $\ell^1$ .

**Exercise IV.27** (The sequence space  $\ell^\infty$ ).

Consider the vector space

$$\ell^\infty = \left\{ x = (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \exists r \geq 0 : \forall n \in \mathbb{N} : |x_n| \leq r \right\}$$

of bounded real sequences (Exercise IV.1). For  $x = (x_n)_{n \in \mathbb{N}} \in \ell^\infty$ , define

$$\|x\|_\infty = \sup \{ |x_n| \mid n \in \mathbb{N} \}. \quad (\text{IV.34})$$

Show that  $\|\cdot\|_\infty$  is a norm on  $\ell^\infty$ .

## Lecture V

### Metric spaces

In this lecture we reach the level of generality at which we will work on almost exclusively from here on: we introduce the notion of a *metric space*. The spaces we have considered so far — the real line, the Euclidean space, more general inner product spaces and normed spaces — all turn out to be at least metric spaces. But there is a vast amount of other relevant examples of metric spaces, too! What is important is that fruitful general theory for metric spaces can be developed, which can then be applied to the all of the specific cases.

The development of the theory will be mainly postponed to the subsequent lectures. The goal of this lecture is to just define metric spaces, give sufficiently many interesting examples of them, and introduce a few first concepts that make sense in general metric spaces. The concepts of open and closed sets in metric spaces turn out to be the most important for the development of the theory, although their importance may not yet be evident at the first sight.

#### V.1. Axioms of metric spaces

Intuitively, a metric space is a space where there is a reasonable notion of *distance* between points. The distances are encoded in a function called a metric. The only requirements, informally described, are: (*symmetry*) that the distance between two points is the same both ways, (*triangle inequality*) that the distance from one point to another cannot be reduced by taking a “shortcut” through a third point, and (*separation*) that the distance between two different points is always strictly positive, while the distance from a point to itself is zero.

The precise definition is the following.

**Definition V.1** (Metric space).

Let  $X$  be a set. A function

$$d: X \times X \rightarrow [0, \infty) \tag{V.1}$$

is a **metric** on  $X$  if the following conditions hold:

**(M-s):** For any  $x, y \in X$  we have (“symmetry”)

$$d(x, y) = d(y, x). \tag{M-s}$$

**(M-Δ):** For any  $x, y, z \in X$  we have (“triangle inequality”)

$$d(x, z) \leq d(x, y) + d(y, z). \tag{M-Δ}$$

**(M-0):** We have (“separation”)

$$d(x, y) = 0 \text{ if and only if } x = y. \tag{M-0}$$

A set  $X$  equipped with a metric  $d$  is called a **metric space**.

To concisely and unambiguously mention both the set  $X$  and the metric  $d$  on it, we may refer to the *pair*  $(X, d)$  as a metric space.

## V.2. Examples of metric spaces

Metric spaces are very general, and most spaces that we have seen so far are in fact also metric spaces. Let us start from something familiar and safe.

**Example V.2** (The real line is a metric space).

On the real line  $\mathbb{R}$ , we use the familiar notion of distances between points, encoded in the metric  $d_{\mathbb{R}}: \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty)$  on the set  $\mathbb{R}$  given by the formula

$$d_{\mathbb{R}}(x, y) = |y - x| \quad \text{for } x, y \in \mathbb{R}. \quad (\text{V.2})$$

Checking the properties (M-s), (M- $\Delta$ ), (M-0) directly is not difficult (it is a good exercise!), but the properties also follow from the more general Remark V.5 below.

We call (V.2) the **standard metric on  $\mathbb{R}$** . Unless otherwise mentioned, we consider the real axis  $\mathbb{R}$  equipped with this metric.

Although the above metric on  $\mathbb{R}$  is generally the reasonable choice that we will use without explicit mention, let us take this opportunity to point out that we *could* equip the set  $\mathbb{R}$  also with some other more exotic metrics.

**Exercise V.1** (Another metric on the real line).

Prove that the formula

$$d(x, y) = \log(1 + |x - y|)$$

also defines a metric  $d: \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty)$  on the set  $\mathbb{R}$ .

How creative can you get with metrics on the line?

**Exercise V.2** (Yet other metrics on the real line).

Find further examples of metrics on  $\mathbb{R}$ , besides those in Example V.2 and Exercise V.1.

Hint: First try to come up with some creative new formula! If that is too hard, a simple solution is to “change the units of distances” by multiplying a given metric<sup>1</sup> by some positive constant. Another idea that always works is to “allow for teleportation at fixed cost”<sup>2</sup> — a precise sense of which we will discuss in Lecture VII. Yet another idea is to define a simple  $\{0, 1\}$ -valued metric (see Example V.7).

Ok, all of this is to illustrate that the same set can be equipped with many possible metrics. But on  $\mathbb{R}$  the standard metric (V.2) is the important one!

Let us then look at the next most familiar case.

<sup>1</sup>Note that you can apply the “change of units” to for example the standard metric or the more exotic metric on Exercise V.1.

<sup>2</sup>Also the “teleportation” modification can be done to either the standard metric or some more exotic metric to start with.

**Example V.3** (The Euclidean spaces are metric spaces).

In the  $d$ -dimensional Euclidean space  $\mathbb{R}^d$ , the usual notion of distances comes from the Euclidean norm  $\|\cdot\|_2$  of (IV.21); the corresponding metric  $d_{\mathbb{R}^d}: \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$  is given by the familiar formula

$$d_{\mathbb{R}^d}(\vec{x}, \vec{y}) = \|\vec{y} - \vec{x}\|_2 = \sqrt{(y_1 - x_1)^2 + \cdots + (y_d - x_d)^2} \quad (\text{V.3})$$

for  $\vec{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$  and  $\vec{y} = (y_1, \dots, y_d) \in \mathbb{R}^d$ . One can check the properties (M-s), (M- $\Delta$ ), (M-0) directly, but they also follow from the more general Remark V.5 below.

When we refer to  $\mathbb{R}^d$  as a Euclidean space, we consider it equipped with this metric (V.3).

Just as with the real line, the Euclidean metric above is the usual reasonable choice, but there are other possible metrics as well. Let us give just one such example for now.

**Example V.4** (The Manhattan metric in the plane).

In the plane  $\mathbb{R}^2$ , the formula

$$d((x, y), (x', y')) = |x' - x| + |y' - y| \quad \text{for } (x, y), (x', y') \in \mathbb{R}^2 \quad (\text{V.4})$$

defines a metric  $d: \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow [0, \infty)$  — the required properties again follow using Remark V.5 for the norm  $\|\cdot\|_1$  of Exercise IV.13.

The metric (V.4) is sometimes referred to as the *Manhattan metric*, because the distance between two points is obtained as if one had to separately move along horizontal “streets” (the term  $|x' - x|$ ) and vertical “avenues” (the term  $|y' - y|$ ).

In the examples above we did not do detailed checks of the properties (M-s), (M- $\Delta$ ), (M-0). The reason is that they can all be handled at once with the following general observation, which in fact immediately gives also many further examples of metric spaces.

**Remark V.5** (All normed spaces are metric spaces).

Suppose that  $(V, \|\cdot\|)$  is a normed space. Then the **metric induced by the norm**  $\|\cdot\|$  is the metric  $d_V: V \times V \rightarrow [0, \infty)$  defined by the formula

$$d_V(\vec{u}, \vec{v}) = \|\vec{v} - \vec{u}\| \quad \text{for } \vec{u}, \vec{v} \in V. \quad (\text{V.5})$$

Let us check that  $d_V$  is indeed a metric on  $V$  in the sense of Definition V.1.

(M-s): Let  $\vec{u}, \vec{v} \in V$ . Noting that  $\vec{u} - \vec{v} = (-1)(\vec{v} - \vec{u})$  and using the homogeneity property (N2) of norms, we straightforwardly get the desired symmetry of  $d_V$

$$d_V(\vec{v}, \vec{u}) = \|\vec{u} - \vec{v}\| = \|(-1)(\vec{v} - \vec{u})\| = |-1| \|\vec{v} - \vec{u}\| = \|\vec{v} - \vec{u}\| = d_V(\vec{u}, \vec{v}).$$

(M- $\Delta$ ): Let  $\vec{u}, \vec{v}, \vec{w} \in V$ . Then using the triangle inequality (N1) for the norm  $\|\cdot\|$ , we get the desired triangle inequality of  $d_V$  as follows

$$\begin{aligned} d_V(\vec{u}, \vec{w}) &= \|\vec{w} - \vec{u}\| && \text{(definition of } d_V) \\ &= \|(\vec{w} - \vec{v}) + (\vec{v} - \vec{u})\| && \text{(added and subtracted } \vec{v}) \\ &\leq \|\vec{w} - \vec{v}\| + \|\vec{v} - \vec{u}\| && \text{(N1)} \\ &= d_V(\vec{v}, \vec{w}) + d_V(\vec{u}, \vec{v}). && \text{(definition of } d_V) \end{aligned}$$

(M-0): Suppose that  $\vec{u}, \vec{v} \in V$  are such that  $d_V(\vec{u}, \vec{v}) = 0$ . By definition of  $d_V$ , this means that  $\|\vec{v} - \vec{u}\| = 0$ . By property (N3) of the norm  $\|\cdot\|$ , this implies that  $\vec{v} - \vec{u} = 0$ . Adding  $\vec{u}$  to both sides we find  $\vec{v} = \vec{u}$ , as desired. The converse implication is easy: we have  $d_V(\vec{v}, \vec{v}) = \|\vec{v} - \vec{v}\| = \|\vec{0}\| = 0$  for any  $\vec{v} \in V$ .

The verification of the properties of metric in Examples V.2, V.3, and V.4 now follow directly by applying Remark V.5 to the normed spaces  $(\mathbb{R}, |\cdot|)$ ,  $(\mathbb{R}^d, \|\cdot\|_2)$ , and  $(\mathbb{R}^2, \|\cdot\|_1)$ , respectively.

We also get lots of other examples — just check again every example of a normed space from Lecture IV (including all examples of inner product spaces, in view of Remark IV.24). Let us just take one example explicitly.

**Example V.6** (A metric on the space of continuous function on a closed interval).

The space  $\mathcal{C}([a, b])$  on continuous real-valued functions on a closed interval  $[a, b]$ , equipped with the supremum norm  $\|\cdot\|_\infty$  of Example IV.27, is a normed space — and thus in particular a metric space by Remark V.5. The metric (V.5) induced by the supremum norm (IV.31) takes the form

$$d(f, g) = \|g - f\|_\infty = \sup_{t \in [a, b]} |g(t) - f(t)| \quad \text{for } f, g \in \mathcal{C}([a, b]). \quad (\text{V.6})$$

Let us finally mention an extremely simple metric you could use on any set.

**Example V.7** (The 0/1 metric).

Let  $X$  be any set. The formula

$$d_{0/1}(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases} \quad \text{for } x, y \in X \quad (\text{V.7})$$

is easily seen to define a metric on  $X$ . It is called the **0/1-metric** on  $X$ . It equips the set  $X$  with a structure that in some cases (discrete spaces) is reasonable, but in other cases is bizarre — there are no points “near each other”, the distance between all pairs of points is the same!

Nevertheless,  $(X, d_{0/1})$  a valid example of a metric space, it has some relevant applications, and it often provides simple and illuminating counterexamples.

## Metric inherited to subsets

Let  $(X, d)$  be a metric space, and  $X' \subset X$  any subset. Then the subset naturally inherits a metric from the whole space — since  $d$  gives distances between all pairs of points in  $X$ , it certainly also gives distances between all pairs of points in the subset  $X' \subset X$ . Formally, the **inherited metric**  $d': X' \times X' \rightarrow [0, \infty)$  on  $X'$  is given by

$$d'(x', y') = d(x', y') \quad \text{for } x', y' \in X' \subset X, \quad (\text{V.8})$$

i.e., the restriction of  $d: X \times X \rightarrow [0, \infty)$  to the subset  $X' \times X' \subset X \times X$ . The properties (M-s), (M- $\Delta$ ), and (M-0) for  $d'$  directly follow as special cases of the corresponding properties of  $d$ .

We now get lots of new examples of metric spaces!

**Example V.8** (Subsets of the real line are metric spaces).

Since the real line  $\mathbb{R}$  is a metric space (Example V.2), all its subsets also become metric spaces with the metrics they inherit by (V.8).

In particular, any of the following are metric spaces:

- intervals such as  $[0, 1] \subset \mathbb{R}$ ,  $(-\infty, 0) \subset \mathbb{R}$ ,  $(-\pi, \pi] \subset \mathbb{R}$ ;
- the set of rational numbers  $\mathbb{Q} \subset \mathbb{R}$ ;
- the set of irrational numbers  $\mathbb{R} \setminus \mathbb{Q} \subset \mathbb{R}$ ;





FIGURE V.1. The unit sphere  $\mathbf{S}^2 \subset \mathbb{R}^3$  inherits a metric from  $\mathbb{R}^3$ , but it also has another natural metric corresponding to the minimal length of paths along the surface.

- the Cantor set  $C \subset \mathbb{R}$  (Appendix B.3);
- etc. etc.

**Example V.9** (Subsets of the plane are metric spaces).

Since the Euclidean plane  $\mathbb{R}^2$  is a metric space (Example V.3), all its subsets also become metric spaces with the metrics they inherit by (V.8).

In particular, all lines, circles, curves, rectangles, triangles, disks, half-spaces, cones, the Koch snowflake, the Sierpinski carpet, ... in the plane  $\mathbb{R}^2$  are metric spaces.

You have to admit that the number of relevant examples of metric spaces is vast.

Since we happen to live on this (approximately) spherical planet called Earth, and we care about distances between places on its surface (say Otaniemi and Paris), let us still address the topic of metrics on the sphere briefly.

**Example V.10** (The unit sphere in  $\mathbb{R}^3$ ).

Consider the unit sphere

$$\mathbf{S}^2 = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1 \right\} \subset \mathbb{R}^3$$

in the 3-dimensional space  $\mathbb{R}^3$ . Since the Euclidean space  $\mathbb{R}^3$  is a metric space (Example V.3), the sphere  $\mathbf{S}^2$  also inherits a metric as its subset. Note, however, that in this metric inherited from  $\mathbb{R}^3$ , the distances between points on the sphere are their shortest distances in the 3-dimensional space (i.e., along straight lines) — see the gray line in Figure V.1.

This notion of distances on  $\mathbf{S}^2$  is relevant for example for neutrinos passing through the Earth (admitting that we only use neutrino detectors relatively close to the surface of the Earth).

**Example V.11** (Another metric on the unit sphere in  $\mathbb{R}^3$ ).

For the purposes of, say, aviation or nautical seafaring, the natural metric on the sphere

$$\mathbf{S}^2 = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1 \right\} \subset \mathbb{R}^3$$

would be given by shortest distances along the surface of the sphere — see the blue path in Figure V.1. The easiest precise definition of this metric is in terms of the infimum of the arclengths of paths in  $\mathbf{S}^2$  connecting the two points, see Appendix B.4, but we leave the details to the interested reader.

Already Examples V.8, V.9, V.10, and V.11 as well as Example V.7 give a lot of metric spaces which are *not* normed spaces. Among these simple examples, Examples V.11 and V.7 are not even subsets of a normed space equipped with the metric induced by the norm as in V.5. The abundance of such relevant examples shows an advantage of the more general concept of metric spaces compared to that of normed spaces.<sup>3</sup>

### V.3. First notions in metric spaces

Let us now introduce some notions that are immediately meaningful in metric spaces. Throughout, let  $(X, \mathbf{d})$  be a metric space.

#### Balls

For  $z \in X$  and  $r > 0$ , the set

$$\mathcal{B}_r(z) = \left\{ x \in X \mid \mathbf{d}(z, x) < r \right\} \quad (\text{V.9})$$

is called the **open ball** of radius  $r$  centered at  $z$ . Note that while in Euclidean spaces these are balls in an ordinary sense, in general such “balls” can be quite different. Note that by (M-0), the center always belongs to the ball,  $z \in \mathcal{B}_r(z)$  for any  $r > 0$ .

Similarly, the **closed ball** of radius  $r \geq 0$  centered at  $z \in X$  is

$$\overline{\mathcal{B}}_r(z) = \left\{ x \in X \mid \mathbf{d}(z, x) \leq r \right\}; \quad (\text{V.10})$$

the only difference to (V.9) being that also points at exactly distance  $r$  from  $z$  are included. In particular the open ball is a subset in the closed ball of the same center and radius,  $\mathcal{B}_r(z) \subset \overline{\mathcal{B}}_r(z)$ . By (M-0) we have  $z \in \overline{\mathcal{B}}_r(z)$  for any  $r \geq 0$ .

Furthermore the set

$$\left\{ x \in X \mid \mathbf{d}(z, x) = r \right\} = \overline{\mathcal{B}}_r(z) \setminus \mathcal{B}_r(z) \quad (\text{V.11})$$

of points exactly at distance  $r > 0$  from  $z$  could be called a *sphere* (compare with Example V.10).

**Example V.12** (Balls on the real line are intervals).

Consider the real line  $\mathbb{R}$  with its standard metric  $\mathbf{d}_{\mathbb{R}}$  as in (V.2). Let  $z \in \mathbb{R}$  and  $r > 0$ .

---

<sup>3</sup>Admittedly, normed spaces have their own advantages, since they have more structure than general metric spaces — in particular they have vector addition and scalar multiplication.

The open ball of radius  $r$  centered at  $z$  now consists of all those real numbers  $x$  whose distance from  $z$  is strictly less than  $r$  — it is an open interval

$$\mathcal{B}_r(z) = \left\{ x \in \mathbb{R} \mid \underbrace{d_{\mathbb{R}}(z, x)}_{|x-z| < r} < r \right\} = (z - r, z + r) \subset \mathbb{R}.$$

The closed ball of radius  $r$  centered at  $z$  consists of all those real numbers  $x$  whose distance from  $z$  is less than or equal to  $r$ ; it is a closed interval

$$\overline{\mathcal{B}}_r(z) = \left\{ x \in \mathbb{R} \mid \underbrace{d_{\mathbb{R}}(z, x)}_{|x-z| \leq r} \leq r \right\} = [z - r, z + r] \subset \mathbb{R}.$$

**Example V.13** (Balls in a function space with the uniform norm).

Consider the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on a closed interval  $[a, b]$ , equipped with the metric induced by the supremum norm as in Example V.6.

Let  $f \in \mathcal{C}([a, b])$  and  $r > 0$ . In view of the expression (V.6) for the metric on  $\mathcal{C}([a, b])$  and the definition of balls (V.9) in metric spaces, the open ball  $\mathcal{B}_r(f)$  of radius  $r$  around the function  $f: [a, b] \rightarrow \mathbb{R}$  can be visualized as consisting of all those continuous functions  $g: [a, b] \rightarrow \mathbb{R}$  whose graphs stay in a “tube” of radius  $r$  from the graph of  $f$ .

**Example V.14** (Balls with respect to the 0/1-metric).

Let  $X$  be a set and  $d_{0/1}$  the 0/1-metric (V.7) on  $X$ . Let  $z \in X$ .

Since all distinct points are at distance 1 from each other, there are no other points besides the center  $z$  itself in any open ball of radius  $r \leq 1$ , i.e., the open ball is a singleton  $\mathcal{B}_r(z) = \{z\}$  whenever  $0 < r \leq 1$ . On the other hand, all of the points in the space are within any radius  $r > 1$  from  $z$ , so the open ball is the whole space  $\mathcal{B}_r(z) = X$  whenever  $r > 1$ .

For closed balls one gets  $\overline{\mathcal{B}}_r(z) = \{z\}$  when  $0 \leq r < 1$ , and  $\overline{\mathcal{B}}_r(z) = X$  when  $r \geq 1$ .

A simple observation which is used frequently is that balls of smaller radius are contained in balls of larger radius with the same center. More precisely, for any  $z \in X$  and for  $r_1 \leq r_2$ , directly from (V.9) and (V.10) we see that  $\mathcal{B}_{r_1}(z) \subset \mathcal{B}_{r_2}(z)$  and  $\overline{\mathcal{B}}_{r_1}(z) \subset \overline{\mathcal{B}}_{r_2}(z)$ .

## Diameter and distance

For  $A \subset X$  a subset, we define the **diameter** as

$$\text{diam}(A) = \sup \left\{ d(a_1, a_2) \mid a_1, a_2 \in A \right\}. \quad (\text{V.12})$$

The idea is that we are trying to choose two points  $a_1, a_2 \in A$  from the set with maximal possible distance between them, but the maximum in general does not need to exist, so we take the supremum instead.

If  $\text{diam}(A) < \infty$ , we say that the set  $A$  is **bounded**.

**Example V.15** (Balls are bounded).

Let  $z \in X$  and  $r > 0$ . What is the diameter of  $A = \mathcal{B}_r(z)$ ?

If  $x_1, x_2 \in \mathcal{B}_r(z)$ , then by definition  $d(z, x_1) < r$  and  $d(z, x_2) < r$ . By triangle inequality (M-Δ) we get

$$d(x_1, x_2) \leq d(x_1, z) + d(z, x_2) < r + r = 2r.$$

In view of the definition (V.12) this shows that

$$\text{diam}(\mathcal{B}_r(z)) \leq 2r.$$

In particular this shows that any open ball  $\mathcal{B}_r(z)$  is bounded.

One should not, however, expect the diameter of a ball of radius  $r$  to be exactly  $2r$  in general. For example in a space with 0/1 metric, if  $0 < r \leq 1$ , we have  $\mathcal{B}_r(z) = \{z\}$  (see Example V.14) and correspondingly  $\text{diam}(\mathcal{B}_r(z)) = \text{diam}(\{z\}) = 0$ .

**Lemma V.16** (Sets contained in balls are bounded).

*If  $A \subset \mathcal{B}_r(z)$  for some  $z \in X$  and  $r > 0$ , then the set  $A$  is bounded.*

*Proof.* If  $A \subset A' \subset X$ , then from definition (V.12) it is clear that  $\text{diam}(A) \leq \text{diam}(A')$ . In particular assuming  $A \subset \mathcal{B}_r(z)$ , we get  $\text{diam}(A) \leq \text{diam}(\mathcal{B}_r(z)) \leq 2r$  by the example above. This shows that  $\text{diam}(A) < \infty$  and establishes boundedness of  $A$ .  $\square$

Also the following converse implication holds, and it is frequently useful.

**Exercise V.3** (Bounded sets are contained in some balls).

Suppose that  $z \in X$ . Show that if  $A \subset X$  is bounded, then there exists some  $r > 0$  (sufficiently large) such that  $A \subset \mathcal{B}_r(z)$ .

Hint: The case  $A = \emptyset$  is trivial (why?). If  $A \neq \emptyset$ , pick some  $a \in A$ , and choose (for example)  $r = \text{diam}(A) + d(z, a) + 1$ . Use the triangle inequality to conclude.

The **distance** between non-empty subsets  $A, B \subset X$  is defined as

$$\text{dist}(A, B) = \inf \left\{ d(a, b) \mid a \in A, b \in B \right\}. \quad (\text{V.13})$$

The idea is that we are trying to choose points  $a \in A$  and  $b \in B$  from the two sets with minimal possible distance among them, but the minimum in general does not need to exist, so we take the infimum instead.

Obviously for any  $a, b \in X$ , the distance between the corresponding singletons is just  $\text{dist}(\{a\}, \{b\}) = d(a, b)$ , so in this way the distance is a more general notion than the metric itself.<sup>4</sup>

We always have  $\text{dist}(A, B) \geq 0$  for non-empty subsets  $A, B \subset X$ , since  $d(a, b) \geq 0$  for all  $a \in A, b \in B$ . If  $A \cap B \neq \emptyset$ , then the existence of a point  $x \in A \cap B$  and the property  $d(x, x) = 0$  implies that  $\text{dist}(A, B) = 0$ . But it is possible to have vanishing distance even between two disjoint sets!

**Example V.17** (Disjoint sets at zero distance).

Consider the real line  $\mathbb{R}$  with its usual metric  $d_{\mathbb{R}}$  (Example V.2). Let  $A = (-\infty, 0] \subset \mathbb{R}$  and  $B = (0, 1]$ . The sets  $A$  and  $B$  are disjoint,  $A \cap B = \emptyset$ . We nevertheless have  $\text{dist}(A, B) = 0$  (think about the precise justification!).

The general notion of course captures also familiar problems such as the following.

**Exercise V.4** (Distance from a circle to a line).

Consider the Euclidean plane  $\mathbb{R}^2$  (Example V.3). Let

$$\begin{aligned} C &= \left\{ (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1 \right\} \subset \mathbb{R}^2, \\ L &= \left\{ (x, y) \in \mathbb{R}^2 \mid y = 2 - x \right\} \subset \mathbb{R}^2. \end{aligned}$$

---

<sup>4</sup>Despite some parallels of properties and interpretation, the distance is *not* a metric on the set  $\mathcal{P}(X)$  of all subsets of  $X$  — it is nonnegative and symmetric, but generally fails the triangle inequality and nondegeneracy, as shown by the examples right afterwards.

Calculate the distance between the circle  $C$  and the line  $L$ , and argue precisely that your answer  $\text{dist}(C, L) = \sqrt{2} - 1$  indeed is the infimum in the definition (V.13).

## V.4. Open and closed sets

The notion of an open subset in a metric space, and the related notion of a closed subset, turn out to lead to an extremely rich theory. All general topology, ultimately, is about open sets. This is probably hard to believe at first sight, but in the chapters that follow we hope you get to appreciate that this simple-looking concept is in fact very profound.

Let  $(X, d)$  be a metric space, throughout.

### Open sets

**Definition V.18** (Open set).

A subset  $U \subset X$  in a metric space  $X$  is **open** if for all  $x \in U$  there exists an  $r_x > 0$  such that  $\mathcal{B}_{r_x}(x) \subset U$ .

In other words, all points of an open set must have a little bit of room around them within the set itself: a small ball around the point is contained in the set. Given a point  $x \in X$ , we call any open set  $U \subset X$  such that  $x \in U$  a **neighborhood** of the point  $x$ . Topology can be seen (almost by its definition) as the study of neighborhoods of points.

The following two examples are trivial — but important!

**Example V.19** (The empty set is open).

The empty set  $\emptyset$  is a subset in any metric space  $X$ . It is obvious (by careful thinking<sup>5</sup>) that  $\emptyset \subset X$  is open in the sense of Definition V.18.

**Example V.20** (The whole space is open).

In any metric space  $X$ , the whole space  $X$  is a subset. It is obvious (by careful thinking<sup>6</sup>) that  $X \subset X$  is open in the sense of Definition V.18.

Ok, let us have our first nontrivial example.<sup>7</sup>

**Example V.21** (Open balls are open sets).

Suppose  $z \in X$  and  $r > 0$ . We claim that the open ball  $\mathcal{B}_r(z) \subset X$  is an open subset.

<sup>5</sup>There are no points in the empty set  $\emptyset$ , so Definition V.18 contains no requirements (there is only one requirement for every point  $x$  of the set), and in particular there is no required property that would not be satisfied! So  $\emptyset$  is open.

<sup>6</sup>Let  $x \in X$ . Take any  $r_x > 0$ , for example  $r_x = 1$ . The open ball  $\mathcal{B}_{r_x}(x)$  is *by definition* (V.9) contained in the space  $X$ , i.e.,  $\mathcal{B}_{r_x}(x) \subset X$ . In view of Definition V.18, this shows that  $X$  is open.

<sup>7</sup>If the phrase “open balls are open sets” sounds tautological, remember that we have given a precise meaning to both the term *open ball* (V.9) and the term *open set* (Definition V.18), and it is the precise definitions that carry logical content, not the words we have chosen to use... The terminology has actually been chosen *because of* the properties that we are just about to establish!

We check this directly from Definition V.18. So let  $x \in \mathcal{B}_r(z)$ . By definition (V.9) this means that  $d(z, x) < r$ . Now let  $r_x = r - d(z, x)$  and note that  $r_x > 0$  because  $d(z, x) < r$ . Now we only need to check that  $\mathcal{B}_{r_x}(x) \subset \mathcal{B}_r(z)$ . So let  $y \in \mathcal{B}_{r_x}(x)$ , which by definition means  $d(x, y) < r_x$ . Using triangle inequality (M- $\Delta$ ), we now get

$$d(z, y) \leq d(z, x) + d(x, y) < d(z, x) + r_x = d(z, x) + (r - d(z, x)) = r,$$

which shows that  $y \in \mathcal{B}_r(z)$ . Since  $y \in \mathcal{B}_{r_x}(x)$  was arbitrary, we conclude  $\mathcal{B}_{r_x}(x) \subset \mathcal{B}_r(z)$ , which is what we needed to check.

**Exercise (✓) V.5** (Punctured space is open).

Let  $z \in X$ . The complement  $X \setminus \{z\}$  of the singleton  $\{z\}$  is called a **punctured space**. Show that the punctured space  $X \setminus \{z\} \subset X$  is open.

**Exercise V.6** (The complement of a closed ball is open).

Let  $z \in X$  and  $r \geq 0$ . Show that the complement  $X \setminus \overline{\mathcal{B}}_r(z)$  of the closed ball  $\overline{\mathcal{B}}_r(z)$  is open.

The following proposition shows how we can construct new open sets out of given ones. Again despite its simplicity, it will turn out unbelievably consequential.

**Proposition V.22** (Unions and intersections of open sets).

- (a) *The union of any collection of open sets in  $X$  is also open.*
- (b) *The intersection of any finite collection of open sets in  $X$  is also open.*

Note the difference: in (a) we allow arbitrary collections of open sets, whereas in (b) we only allow finite collections of open sets! The following example shows why.

**Example V.23** (Infinite intersections of open sets are not always open).

Consider the metric space  $(\mathbb{R}, d_{\mathbb{R}})$  — the real line with its standard metric.

For  $n \in \mathbb{N}$ , the open interval  $U_n = (-\frac{1}{n}, \frac{1}{n})$  is just the open ball of radius  $\frac{1}{n}$  centered at  $0 \in \mathbb{R}$  (recall Example V.12), and therefore  $U_n \subset \mathbb{R}$  is an open set (by Example V.21). The intersection of the countably infinite collection  $(U_n)_{n \in \mathbb{N}}$  of open sets is  $\bigcap_{n \in \mathbb{N}} U_n = \{0\}$ , a singleton (the reasoning is similar to Example I.7). But the singleton  $\{0\} \subset \mathbb{R}$  is not an open set!<sup>8</sup>

This shows that the requirement of finiteness of the collection of open sets is needed in Proposition V.22(b) — the conclusion is not generally true for infinite collections (not even countably infinite).

*Proof of Proposition V.22.* We prove statements (a) and (b) separately.

*proof of (a):* Let  $(U_j)_{j \in J}$  be a collection of open sets  $U_j \subset X$ . Denote the union of this collection by

$$U = \bigcup_{j \in J} U_j.$$

We prove that  $U \subset X$  is open, directly from Definition V.18. So let  $x \in U = \bigcup_{j \in J} U_j$ . By definition of the union, there exists some  $j_0 \in J$  such that  $x \in U_{j_0}$ . Then since  $U_{j_0} \subset X$  was assumed open, there exists some  $r > 0$  such that  $\mathcal{B}_r(x) \subset U_{j_0}$ . But now

$$\mathcal{B}_r(x) \subset U_{j_0} \subset \bigcup_{j \in J} U_j = U.$$

The existence of such an  $r > 0$  for any  $x \in U$  by definition shows that  $U$  is open.

---

<sup>8</sup>Indeed, there is no  $r > 0$  such that the open ball  $\mathcal{B}_r(0)$  of radius  $r$  centered at  $0 \in \{0\}$  would not contain also other real numbers besides 0: we always have  $\mathcal{B}_r(0) \not\subset \{0\}$ .

*proof of (b):* Let  $U_1, U_2, \dots, U_m \subset X$  be open sets (finitely many!). Consider their intersection

$$V = U_1 \cap U_2 \cap \dots \cap U_m.$$

We prove that  $V \subset X$  is open, directly from Definition V.18. So let  $x \in V = \bigcap_{j=1}^m U_j$ . By definition of the intersection, for all  $j \in \{1, \dots, m\}$  we have  $x \in U_j$ . Then since each  $U_j \subset X$  was assumed open, there exists  $r_j > 0$  such that  $\mathcal{B}_{r_j}(x) \subset U_j$ . Now let  $r = \min \{r_1, \dots, r_m\}$ . We have  $r > 0$ , as the minimum of these finitely many positive numbers. For each  $j \in \{1, \dots, m\}$ , since  $r \leq r_j$ , we have  $\mathcal{B}_r(x) \subset \mathcal{B}_{r_j}(x) \subset U_j$ . By definition of intersection, this implies  $\mathcal{B}_r(x) \subset \bigcap_{j=1}^m U_j = V$ . The existence of such an  $r > 0$  for any  $x \in V$  by definition shows that  $V$  is open.  $\square$

**Example V.24** (Open sets in the 0/1-metric).

Let  $X$  be a set. Consider the metric space  $(X, d_{0/1})$  obtained by equipping  $X$  with the 0/1-metric of Example V.7.

For any  $z \in X$ , we observe based on Example V.14 that the singleton  $\{z\}$  is in fact a certain open ball: for  $r = 1$  (or in fact any  $0 < r \leq 1$ ) we have  $\{z\} = \mathcal{B}_r(z)$ . In particular it then follows from Example V.21 that the singleton  $\{z\} \subset X$  is an open set.

Now consider any subset  $A \subset X$ . Of course the subset is the union of the singletons of its points,  $A = \bigcup_{a \in A} \{a\}$ , and since each singleton  $\{a\} \subset X$  is open, this union is open by Proposition V.22(i). We conclude that *every* subset  $A \subset X$  is open in a space equipped with the 0/1-metric!

### *Isolated points and discrete topology*

Example V.24 shows some peculiar properties of the 0/1-metric.

In general, in a metric space  $(X, d)$ , if  $z \in X$  is a point such that the singleton  $\{z\}$  is open, then we say that  $z$  is an **isolated point**. The first observation in Example V.24 therefore says that in a space with the 0/1-metric, every point is an isolated point.

If in a metric space  $(X, d)$  every subset  $A \subset X$  is an open set, then the space is said to have (the) **discrete topology**. The second observation in Example V.24 therefore says that a space equipped with the 0/1-metric has discrete topology. Just like in the example, it is easy to see that a space has discrete topology if and only if its every point is an isolated point.

Despite the fact that we thus first encountered discrete topology in a somewhat exotic space, there are actually many quite familiar examples of it.

**Exercise V.7** (Finite metric spaces have discrete topology).

Suppose that  $X$  is a finite set, and  $d$  is a metric on  $X$ . Show that every point  $z \in X$  is an isolated point, and that  $(X, d)$  has discrete topology.

**Exercise V.8** (The set of integers has discrete topology).

Consider the real line  $\mathbb{R}$  with its standard metric  $d_{\mathbb{R}}$ . Since the set  $\mathbb{Z} \subset \mathbb{R}$  of integers is a subset in the metric space  $(\mathbb{R}, d_{\mathbb{R}})$ , it becomes a metric space with the metric inherited from  $\mathbb{R}$  by (V.8). Show that every point  $n \in \mathbb{Z}$  is an isolated point in  $\mathbb{Z}$ , and that  $\mathbb{Z}$  has discrete topology.

Do we have similar conclusions for  $\mathbb{N} \subset \mathbb{R}$ ?

### **Open sets in subspaces of metric spaces**

The following simple observation is the foundation of subspace topology.

**Lemma V.25** (Open sets in a subspace of a metric space).

*Let  $(X, d)$  be a metric space, and let  $X' \subset X$  be a subset, which we equip with the inherited metric  $d'$  given by (??).*

*Then a subset  $U' \subset X'$  is open if and only if there exists an open set  $U \subset X$  such that  $U' = U \cap X'$ .*

*Proof.* We prove the “if” and “only if” implications separately.

*proof of “only if”:* **UNDER CONSTRUCTION!**

*proof of “if”:* **UNDER CONSTRUCTION!** □

## Closed sets

The definition of a closed set directly refers to that of an open set (Definition V.18).

**Definition V.26** (Closed set).

A subset  $A \subset X$  in a metric space  $X$  is **closed** if its complement  $X \setminus A \subset X$  is open.

Let us again start by two trivial but important examples.

**Example V.27** (The empty set is closed).

The empty set  $\emptyset$  is a subset in any metric space  $X$ . It is closed, since its complement  $X \setminus \emptyset = X$  is the whole space, which is open by Example V.20.

**Example V.28** (The whole space is closed).

In a metric space  $X$ , the whole space is a subset  $X \subset X$ . It is closed, since its complement  $X \setminus X = \emptyset$  is the empty set, which is open by Example V.19.

The first nontrivial example is the following.

**Example V.29** (Closed balls are closed sets).

Let  $z \in X$  and  $r \geq 0$ . The closed ball  $\overline{B}_r(z) \subset X$  is closed, since by Exercise V.6 its complement  $X \setminus \overline{B}_r(z) \subset X$  is open.

**Example V.30** (Singletons are closed sets).

Let  $z \in X$ . The singleton  $\{z\} \subset X$  is closed, since by Exercise V.5 its complement  $X \setminus \{z\} \subset X$  is open.

As an exercise, you can prove the following directly from definition.

**Exercise V.9** (Finite sets are closed).

Let  $A = \{z_1, \dots, z_m\} \subset X$  be a finite subset in a metric space  $X$ . Prove that  $A \subset X$  is closed.

**Proposition V.31** (Unions and intersections of closed sets).

- (a) *The intersection of any collection of closed sets in  $X$  is also closed.*
- (b) *The union of any finite collection of closed sets in  $X$  is also closed.*



**Exercise V.10** (Proof of Proposition V.31).

Prove Proposition V.31.

Hint: Use Exercise I.6 and the corresponding result for open sets.

You can now consider Exercise V.9 again, in view of Proposition V.31 and Example V.30. The right tools make things easier.

### Warning: clopen sets!

You should *not* think of open and closed as opposites!<sup>9</sup> They are mathematical terms whose meaning was given in Definitions V.18 and V.26.

There are, in particular, sets which are both open and closed.

**Remark V.32** (Clopen sets).

According to Examples V.19 and V.27, the empty set  $\emptyset \subset X$  is both open and closed.

According to Examples V.20 and V.28, the whole space  $X \subset X$  is both open and closed.

Depending on the metric space, there can exist also other subsets which are both open and closed.<sup>10</sup> Such sets are sometimes called **clopen** (although this notion is not needed often).

In any metric space  $(X, d)$  with discrete topology<sup>11</sup>, any subset  $A \subset X$  is open (this is the definition of discrete topology!), but also its complement  $X \setminus A \subset X$  is open (it is also a subset!), so  $A$  is by definition also closed.

There are also sets which are neither open nor closed.

**Example V.33** (The set of rational numbers is neither open nor closed on the real line).

Consider the real line  $\mathbb{R}$  with its usual metric, and consider the subset  $\mathbb{Q} \subset \mathbb{R}$  of rational numbers in it. We claim that  $\mathbb{Q}$  is neither open nor closed.

To see that  $\mathbb{Q}$  is not open, take a rational number  $q \in \mathbb{Q}$ , for example  $q = 0$ . Now for an arbitrary  $r > 0$ , the open ball  $\mathcal{B}_r(q) = (q - r, q + r)$  is a non-empty open interval, and thus by Theorem II.23 contains some irrational numbers and therefore we have  $\mathcal{B}_r(q) \not\subset \mathbb{Q}$ . This shows that  $\mathbb{Q}$  is not open.

To see that  $\mathbb{Q}$  is not closed, we must by definition show that its complement  $\mathbb{R} \setminus \mathbb{Q}$  is not open. So take an irrational number  $z \in \mathbb{R} \setminus \mathbb{Q}$ , for example  $z = \sqrt{2}$ . Now for an arbitrary  $r > 0$ , the open ball  $\mathcal{B}_r(z) = (z - r, z + r)$  is a non-empty open interval, and thus by Theorem II.21 contains some rational numbers and therefore we have  $\mathcal{B}_r(z) \not\subset \mathbb{R} \setminus \mathbb{Q}$ . This shows that  $\mathbb{R} \setminus \mathbb{Q}$  is not open, i.e., that  $\mathbb{Q}$  is not closed.

**Exercise V.11** (Half-open intervals are neither open nor closed on the real line).

Let  $a < b$ . Show that  $(a, b] \subset \mathbb{R}$  is neither open nor closed.

Check that the same holds also for  $[a, b) \subset \mathbb{R}$ ,  $[a, \infty) \subset \mathbb{R}$ , and  $(-\infty, b] \subset \mathbb{R}$ .

<sup>9</sup>Sometimes this warning is summarized as “sets are not doors”. A door, arguably, is always either open or closed — whereas subsets in a metric space can be both open and closed simultaneously, or neither open or closed.

<sup>10</sup>This in fact amounts to the disconnectedness of the space — which we will define and study in Lecture XII.

<sup>11</sup>Recall that examples of spaces with discrete topology include the set  $\mathbb{Z}$  of integers, the set  $\mathbb{N}$  of natural numbers, any finite metric space  $X$ , and any set  $X$  equipped with the 0/1 metric.

To reiterate, *open* and *closed* mean exactly what they were defined to mean. They are *not* the opposites of each other!

### V.5. ♥ Interior, exterior, and boundary

The following notions are frequently used in calculus, and various areas of analysis and its applications. They are, however, not strictly necessary for the theory covered in this course. The definitions below should be easy to grasp once one has a feeling for what metric spaces are, but if the amount of definitions up until now in this chapter seems overwhelming already, then one can postpone studying these concepts until they are needed in other courses.

Given a subset  $A$  of a metric space  $X$ , an often useful idea is to classify the points  $x$  of the space  $X$  according to how they lie in relation to the subset  $A$ : whether they are well inside, well outside, or neither really. The precise definitions and terminology are as follows.

**Definition V.34** (Interior and exterior).

A point  $x \in X$  is said to be an **interior point** of  $A \subset X$  if for some  $r > 0$  we have  $\mathcal{B}_r(x) \subset A$ . A point  $x$  is said to be an **exterior point** of  $A$  if for some  $r > 0$  we have  $\mathcal{B}_r(x) \subset X \setminus A$ . The set of all interior points of  $A$  is denoted by  $A^\circ$ , and the set of all exterior points by  $\text{ext}(A)$ .

From the definition it is clear that  $A^\circ \subset A$  and  $\text{ext}(A) \subset X \setminus A$ . In particular the interior and the exterior are mutually disjoint,  $A^\circ \cap \text{ext}(A) = \emptyset$ .

**Definition V.35** (Boundary).

The **boundary** of subset  $A \subset X$  is defined as

$$\partial A = X \setminus (A^\circ \cup \text{ext}(A))$$

The points  $x \in \partial A$  are called **boundary points** of  $A \subset X$ .

In other words, a point  $x$  is a boundary point of  $A$  if it is neither an interior point nor an exterior point. The following characterization is therefore obvious (consider the contrapositive).

**Lemma V.36** (A direct characterization of boundary points).

A point  $x \in X$  is a boundary point of  $A \subset X$  if and only if for all  $r > 0$  the ball  $\mathcal{B}_r(x)$  contains both a point of  $A$  and a point of its complement, i.e.,  $\mathcal{B}_r(x) \cap A \neq \emptyset$  and  $\mathcal{B}_r(x) \cap (X \setminus A) \neq \emptyset$ .

In summary we have defined the interior, the exterior, and the boundary of  $A$  so that they form a partition

$$X = A^\circ \cup \text{ext}(A) \cup \partial A.$$

of the space to three mutually disjoint subsets.

*Examples***Example V.37** (The interior of an interval).**UNDER CONSTRUCTION!****Example V.38** (The boundary of an interval).**UNDER CONSTRUCTION!****Proposition V.39** (The interior and exterior are open).*Let  $A \subset X$ . Then both  $A^\circ$  and  $\text{ext}(A)$  are open sets in  $X$ .**Proof.* **UNDER CONSTRUCTION!**

□

**Corollary V.40** (The boundary is closed).*Let  $A \subset X$ . Then the boundary  $\bar{A}$  is a closed set in  $X$ .**Proof.* By Proposition V.39, both  $A^\circ$  and  $\text{ext}(A)$  are open, so their union  $A^\circ \cup \text{ext}(A)$  is also open by Proposition V.22(a). The boundary  $\partial A = X \setminus (A^\circ \cup \text{ext}(A))$  is by definition the complement of this open set, so it is closed (by definition). □**V.6. ♥ Closure**

At this stage we will define the closure of a subset in terms of the above notions.

**Definition V.41** (Closure).The **closure** of a subset  $A \subset X$  is  $\bar{A} = A \cup \partial A$ .**UNDER CONSTRUCTION!**



## Lecture VI

### Continuity of functions

Throughout this section, let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces. We will consider functions

$$f: X \rightarrow Y.$$

Our main task is to extend the notion of continuity, which is familiar in simpler setups (for example real valued functions on the real line), to this general setup. The general definition, an “ $\varepsilon$ - $\delta$  condition”, intuitively requires that if one does not change the argument  $x$  of a continuous function  $f$  by too much, then the value  $f(x)$  of the function does not change by too much either.

It is worth pointing out this “ $\varepsilon$ - $\delta$  condition” that is taken as the general definition of continuity does not immediately reduce to the definition of continuity that was used in the special case of real functions in Chapter III. However, in this and subsequent lectures we will also obtain a number of equivalent characterizations of the general notion of continuity, and one such characterization in Chapter VIII will in particular show that the general definition indeed is logically equivalent, in the special case of real functions, to the definition used before.

Continuous functions are arguably the most important topological notion: they are in a sense exactly those functions which “respect the topology”.<sup>1</sup> In later chapters we will study what happens to various topological properties of spaces when continuous functions are applied: we will show, e.g., that the image of a connected space under a continuous function is also connected (Chapter XII), that the image of a compact space under a continuous function is also compact (Chapter XII), etc. But let us not get ahead of ourselves. We must first define the notion of continuity! The definition will be followed by the first simple examples of continuous functions. With the notion of Lipschitz-continuity, we start to get more substantial examples in many of the spaces that have been introduced in earlier chapters. And to finish this general chapter about continuity, we provide an equivalent characterization of continuity

---

<sup>1</sup>To provide some context by way of analogies and points of comparison, let us make the following meta-mathematical observation. Whenever some new mathematical structure is defined, it becomes natural to ask what are the mappings (=functions) which either “respect” the structure or “(fully) preserve” the structure. An example from linear algebra would be vector spaces: the functions that respect the vector space structure are linear maps, and functions which preserve the vector space structure are bijective linear maps. An example from algebra would be groups: the functions that respect the group structure are group homomorphisms, and functions which preserve the group structure are group isomorphisms. An example from differential geometry is that of a smooth manifold: the mappings that respect the manifold structure are smooth functions, and mappings which preserve the manifold structure are diffeomorphisms. The reader should become attentive to such examples in whatever topic of mathematics (s)he studies. For the present course, although we focus on the more concrete case of metric spaces, the core notion is topological spaces. The mappings which respect this topological structure, then, are continuous functions. In the next lecture we will study mappings which (fully) preserve the topological structure: these are called homeomorphisms.

phrased purely in terms of open sets. The characterization is more practical for certain purposes (although it looks more abstract than the “ $\varepsilon$ - $\delta$  condition”), and fundamentally demonstrates that continuity is a genuinely topological concept.

### VI.1. Definition of continuity

The following general definition of continuity makes sense whenever both the domain  $X$  and the codomain  $Y$  of a function are equipped with metrics, denoted  $\mathbf{d}_X$  and  $\mathbf{d}_Y$ , respectively.

**Definition VI.1** (Continuity).

Let  $f: X \rightarrow Y$  be a function, and  $x \in X$  a point. The function  $f$  is said to be **continuous at  $x$**  if for any  $\varepsilon > 0$  there exists a  $\delta > 0$  such that we have  $\mathbf{d}_Y(f(x'), f(x)) < \varepsilon$  whenever  $x' \in X$  and  $\mathbf{d}_X(x', x) < \delta$ .

We say that  $f: X \rightarrow Y$  is a **continuous function** if  $f$  is continuous at every point  $x \in X$ .

Here you should really care about very small numbers  $\varepsilon > 0$  and  $\delta > 0$ . Indeed, in more intuitive terms, the definition of continuity at  $x$  should be thought of as follows. However small positive amount  $\varepsilon > 0$  we are willing to allow the values of our function  $f$  to move, movement by less than this much can be guaranteed by requiring that the arguments move by less than some small amount  $\delta > 0$ . A crucial thing to note is that  $\delta$  is allowed to depend on  $\varepsilon$  (as well as on the point  $x$  where continuity is considered).<sup>2</sup> The less we are willing to let the values move, the less we can allow the arguments to move — i.e., the smaller the given  $\varepsilon > 0$ , the smaller we should choose  $\delta$ .

**Remark VI.2** (Rewriting the  $\varepsilon$ - $\delta$  condition in logically equivalent forms).

We may rewrite the above definitions in logical notation as follows:

$f$  is continuous at  $x$  means

$$\forall \varepsilon > 0 : \exists \delta > 0 : \forall x' \in X : \quad \mathbf{d}_X(x', x) < \delta \Rightarrow \mathbf{d}_Y(f(x'), f(x)) < \varepsilon, \quad (\text{VI.1})$$

and

$f$  is a continuous function means

$$\forall x \in X : \forall \varepsilon > 0 : \exists \delta > 0 : \forall x' \in X : \mathbf{d}_X(x', x) < \delta \Rightarrow \mathbf{d}_Y(f(x'), f(x)) < \varepsilon. \quad (\text{VI.2})$$

This may be further rewritten in various useful ways using the notion of balls, and the images of balls in  $X$  or the preimages of balls in  $Y$ . Specifically, we observe<sup>3</sup> that (VI.1) is

<sup>2</sup>Just note the order of quantifiers: for all  $\varepsilon > 0$  there exists a  $\delta > 0$  (which may depend on the earlier chosen  $\varepsilon$ ) such that...

<sup>3</sup>The equivalence of (VI.1) and (VI.1a) is seen by noting that by the definition of open balls,  $\mathbf{d}_X(x', x) < \delta$  is equivalent to  $x' \in \mathcal{B}_\delta(x)$ , while  $\mathbf{d}_Y(f(x'), f(x)) < \varepsilon$  is equivalent to  $f(x') \in \mathcal{B}_\varepsilon(f(x))$ . The equivalence with (VI.1b) and (VI.1c) is then seen by noting that the implication  $x' \in \mathcal{B}_\delta(x) \Rightarrow f(x') \in \mathcal{B}_\varepsilon(f(x))$  is equivalent to  $f[\mathcal{B}_\delta(x)] \subset \mathcal{B}_\varepsilon(f(x))$  by definition of images, and this in turn is equivalent to  $\mathcal{B}_\delta(x) \subset f^{-1}[\mathcal{B}_\varepsilon(f(x))]$  by definition of preimages.

equivalent to any of the following:

$$\forall \varepsilon > 0 : \exists \delta > 0 : \quad x' \in \mathcal{B}_\delta(x) \Rightarrow f(x') \in \mathcal{B}_\varepsilon(f(x)), \quad (\text{VI.1a})$$

$$\forall \varepsilon > 0 : \exists \delta > 0 : \quad f[\mathcal{B}_\delta(x)] \subset \mathcal{B}_\varepsilon(f(x)), \quad (\text{VI.1b})$$

$$\forall \varepsilon > 0 : \exists \delta > 0 : \quad \mathcal{B}_\delta(x) \subset f^{-1}[\mathcal{B}_\varepsilon(f(x))]. \quad (\text{VI.1c})$$

Similarly, of course, by simply preceding each of these with the quantifier  $\forall x \in X$ , we obtain equivalent ways to phrase the condition (VI.2).

At this point, let us give just two extremely simple examples, without even full details (the details are good exercises, and we will anyway revisit these simple examples more carefully with a later characterization of continuity).

**Example VI.3** (Constant function is continuous).

Let  $X, Y$  be any metric spaces, and  $c \in Y$ . Consider the constant function  $f: X \rightarrow Y$  given by

$$f(x) = c \quad \text{for all } x \in X.$$

We claim that  $f$  is continuous. Indeed, we leave it for the reader to verify more formally that at any point  $x$ , for any given  $\varepsilon > 0$ , we can in fact use any  $\delta > 0$  (for example  $\delta = 1$  works) — the values of a constant function in fact never move at all, however much the argument is moved.

**Example VI.4** (Identity function is continuous).

Let  $X$  be any metric space. Consider the identity function  $\text{id}: X \rightarrow X$  given by

$$\text{id}(x) = x \quad \text{for } x \in X.$$

We claim that  $\text{id}$  is continuous. Indeed, we leave it for the reader to verify more formally that at any point  $x$ , for any given  $\varepsilon > 0$ , we can in fact use exactly  $\delta = \varepsilon > 0$  — the values of the identity function move by only (and exactly) as much as the argument is moved.

## Lipschitz continuity

The general  $\varepsilon$ - $\delta$  definition of continuity (Definition VI.1) is qualitative. It does not quantitatively say exactly how much of movement of the argument corresponds to how much of movement of the values (it merely asserts that the latter can be made arbitrarily small by choosing the former sufficiently small). There are various quantitative notions of continuity, which are of practical importance especially in analysis, and which also serve as convenient sufficient conditions by which we can often verify continuity more concretely. The simplest and most important of these is Lipschitz-continuity.

**Definition VI.5** (Lipschitz property).

Let  $M \geq 0$ . A function  $f: X \rightarrow Y$  is said to be  **$M$ -Lipschitz** if for all  $x_1, x_2 \in X$  we have

$$d_Y(f(x_1), f(x_2)) \leq M d_X(x_1, x_2). \quad (\text{VI.3})$$

The parameter  $M$  in the above definition is often called the **Lipschitz constant**. Equation (VI.3) has the interpretation that the mapping  $f$  cannot stretch distances by more than a factor  $M$ .

Let us prove that the Lipschitz property above is a sufficient condition for continuity (justifying the term Lipschitz continuity).

**Proposition VI.6** (Lipschitz functions are continuous).

*If  $f: X \rightarrow Y$  is  $M$ -Lipschitz, then  $f$  is continuous.*

*Proof.* Assume that  $f$  is  $M$ -Lipschitz, and without loss of generality assume that  $M > 0$ .<sup>4</sup> To show continuity, we must verify the condition in Definition VI.1. Let therefore  $x \in X$ , and let  $\varepsilon > 0$  be given. We claim that the choice  $\delta = \frac{\varepsilon}{M} > 0$  works. Indeed, if  $d_X(x', x) < \delta = \frac{\varepsilon}{M}$ , then by the Lipschitz property (VI.3) we get

$$d_Y(f(x'), f(x)) \leq M d_X(x', x) < M \delta = M \frac{\varepsilon}{M} = \varepsilon.$$

Since this worked for an arbitrary  $\varepsilon > 0$ , continuity of  $f$  at  $x$  is established. Since the point  $x \in X$  was arbitrary, the continuity of the function  $f: X \rightarrow Y$  is established.  $\square$

### *Real functions with bounded derivative*

For real functions, the Lipschitz property is closely related to the boundedness of the derivative. Real functions with bounded derivative constitute a common and important class of Lipschitz functions.

**Example VI.7** (Functions with bounded derivative are Lipschitz).

Let  $I \subset \mathbb{R}$  be an interval and  $f: I \rightarrow \mathbb{R}$  a continuously differentiable function such that  $|f'(x)| \leq M$  for all  $x \in I$ . If  $x_1, x_2 \in I$  with  $x_1 < x_2$ , then the Mean Value Theorem states that for some  $x \in (x_1, x_2) \subset I$  we have

$$f'(x) = \frac{f(x_2) - f(x_1)}{x_2 - x_1}.$$

We may rearrange this to the form  $f(x_2) - f(x_1) = f'(x)(x_2 - x_1)$ , and then take absolute values and use the bound  $|f'(x)| \leq M$  for the derivative to get

$$|f(x_2) - f(x_1)| = |f'(x)| |x_2 - x_1| \leq M |x_2 - x_1|.$$

The metric on  $\mathbb{R}$  is taken to be the usual one, so  $|f(x_2) - f(x_1)| = d_{\mathbb{R}}(f(x_1), f(x_2))$ , and the interval  $I \subset \mathbb{R}$  inherits a metric from the real axis so that  $|x_2 - x_1| = d_I(x_1, x_2)$ . Thus the conclusion is exactly of the form (VI.3)

$$d_{\mathbb{R}}(f(x_1), f(x_2)) \leq M d_I(x_1, x_2),$$

and it is easy to see that this inequality also holds with  $x_1 > x_2$  (just rename the variables) and with  $x_1 = x_2$  (both sides are zero). Therefore  $f$  is  $M$ -Lipschitz.

An important particular case of the above is the following.

**Example VI.8** (Continuously differentiable functions on closed intervals are Lipschitz).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and  $f: [a, b] \rightarrow \mathbb{R}$  a continuously differentiable function. Then the continuous function  $f': [a, b] \rightarrow \mathbb{R}$  on the closed interval  $[a, b]$  is bounded (according to Theorem III.13), so there exists some  $M \geq 0$  such that  $|f'(x)| \leq M$  for all  $x \in [a, b]$ , and the reasoning in Example VI.7 applies. Therefore any continuously differentiable function on a closed interval is automatically Lipschitz (with some Lipschitz constant  $M$ ).

The assumption that the interval is closed was important in the previous example.

<sup>4</sup>The case  $M = 0$  is trivial — it requires that the function is constant. In any case if a function is  $M$ -Lipschitz for some  $M$ , then it is also  $M'$ -Lipschitz for any  $M' > M$ , so we may assume the Lipschitz constant chosen positive.



**Exercise VI.1** (A continuously differentiable function on an interval).

Consider the function  $f: (-1, 1) \rightarrow \mathbb{R}$  defined by

$$f(x) = \frac{1}{1-x^2} \quad \text{for } x \in (-1, 1).$$

- (a) Is the function  $f: (-1, 1) \rightarrow \mathbb{R}$  continuous?
- (b) Calculate the derivative  $f'(x)$ . Is  $f': (-1, 1) \rightarrow \mathbb{R}$  continuous?
- (c) Is the function  $f: (-1, 1) \rightarrow \mathbb{R}$  Lipschitz continuous (for some Lipschitz constant  $M$ )?

*Hint:* For the purposes of parts (a)-(b) of this exercise, let us admit that the continuity of  $f$  and  $f'$  can be understood in the sense of Definition III.7. Later in Theorem ?? it will be justified that this is indeed equivalent to the  $\varepsilon$ - $\delta$  definition of continuity.

**Exercise VI.2** (The square root function is not Lipschitz).

Prove that the function

$$q: [0, 1] \rightarrow [0, 1] \quad q(x) = \sqrt{x} \quad \text{for } x \in [0, 1]$$

is not  $K$ -Lipschitz for any  $K \geq 0$  (although it is continuous).

Why can't we use Example VI.8 in this case?

### More examples of Lipschitz functions

Let us give some examples of Lipschitz functions, where the space  $X$  on which the functions are defined is not just the real axis or an interval.

In the following example, the domain  $X$  of our function is a function space. So we are considering functions of functions (albeit still quite simple ones, more complicated examples will follow).

**Example VI.9** (The evaluation map on a function space).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and, as in Example IV.27, consider the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on this interval equipped with the supremum-norm  $\|\cdot\|_\infty$  and the metric induced by this norm.

Fix a point  $z \in [a, b]$ . Then we may define a real-valued function

$$\text{ev}_z: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$$

on this function space by the formula

$$\text{ev}_z(f) = f(z) \quad \text{for } f \in \mathcal{C}([a, b]),$$

i.e., the value of the function  $\text{ev}_z$  at the point  $f$  (note that points of a function space are themselves functions) is the value of  $f$  at the chosen point  $z \in [a, b]$ . This function  $\text{ev}_z$  is called **evaluation at point  $z$** .

We claim that  $\text{ev}_z: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  is continuous, in fact Lipschitz continuous with constant  $M = 1$ . To see this, let  $f_1, f_2 \in \mathcal{C}([a, b])$  be two points in the function space — thus themselves functions  $f_1, f_2: [a, b] \rightarrow \mathbb{R}$ . On the codomain  $\mathbb{R}$  of the function  $\text{ev}_z$  we use the usual metric, so the distance of the values is

$$d_{\mathbb{R}}(\text{ev}_z(f_1), \text{ev}_z(f_2)) = |\text{ev}_z(f_2) - \text{ev}_z(f_1)| = |f_2(z) - f_1(z)|.$$

By the vector space structure on  $\mathcal{C}([a, b])$  we have  $f_2(z) - f_1(z) = (f_2 - f_1)(z)$ , the value of the function  $f_2 - f_1 \in \mathcal{C}([a, b])$  at  $z$ . The distance of the values can then be estimated as

$$\begin{aligned} d_{\mathbb{R}}(\text{ev}_z(f_1), \text{ev}_z(f_2)) &= |(f_2 - f_1)(z)| \\ &\leq \sup_{x \in [a, b]} |(f_2 - f_1)(x)| = \|f_2 - f_1\|_\infty = d_{\mathcal{C}([a, b])}(f_1, f_2). \end{aligned}$$

This is the Lipschitz property of the function  $\text{ev}_z: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  with Lipschitz constant  $M = 1$ . In particular it follows from Proposition VI.6 that evaluation  $\text{ev}_z$  is a continuous function.

The next exercise features another function on the same function space.

**Exercise VI.3** (The maximum map on a function space).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and consider again the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on this interval equipped with the supremum-norm  $\|\cdot\|_\infty$ . For any  $f \in \mathcal{C}([a, b])$ , the maximum  $\max_{x \in [a, b]} f(x)$  exists, and we can thus define a function

$$\text{Max}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$$

defined on this function space by the formula

$$\text{Max}(f) := \max_{x \in [a, b]} f(x) \quad \text{for } f \in \mathcal{C}([a, b]).$$

Show that  $\text{Max}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  is 1-Lipschitz, and conclude that it is in particular continuous.

In the following, the domain  $X$  is an inner product space (with its natural metric).

**Example VI.10** (Inner product is continuous in either of its arguments).

Let  $V$  be an inner product space, with inner product denoted by  $\langle \cdot, \cdot \rangle$ . Fix a vector  $w \in V$ . Define a function

$$f: V \rightarrow \mathbb{R}$$

by the formula

$$f(v) = \langle w, v \rangle \quad \text{for } v \in V.$$

We claim that this function  $f: V \rightarrow \mathbb{R}$  is Lipschitz continuous with  $M = \|w\|$ .

Recall first that the domain  $V$  of the above function  $f$  is naturally a metric space with the metric  $d_V(u, v) = \|u - v\| = \sqrt{\langle u - v, u - v \rangle}$  (combine Remarks IV.24 and V.5). Also the codomain  $\mathbb{R}$  of the function  $f$  is naturally a metric space, with the usual metric  $d_{\mathbb{R}}(x, y) = |x - y|$  of the real axis. Now for  $v_1, v_2 \in V$ , calculate the quantity in the definition of Lipschitz property and estimate it as follows

$$\begin{aligned} d_{\mathbb{R}}(f(v_1), f(v_2)) &= |f(v_1) - f(v_2)| && \text{(definition of } d_{\mathbb{R}}) \\ &= |\langle w, v_1 \rangle - \langle w, v_2 \rangle| && \text{(defining formula for } f) \\ &= |\langle w, v_1 - v_2 \rangle| && \text{(linearity of inner product)} \\ &\leq \|w\| \|v_1 - v_2\| && \text{(Cauchy-Schwarz inequality)} \\ &\leq \|w\| d_V(v_1, v_2). && \text{(definition of } d_V) \end{aligned}$$

This is precisely the Lipschitz property  $d_{\mathbb{R}}(f(v_1), f(v_2)) \leq M d_V(v_1, v_2)$  with Lipschitz constant  $M = \|w\|$ .

Since  $v \mapsto \langle w, v \rangle$  has the Lipschitz property, it is continuous by Proposition VI.6. This shows that the inner product is a continuous function of its second argument. But by symmetry of inner product,  $\langle w, v \rangle = \langle v, w \rangle$ , also  $v \mapsto \langle v, w \rangle$  is continuous. Thus the inner product is continuous in either of its arguments.<sup>5</sup>

One special case (albeit one that may seem a bit abstract at first) of Example VI.10 in a function space is the following. Again the domain is a certain function space, so we are considering functions of functions.

**Example VI.11** (♥ Fourier coefficients of a function depend continuously on the function).

The Fourier series coefficients of a function  $f \in \mathcal{C}([-\pi, \pi])$  are up to a normalization constant

<sup>5</sup>The stronger statement that the inner product is “jointly continuous” in its two arguments is also true, i.e., the function  $V \times V \rightarrow \mathbb{R}$  defined by  $(v, w) \mapsto \langle w, v \rangle$  is continuous. But for this, we would first need to understand the product space  $V \times V$  as a metric space (which will have to wait until Section VII.4).

the  $L^2$ -inner products of the function  $f$  with the corresponding trigonometric basis functions of Exercise IV.12.<sup>6</sup> Therefore, applying Example VI.10, it follows that each Fourier coefficient is a continuous function of the function  $f$ , when the space of functions  $\mathcal{C}([-\pi, \pi])$  is equipped with the metric induced by the norm induced by the  $L^2$ -inner product (see Example IV.13).

Since there are (countably) infinitely many basis functions, and corresponding Fourier series coefficients for each basis function, this example featured (countably) infinitely many continuous functions of continuous functions.

Also the next exercise deals with functions of functions, but with a norm that does not come from an inner product.

**Exercise (#) VI.4** (The integral map on a function space).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and consider again the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on this interval equipped with the supremum-norm  $\|\cdot\|_\infty$ . For any  $f \in \mathcal{C}([a, b])$ , the integral  $\int_a^b f(x) dx$  is well-defined. We can thus define a function

$$\text{Int}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$$

defined on this function space by the formula

$$\text{Int}(f) := \int_a^b f(x) dx \quad \text{for } f \in \mathcal{C}([a, b]).$$

Show that  $\text{Int}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  is  $M$ -Lipschitz for  $M = b - a$ , and conclude that it is in particular continuous.

Hint: In this exercise, you may need the triangle inequality and monotonicity for integrals. The triangle inequality for integrals states that

$$\left| \int_\alpha^\beta g(t) dt \right| \leq \int_\alpha^\beta |g(t)| dt$$

whenever  $g: [\alpha, \beta] \rightarrow \mathbb{R}$  is a Riemann-integrable function (in particular if  $g$  is continuous). Monotonicity of integrals states that if  $g, h: [\alpha, \beta] \rightarrow \mathbb{R}$  are Riemann-integrable functions such that  $g(t) \leq h(t)$  for all  $t \in [\alpha, \beta]$ , then

$$\int_\alpha^\beta g(t) dt \leq \int_\alpha^\beta h(t) dt.$$

**Exercise (#) VI.5** (Revisiting functions on functions spaces).

Consider again the space  $\mathcal{C}([a, b])$ , but now equipped with the norm  $\|\cdot\|_1$  and the metric induced by it. The functions  $\text{ev}_z, \text{Max}, \text{Int}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  of the Example VI.9, Exercise VI.3, and Exercise VI.4 are still well-defined. One of these three is still Lipschitz continuous with the new choice of metric: which one?

(Extra: Why are the other two not continuous?)

**Exercise (#) VI.6** (Re-revisiting functions on functions spaces).

Consider again the space  $\mathcal{C}([a, b])$ , but now equipped with the norm  $\|\cdot\|_2$  and the metric induced by it. The functions  $\text{ev}_z, \text{Max}, \text{Int}: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  of the Example VI.9, Exercise VI.3, and Exercise VI.4 are still well-defined. One of these three is still Lipschitz continuous with the new choice of metric: which one?

(Extra: Why are the other two not continuous?)

Hint: Recall Example VI.10.

---

<sup>6</sup>The same exercise also contains the key idea for *why* the Fourier coefficients are defined as such inner products.

*Distance to a point*

On any metric space  $(X, \mathbf{d})$ , distances (defined by the metric  $\mathbf{d}$ ) give rise to natural functions. We next address the continuity of these. The key lemma is the following lower bound counterpart of the triangle inequality (M- $\Delta$ ).

**Lemma VI.12** (Triangle inequality lower bound).

Let  $(X, \mathbf{d})$  be a metric space, and  $x, y, z \in X$  any three points. Then we have

$$\mathbf{d}(x, z) \geq |\mathbf{d}(x, y) - \mathbf{d}(y, z)|. \quad (\text{VI.4})$$

*Proof.* Note that  $|\mathbf{d}(x, y) - \mathbf{d}(y, z)|$  equals either  $\mathbf{d}(x, y) - \mathbf{d}(y, z)$  or  $\mathbf{d}(y, z) - \mathbf{d}(x, y)$ , depending on the sign of the quantity inside the absolute value. We will separately prove both

$$\mathbf{d}(x, y) - \mathbf{d}(y, z) \leq \mathbf{d}(x, z) \quad \text{and} \quad (\text{VI.5})$$

$$\mathbf{d}(y, z) - \mathbf{d}(x, y) \leq \mathbf{d}(x, z). \quad (\text{VI.6})$$

Together these will show the asserted inequality  $|\mathbf{d}(x, y) - \mathbf{d}(y, z)| \leq \mathbf{d}(x, z)$ .

First apply the triangle inequality (M- $\Delta$ ) to the three points  $x, y, z \in X$  in the form

$$\mathbf{d}(x, y) \leq \mathbf{d}(x, z) + \mathbf{d}(z, y) \quad (\text{“going via } z \text{ is not a shortcut from } x \text{ to } y\text{”}).$$

Subtracting  $\mathbf{d}(z, y)$  from both sides yields  $\mathbf{d}(x, y) - \mathbf{d}(y, z) \leq \mathbf{d}(x, z)$ . Since  $\mathbf{d}(z, y) = \mathbf{d}(y, z)$  by (M-s), this establishes (VI.5) — the first of the two inequalities we set out to prove.

Then apply the triangle inequality (M- $\Delta$ ) to the three points  $x, y, z \in X$  in the form

$$\mathbf{d}(y, z) \leq \mathbf{d}(y, x) + \mathbf{d}(x, z) \quad (\text{“going via } x \text{ is not a shortcut from } y \text{ to } z\text{”}).$$

Subtracting  $\mathbf{d}(y, x)$  from both sides yields  $\mathbf{d}(y, z) - \mathbf{d}(y, x) \leq \mathbf{d}(x, z)$ . Since  $\mathbf{d}(y, x) = \mathbf{d}(x, y)$  by (M-s), this establishes (VI.6) — the second of the two inequalities we set out to prove.

The proof is thus complete.  $\square$

**Example VI.13** (Distances to a given point are 1-Lipschitz).

Let  $(X, \mathbf{d})$  be a metric space and let  $a \in X$  be a point in the space. Consider the function

$$f: X \rightarrow [0, \infty)$$

defined by the distance to the given point  $a$ ,

$$f(x) = \mathbf{d}(x, a) \quad \text{for } x \in X.$$

We claim that this function is 1-Lipschitz, and therefore in particular a continuous function  $f: X \rightarrow [0, \infty)$ .

On the codomain  $[0, \infty) \subset \mathbb{R}$  of this function we use the metric inherited from the real axis, so for any two points  $x_1, x_2 \in X$  of the domain, the distance between the corresponding values is

$$\mathbf{d}_{\mathbb{R}}(f(x_1), f(x_2)) = |f(x_2) - f(x_1)| = |\mathbf{d}(x_2, a) - \mathbf{d}(x_1, a)| \leq \mathbf{d}(x_1, x_2)$$

where the last inequality follows from (VI.4) with  $x = x_2$ ,  $y = a$ ,  $z = x_1$ . This shows that  $x \mapsto \mathbf{d}(x, z)$  is 1-Lipschitz  $X \rightarrow [0, \infty)$ .

**Exercise (✓) VI.7** (The norm is a continuous function).

Let  $(V, \|\cdot\|)$  be a normed space. Consider the norm, seen as a function

$$v \mapsto \|v\|.$$

Why is this function  $V \rightarrow [0, \infty)$  continuous?

## ♥ Other quantitative notions of continuity

UNDER CONSTRUCTION!

## VI.2. Characterization of continuity

It turns out that the  $\varepsilon$ - $\delta$  definition of continuity (Definition VI.1) is equivalent to a condition formulated purely in terms of open sets. Let us just briefly mention why this is significant, although we will not elaborate further now. It means that the notion of continuity of a function only depends on the metrics used on its domain and codomain only to the extent that these metrics affect which sets are open. Properties that can be phrased purely in terms of open sets are topological. Namely, in general topology the notion of open sets is taken as the axiomatic starting point. What we below see as a characterization of continuity between metric spaces is then taken as the definition of continuity between topological spaces (the  $\varepsilon$ - $\delta$  definition of course could not be used in the absence of metrics).

Besides thus enabling (further) generalization of the notion of continuity, the characterization below has concrete benefits as well. It occasionally serves as an easier way to verify that a given function is continuous (for an example of this, see Proposition VI.20). Even more commonly it is used with functions that are known to be continuous, to give quick arguments that certain sets defined in terms of those functions are open (or closed). We will provide examples of such reasoning soon, and almost every lecture from here on contains further examples.

Note that the following characterizes the continuity of a function, rather than just the continuity of a function at a given point.

**Theorem VI.14** (Characterization of continuity with open (and closed) sets).

*Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f: X \rightarrow Y$  a function. Then the following are equivalent:*

- (i) *the function  $f: X \rightarrow Y$  is continuous (in the sense of Definition VI.1),*
- (ii) *for every open set  $V \subset Y$ , the preimage  $f^{-1}[V] \subset X$  is open,*
- (iii) *for every closed set  $A \subset Y$ , the preimage  $f^{-1}[A] \subset X$  is closed.*

*Proof.* We will show the equivalence of all three conditions by separately proving the implications (i)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (i), and the equivalence (ii)  $\Leftrightarrow$  (iii).

*proof of (i)  $\Rightarrow$  (ii):* Suppose that  $f: X \rightarrow Y$  is continuous. Let  $V \subset Y$  be an open set. We must show that the preimage  $f^{-1}[V] \subset X$  is open, i.e., we must find neighborhoods for its points contained in the set itself. So let  $x \in f^{-1}[V]$  be a point. By definition of preimages, this means  $f(x) \in V$ . By openness of  $V$  there exists some  $\varepsilon > 0$  such that  $\mathcal{B}_\varepsilon(f(x)) \subset V$ . According to the reformulation (VI.1c) of continuity of  $f$  at the point  $x$ , there then exists a  $\delta > 0$  such that  $\mathcal{B}_\delta(x) \subset f^{-1}[\mathcal{B}_\varepsilon(f(x))] \subset f^{-1}[V]$ . Thus  $\mathcal{B}_\delta(x)$  is a neighborhood of  $x$  contained in  $f^{-1}[V]$ . We conclude that  $f^{-1}[V] \subset X$  is open.

*proof of (ii)  $\Rightarrow$  (i):* Suppose that the preimage  $f^{-1}[V]$  of each open set  $V \subset Y$  is open. We must show that  $f$  is continuous. So let  $x \in X$ , and let  $\varepsilon > 0$ . The open ball  $\mathcal{B}_\varepsilon(f(x)) \subset Y$  is an open set, so by assumption its preimage  $f^{-1}[\mathcal{B}_\varepsilon(f(x))] \subset X$  is also open. Note that we have  $x \in f^{-1}[\mathcal{B}_\varepsilon(f(x))]$ , since  $f(x) \in \mathcal{B}_\varepsilon(f(x))$ . Therefore by openness of  $f^{-1}[\mathcal{B}_\varepsilon(f(x))]$ , there exists some  $\delta > 0$  such that  $\mathcal{B}_\delta(x) \subset f^{-1}[\mathcal{B}_\varepsilon(f(x))]$ . According to the reformulation (VI.1c), we have thus shown the continuity of  $f$  at the point  $x$ . The point  $x \in X$  was arbitrary, so the continuity of  $f: X \rightarrow Y$  follows.

*proof of (ii)  $\Leftrightarrow$  (iii):* The equivalence of (ii) and (iii) is easy to see by passing to complements.

For example, assume that (ii) holds, i.e., that the preimage  $f^{-1}[V]$  of every open set  $V \subset Y$  is open. We must then show that the preimage of any closed set in  $Y$  is closed. So let

$A \subset Y$  be closed. To show that its preimage  $f^{-1}[A] \subset X$  is closed, we must by definition show that its complement  $X \setminus f^{-1}[A]$  is open. But by routine set theory, the complement of the preimage is the preimage of the complement,  $X \setminus f^{-1}[A] = f^{-1}[Y \setminus A]$ . However, the complement  $Y \setminus A$  of the closed set  $A$  is by definition open. The preimage  $f^{-1}[Y \setminus A]$  of this open set is by assumption (ii) open. We have thus shown that the preimage  $f^{-1}[A]$  of an arbitrary closed set  $A \subset Y$  is closed.

The other direction of this equivalence is entirely similar.  $\square$

**Remark VI.15** (The images of open sets under continuous functions are not always open).  
**UNDER CONSTRUCTION!**

To familiarize ourselves with this characterization, let us revisit some very simple examples of continuous functions in view of it (this time in fact providing all details).

**Example VI.16** (Constant functions are continuous, again).

Let  $X, Y$  be metric spaces, and  $c \in Y$ . Consider the constant function  $f: X \rightarrow Y$  given by

$$f(x) = c \quad \text{for all } x \in X,$$

as in Example VI.3.

Suppose that  $V \subset Y$  is an open subset, and consider the preimage  $f^{-1}[V] \subset X$ . If  $c \in V$ , then the preimage is the whole space,  $f^{-1}[V] = X$ . If  $c \notin V$ , then the preimage is empty,  $f^{-1}[V] = \emptyset$ . Both the whole space and the empty set are open subsets of  $X$ , so the preimage  $f^{-1}[V] \subset X$  is in either case open. Continuity of  $f$  now follows by the characterization of Theorem VI.14.

**Example VI.17** (Identity function is continuous, again).

Let  $X$  be a metric space, and consider the identity function  $\text{id}: X \rightarrow X$  given by

$$\text{id}(x) = x \quad \text{for } x \in X,$$

as in Example VI.4.

Note that the preimage of any subset  $V \subset X$  under the identity function  $\text{id}$  is the subset itself,  $\text{id}^{-1}[V] = V$ . Therefore in particular for an open subset  $V \subset X$ , the preimage  $\text{id}^{-1}[V] = V$  is also open.<sup>7</sup> Continuity of  $\text{id}$  now follows by the characterization of Theorem VI.14.

**Exercise VI.8** (Inclusion is continuous).

Let  $X$  be a metric space and  $X' \subset X$  a subset. Recall that  $X'$  becomes a metric space with the metric it inherits from  $X$ . Consider the inclusion function  $\iota: X' \rightarrow X$  given by

$$\iota(x') = x' \quad \text{for } x' \in X'.$$

Show that  $\iota: X' \rightarrow X$  is continuous.

### *Verifying the openness and closedness of sets*

The following examples illustrate a very common and practical way of verifying that sets defined by conditions involving continuous functions are either open or closed.

Let us start with a little bit to think about.

**Exercise VI.9** (A closed ball is a closed set, again).

With the tools that we now have (particularly Theorem VI.14 and Example VI.13), how

---

<sup>7</sup>It is worth noting that we are not merely using the same set  $X$  as the domain and codomain of the function, but we are moreover equipping it with the same metric, and consequently the same collection of open sets. We later consider the possibility of equipping the domain  $X$  and codomain  $X$  (same set) with different metrics.

would you prove that the closed ball  $\overline{B}_r(x) \subset X$  in a metric space  $(X, d)$  is a closed set? Compare with Example V.29.

**Example VI.18** (An ellipse).

Let  $a, b > 0$ . Consider the ellipse in the plane

$$E = \left\{ (x, y) \in \mathbb{R}^2 \mid \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}.$$

We claim that  $E \subset \mathbb{R}^2$  is a closed subset. To see this, note that its definition involves the function

$$f: \mathbb{R}^2 \rightarrow \mathbb{R} \quad \text{given by} \quad f(x, y) = \frac{x^2}{a^2} + \frac{y^2}{b^2}.$$

In fact, by its very definition, the ellipse is the preimage

$$E = \left\{ (x, y) \in \mathbb{R}^2 \mid f(x, y) = 1 \right\} = f^{-1}[\{1\}]$$

of the singleton  $\{1\} \subset \mathbb{R}$ .

The function  $f$  is a polynomial. For the purposes of this example, let us consider it known that polynomials are continuous functions — this easy result will be proven later in this course as well. The singleton  $\{1\} \subset \mathbb{R}$  is a closed set (apply Exercise V.9). Thus the ellipse  $E$  is closed, as the preimage of the closed set  $\{1\} \subset \mathbb{R}$  under the continuous function  $f$ .

Completely similarly we could conclude that the outside region of the ellipse  $E$ ,

$$\left\{ (x, y) \in \mathbb{R}^2 \mid \frac{x^2}{a^2} + \frac{y^2}{b^2} > 1 \right\} = f^{-1}[(1, \infty)],$$

is open, as the preimage of the open set  $(1, \infty) \subset \mathbb{R}$  under the same continuous function  $f$ . The case of the inside region of the ellipse is similar.

**Example VI.19** (Subspace of functions vanishing at the endpoints of an interval).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and consider the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on this interval equipped with the supremum-norm  $\|\cdot\|_\infty$  and the metric induced by this norm.

Consider also the subset

$$A = \left\{ f \in \mathcal{C}([a, b]) \mid f(a) = 0, f(b) = 0 \right\} \subset \mathcal{C}([a, b])$$

consisting of those functions which vanish at both endpoints,  $a$  and  $b$ .

We claim that this set  $A \subset \mathcal{C}([a, b])$  is closed. To see this, recall first that  $f(a) = \text{ev}_a(f)$  and  $f(b) = \text{ev}_b(f)$ , where  $\text{ev}_z: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  are the evaluation functions studied in Example VI.9. In terms of these, the set  $A$  can be written as an intersection of preimages

$$\begin{aligned} A &= \left\{ f \in \mathcal{C}([a, b]) \mid \text{ev}_a(f) = 0, \text{ev}_b(f) = 0 \right\} \\ &= \text{ev}_a^{-1}[\{0\}] \cap \text{ev}_b^{-1}[\{0\}]. \end{aligned}$$

The singleton  $\{0\} \subset \mathbb{R}$  is closed, so since  $\text{ev}_a, \text{ev}_b: \mathcal{C}([a, b]) \rightarrow \mathbb{R}$  are continuous, the preimages  $\text{ev}_a^{-1}[\{0\}], \text{ev}_b^{-1}[\{0\}] \subset \mathcal{C}([a, b])$  are also closed. Therefore  $A \subset \mathcal{C}([a, b])$  is closed, as the intersection of two closed sets.

**Exercise VI.10** (Cone of positive functions on a closed interval\*).

Let  $[a, b] \subset \mathbb{R}$  be a closed interval and, as in Example IV.27, consider the space  $\mathcal{C}([a, b])$  of continuous real-valued functions on this interval equipped with the supremum-norm  $\|\cdot\|_\infty$  and the metric induced by this norm.

Consider the subset

$$P = \left\{ f \in \mathcal{C}([a, b]) \mid f(x) > 0 \text{ for all } x \in [a, b] \right\} \subset \mathcal{C}([a, b])$$

consisting of all positive functions. Show that  $P \subset \mathcal{C}([a, b])$  is an open set.

Hint: You may want to apply a small modification of Exercise VI.3.

*Composition of continuous functions*

We have so far still had only quite limited examples of continuous functions. To deal with more intricate ones, it is important to be able to construct new continuous functions from simpler building blocks. The operation of composition of functions is one of the most important constructions of this kind.

**Proposition VI.20** (Compositions of continuous functions are continuous).

Let  $(X, d_X)$ ,  $(Y, d_Y)$ ,  $(Z, d_Z)$  be three metric spaces, and let

$$f: X \rightarrow Y \quad \text{and} \quad g: Y \rightarrow Z$$

be continuous functions. Then the composition  $g \circ f: X \rightarrow Z$ , defined by

$$(g \circ f)(x) := g(f(x)) \quad \text{for } x \in X,$$

is also a continuous function.

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

$$\quad \quad \quad \searrow \quad \quad \nearrow$$

$$\quad \quad \quad g \circ f$$

*Proof.* By the characterization of Theorem VI.14, it suffices to show that for all open subsets  $W \subset Z$ , the preimages  $(g \circ f)^{-1}[W] \subset X$  are also open. Observe first the general set theoretic fact that  $(g \circ f)^{-1}[W] = f^{-1}[g^{-1}[W]]$  for any subset  $W \subset Z$ .

Let  $W \subset Z$  be open. By the assumed continuity of  $g$  we have that its preimage  $V = g^{-1}[W] \subset Y$  is also open. Then by the assumed continuity of  $f$  we have that the preimage  $f^{-1}[V] \subset X$  of this open set is also open. On the other hand, we have

$$f^{-1}[V] = f^{-1}[g^{-1}[W]] = (g \circ f)^{-1}[W].$$

We conclude that  $(g \circ f)^{-1}[W] \subset X$  is open. This shows that the composed function  $g \circ f: X \rightarrow Z$  is continuous.  $\square$

If you write down a proof of Proposition VI.20 directly using the  $\varepsilon$ - $\delta$  definition of continuity, you will notice that it becomes lengthier and more convoluted. The simplicity of proofs like these is one advantage of the characterization of continuity in Theorem VI.14.

### VI.3. Exercises

**Exercise VI.11.** Let  $(X, d)$  be a metric space and  $(E, \|\cdot\|)$  a normed space. Complete the proof of the following statement.

**Claim.** If  $f, g: X \rightarrow E$  are two continuous functions, then the function  $f + g: X \rightarrow E$  is also continuous.

*Remark:* The function  $f + g: X \rightarrow E$  is the pointwise sum of the vector valued functions  $f$  and  $g$ , defined by the formula  $(f + g)(x) = f(x) + g(x)$  (the right hand side is vector addition in vector space  $E$ ).

**Exercise VI.12** (A continuous function defined by an integral).

Define a function  $f: \mathcal{C}([0, 10]) \rightarrow \mathcal{C}([0, 10])$  by setting

$$[f(x)](t) = \int_0^t s x(s) \, ds, \quad \text{for } x \in \mathcal{C}([0, 10]) \text{ and } t \in [0, 10].$$



(Here  $f(x)$  is a function  $[0, 10] \rightarrow \mathbb{R}$ , whose value at  $t \in [0, 10]$  is obtained by the above formula.)

Prove that  $f$  is  $K$ -Lipschitz for a suitable  $K \geq 0$ , when the space  $\mathcal{C}([0, 10])$  (both as the domain and codomain of  $f$ ) is equipped with the metric induced by the norm  $\|\cdot\|_\infty$  — i.e., show that for all  $x, y \in \mathcal{C}([0, 10])$  we have

$$\|f(x) - f(y)\|_\infty \leq K \|x - y\|_\infty.$$

Hint: “The triangle equality for integrals”  $\left| \int_a^b h(s) \, ds \right| \leq \int_a^b |h(s)| \, ds$  can be used here.

**Exercise VI.13** (Taking the derivative is not Lipschitz with respect to the uniform norm).

Let

$$\mathcal{C}^1([-1, 1]) = \{x: [-1, 1] \rightarrow \mathbb{R} \mid x' \text{ is continuous}\}$$

be the set of all continuously differentiable functions on the interval  $[-1, 1]$ . Interpret it as a subset  $\mathcal{C}^1([-1, 1]) \subset \mathcal{C}([-1, 1])$ , where the space  $\mathcal{C}([-1, 1])$  of continuous functions is equipped again with the metric induced by the norm  $\|\cdot\|_\infty$ . Consider the function  $g: \mathcal{C}^1([-1, 1]) \rightarrow \mathcal{C}([-1, 1])$  given by

$$[g(x)](t) = x'(t), \quad \text{for } x \in \mathcal{C}^1([-1, 1]) \text{ and } t \in [-1, 1].$$

Show that  $g$  is not  $K$ -Lipschitz for any  $K \geq 0$ .

Hint: In the Lipschitz condition, choose  $y = \vec{0}$  (zero function), in which case it suffices to find a sequence  $(x_n)_{n \in \mathbb{N}}$  of functions for which  $\|x_n\|_\infty \leq 1$  for all  $n \in \mathbb{N}$ , but  $\|x'_n\|_\infty \rightarrow \infty$  as  $n \rightarrow \infty$ .

**Exercise VI.14** (Verifying openness and closedness).

Show that the set

$$U = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x^2 < y^2 + z^2 - xyz + 3 \right\}$$

is open and that the set

$$F = \left\{ (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 2 \text{ and } x \leq 1 + \frac{1}{3} \sin(y) \right\}$$

is closed.

Hint: Express the sets  $U$  and  $F$  appropriately using preimages; for the latter it is best to use two different functions and an intersection of preimages.

The continuity of the functions involved can be considered known.

**Exercise VI.15** (Verifying closedness of a set of probability densities).

Consider the space  $\mathcal{C}([-1, 1])$  of all continuous functions  $f: [-1, 1] \rightarrow \mathbb{R}$ , equipped with the metric induced by the sup-norm  $\|\cdot\|_\infty$ . Consider the subset<sup>8</sup>

$$D = \left\{ p \in \mathcal{C}([-1, 1]) \mid p(x) \geq 0 \, \forall x \in [-1, 1], \int_{-1}^1 p(x) \, dx = 1 \right\}.$$

Show that  $D \subset \mathcal{C}([-1, 1])$  is a closed set.

Hint: You may use the facts that evaluation functions  $f \mapsto f(x)$  (for an arbitrary  $x \in [-1, 1]$ ), and the integration function  $f \mapsto \int_{-1}^1 f(x) \, dx$  are continuous functions  $\mathcal{C}([-1, 1]) \rightarrow \mathbb{R}$  with the chosen metric. Otherwise the ideas are similar to Exercise VI.14.

<sup>8</sup>This subset  $P$  could be interpreted as the set of all continuous probability density functions supported on the interval  $[-1, +1]$ .



## Lecture VII

# Topology and homeomorphisms

In this chapter we define the general notion of *topology*, and consider *homeomorphisms* — mappings which preserve the topology. We define topological equivalence of metrics, and show how Cartesian products of metric spaces can be equipped with various natural metrics which are topologically equivalent.

### VII.1. ♥ General topology

UNDER CONSTRUCTION!

### VII.2. Homeomorphism

If a metric space  $X$  can be mapped continuously and surjectively another metric space  $Y$ , then one can interpret that the “shape” of the space  $X$  can be continuously deformed to that of  $Y$  without tearing anything apart (no discontinuities). If the mapping has an inverse function which is also continuous, and thus intuitively also continuously deforms  $Y$  to  $X$  without tearing anything apart, the two “shapes” can be mutually obtained from each other by continuous deformations. This relation among spaces is known as homeomorphism. The precise definition is the following.

**Definition VII.1** (Homeomorphism).

Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces.<sup>1</sup> A function  $f: X \rightarrow Y$  is called a **homeomorphism** if it is bijective and both  $f: X \rightarrow Y$  and its inverse  $f^{-1}: Y \rightarrow X$  are continuous. If a homeomorphism  $f: X \rightarrow Y$  exists, the spaces  $X$  and  $Y$  are said to be **homeomorphic**, and we denote  $X \approx Y$ .

**Lemma VII.2** (Homeomorphism is an equivalence relation).

*Homeomorphism is an equivalence relation among metric spaces<sup>2</sup>, i.e., the following properties hold:*

- (i) *For any metric space  $X$  we have  $X \approx X$ .*
- (ii) *If  $X \approx Y$ , then we also have  $Y \approx X$ .*
- (iii) *If  $X \approx Y$  and  $Y \approx Z$ , then we also have  $X \approx Z$ .*

*Proof.* Each of the properties is very easy to prove:

---

<sup>1</sup>Since one can define continuity of functions more generally between topological spaces, the definition of homeomorphism is readily meaningful in the generality of topological spaces as well. In this course, however, we focus on the case of metric spaces only.

<sup>2</sup>More generally, homeomorphism is an equivalence relation among topological spaces. The proof remains unchanged.

- (i): The identity function  $\text{id}_X: X \rightarrow X$  is bijective, continuous, and its inverse  $\text{id}_X^{-1} = \text{id}_X$  is also continuous. Therefore  $\text{id}_X$  provides a homeomorphism  $X \approx X$ .
- (ii): If  $f: X \rightarrow Y$  is a homeomorphism, then also its inverse  $f^{-1}: Y \rightarrow X$  is a homeomorphism:  $f^{-1}$  is bijective (since  $f$  is bijective),  $f^{-1}$  is continuous (as the inverse of the homeomorphism  $f$ ), and its inverse  $(f^{-1})^{-1} = f$  is continuous (since  $f$  is a homeomorphism).
- (iii): If  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  are homeomorphisms, then the composition  $g \circ f: X \rightarrow Z$  is also a homeomorphism:  $g \circ f$  is bijective (Exercises I.8 and I.9),  $g \circ f$  is continuous (as the composition of the continuous functions  $f$  and  $g$ ), and its inverse  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$  is continuous (as the composition of the continuous functions  $g^{-1}$  and  $f^{-1}$ ).  $\square$

Let us start with some very simple examples of homeomorphisms among intervals on the real line. In all of the following, as usual, we equip the intervals with the standard metric they inherit from the real line.

**Example VII.3** (All nondegenerate closed intervals are mutually homeomorphic).

Let  $[a_1, b_1] \subset \mathbb{R}$  and  $[a_2, b_2] \subset \mathbb{R}$  be two closed intervals which are both nondegenerate,  $a_1 < b_1$  and  $a_2 < b_2$ . Then the function

$$f: [a_1, b_1] \rightarrow [a_2, b_2] \quad f(x) = a_2 + \frac{b_2 - a_2}{b_1 - a_1}(x - a_1) \quad \text{for } x \in [a_1, b_1]$$

is a homeomorphism: it is clearly bijective and  $M$ -Lipschitz with  $M = \frac{b_2 - a_2}{b_1 - a_1} > 0$ , and its inverse  $y \mapsto a_1 + \frac{b_1 - a_1}{b_2 - a_2}(y - a_2)$  is  $\frac{1}{M}$ -Lipschitz, so both  $f$  and  $f^{-1}$  are continuous (by Proposition VI.6). This shows that any two nondegenerate closed intervals are homeomorphic to each other,  $[a_1, b_1] \approx [a_2, b_2]$ .

**Exercise (✓) VII.1** (All bounded open intervals are mutually homeomorphic).

Show that if  $a_1 < b_1$  and  $a_2 < b_2$ , then the bounded open intervals  $(a_1, b_1) \subset \mathbb{R}$  and  $(a_2, b_2) \subset \mathbb{R}$  are homeomorphic,  $(a_1, b_1) \approx (a_2, b_2)$ .

**Exercise (✓) VII.2** (Bounded half-open intervals are mutually homeomorphic).

Show that if  $a_1 < b_1$  and  $a_2 < b_2$ , then there are homeomorphisms  $(a_1, b_1] \approx (a_2, b_2]$  and also  $(a_1, b_1] \approx [a_2, b_2)$ .

**Exercise VII.3** (Bounded and an unbounded half-open intervals are homeomorphic).

Show that  $[0, 1)$  and  $[0, \infty)$  are homeomorphic.

*Hint:* You can use the fact that rational functions are continuous. This was stated in Corollary III.11, and we admit for now that the definitions of continuity given in Chapters III and VI agree. This will be seen in detail in Chapter VIII.

**Remark VII.4** (All bounded and unbounded half-open intervals are homeomorphic).

By Exercise VII.3 we have a homeomorphism  $[0, \infty) \approx [0, 1)$ . By Exercise VII.2 we also have homeomorphism  $[0, 1) \approx [a, b)$  and  $[0, 1) \approx (a, b]$  for any  $a < b$ . By Lemma VII.2(iii) we get that the non-negative real axis  $[0, \infty)$  is homeomorphic to any bounded open interval,  $[0, \infty) \approx [a, b)$  and  $[0, \infty) \approx (a, b]$ . It is very easy to also see that  $[0, \infty) \approx [a, \infty)$  and  $[0, \infty) \approx (-\infty, b]$ , so in fact all bounded and unbounded half-open intervals are mutually homeomorphic.

Similarly one can prove that bounded open intervals and unbounded open intervals are homeomorphic. Let us provide one such homeomorphism.

**Example VII.5** (A homeomorphism from an open interval to the real line).

Consider the open interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ . The tangent function

$$\tan: \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$$

is bijective; its inverse is the arcustangent  $\arctan: \mathbb{R} \rightarrow (-\frac{\pi}{2}, \frac{\pi}{2})$ . We consider it known that both  $\tan$  and  $\arctan$  are continuous.<sup>3</sup> Therefore we get a homeomorphism  $(-\frac{\pi}{2}, \frac{\pi}{2}) \approx \mathbb{R}$ .

One might suspect that closed intervals are not homeomorphic to open intervals, and neither are homeomorphic to half-open intervals. This is indeed the case. You could try to prove it directly, but after Chapters XI and XII we have better general tools which make the proofs significantly easier.

All of the above examples are very easy homeomorphisms. Let us now give one example of something natural that turns out *not* to be a homeomorphism.

**Example VII.6** (A continuous map from interval to circle is not a homeomorphism).

Consider the unit circle

$$\mathbf{S}^1 = \left\{ (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1 \right\}.$$

and the half-open interval  $[0, 2\pi)$ . It is very natural to define a mapping

$$f: [0, 2\pi) \rightarrow \mathbf{S}^1 \quad f(\theta) = (\cos(\theta), \sin(\theta)) \quad \text{for } \theta \in [0, 2\pi).$$

This  $f$  is bijective. The trigonometric functions  $\theta \mapsto \cos(\theta)$  and  $\theta \mapsto \sin(\theta)$  have derivatives bounded by 1 and as such are 1-Lipschitz (see Example VI.7), and from this it is not difficult to show that also  $f$  is  $M$ -Lipschitz ( $M = 1$  works but  $M = 2$  is slightly easier to prove) and in particular  $f$  is continuous.

But  $f$  is *not* a homeomorphism between  $[0, 2\pi)$  and  $\mathbf{S}^1$ !

Indeed, the inverse  $f^{-1}: \mathbf{S}^1 \rightarrow [0, 2\pi)$  is not continuous. In  $[0, 2\pi)$ , consider an open ball<sup>4</sup> of small positive radius  $\varepsilon < 2\pi$  — it is of the form  $[0, \varepsilon) \subset [0, 2\pi)$ . The image  $f[[0, \varepsilon)) \subset \mathbf{S}^1$  contains  $(1, 0) = f(0) \in \mathbf{S}^1$ , but does not contain the point  $(\cos(2\pi - \delta), \sin(2\pi - \delta)) = f(2\pi - \delta) \in \mathbf{S}^1$  for any small  $\delta \in (0, 2\pi - \varepsilon)$ . Since these points are at distances  $d(f(0), f(2\pi - \delta)) < M\delta$  from  $f(0) = (1, 0)$ , we see that no open ball in  $\mathbf{S}^1$  centered at  $f(0)$  is contained in  $f[[0, \varepsilon))$ , so the image  $f[[0, \varepsilon)) \subset \mathbf{S}^1$  is not open. Applying this to preimages under the inverse function  $f^{-1}: \mathbf{S}^1 \rightarrow [0, 2\pi)$ , we see that  $(f^{-1})^{-1}[[0, \varepsilon)) = f[[0, \varepsilon))$  is not open, and since  $[0, \varepsilon) \subset [0, 2\pi)$  was an open set, this shows (by Theorem VI.14) that the inverse  $f^{-1}: \mathbf{S}^1 \rightarrow [0, 2\pi)$  is not a continuous function. In particular,  $f$  is not a homeomorphism.

The above reasoning only shows that this particular function  $f$  is not a homeomorphism between the half-open interval  $[0, 2\pi)$  and the circle  $\mathbf{S}^1$  — it does not say that  $[0, 2\pi)$  and  $\mathbf{S}^1$  are not homeomorphic (i.e., that no homeomorphism exists between them). In Chapter XI we will have good tools to prove that such a homeomorphism indeed is impossible, and we have  $[0, 2\pi) \not\approx \mathbf{S}^1$ . The intuitive idea, however, should be clear: a bijection in the direction  $\mathbf{S}^1 \rightarrow [0, 2\pi)$  cannot be continuous — it has to tear apart the circle at some point.

Since all of the above examples have pertained to merely “one-dimensional spaces” (“lines and curves”), let us take one slightly different example.

**Exercise VII.4** (The open unit disk is homeomorphic to the plane).

Prove that the open unit disk  $\mathcal{B}_1(\vec{0}) \subset \mathbb{R}^2$  in the Euclidean plane  $\mathbb{R}^2$  and the whole plane  $\mathbb{R}^2$  are homeomorphic,  $\mathcal{B}_1(\vec{0}) \approx \mathbb{R}^2$ .

<sup>3</sup>One way to see this is to observe that restricted to closed intervals of their respective domains of definition,  $\tan$  and  $\arctan$  are continuously differentiable, and therefore Lipschitz by Example VI.8.

<sup>4</sup>Note that we are taking open balls in the metric space  $[0, 2\pi)$  equipped with the metric it inherits from  $\mathbb{R}$ . Following the ideas of Lemma V.25, these open balls are just intersections of open balls on  $\mathbb{R}$  with  $[0, 2\pi) \subset \mathbb{R}$ . In particular the ball of radius  $\varepsilon < 2\pi$  centered at 0 is  $\mathcal{B}_\varepsilon(0) = (-\varepsilon, \varepsilon) \cap [0, 2\pi) = [0, \varepsilon)$ .

*Hint:* One idea is a radial mapping in polar coordinates, using Exercise VII.3. Detailed proofs of continuity may require some work until we develop better tools later. . .

*Fun: homeomorphism classes of letters*

Let us borrow a fun example from [Väi99]: consider the capital letters

A, B, C, D, E, F, G, H, I, J, K, L, M  
N, O, P, Q, R, S, T, U, V, W, X, Y, Z

as subsets of the plane  $\mathbb{R}^2$ , so they in particular inherit a metric and become metric spaces. Which of these are homeomorphic?

Some “obvious” homeomorphisms are  $D \approx O$  (both homeomorphic to the circle  $\mathbf{S}^1$ ) and  $J \approx S$  (both homeomorphic to the interval  $[0, 1]$ ) and  $Y \approx T$ . Let us not spoil any more of the fun — you should try to classify the letters<sup>5</sup> into equivalence classes under homeomorphisms (feel free to resort to intuitive arguments, since we did not even give precise definitions of the subsets corresponding to the letters. . .).

*The quintessential topology joke*

The donut is homeomorphic to the coffee cup.

<https://www.shapeways.com/product/6CJQ9GXWW/topology-joke>

## Homeomorphism as mappings that preserve topology

Observe the following.

**Proposition VII.7** (A homeomorphism preserves the openness of sets).

*Let  $f: X \rightarrow Y$  be a homeomorphism. Then a subset  $A \subset X$  is open if and only if its image  $f[A] \subset Y$  is open.*

*Proof.* Suppose that  $A \subset X$  is such that  $f[A] \subset Y$  is open. Then by Theorem VI.14 and the continuity of  $f: X \rightarrow Y$ , the preimage  $f^{-1}[f[A]] \subset X$  is also open. By bijectivity of  $f$  we have  $f^{-1}[f[A]] = A$ , so  $A \subset X$  is open.

Suppose then that  $A \subset X$  is open. Then by Theorem VI.14 and the continuity of the inverse function  $f^{-1}: Y \rightarrow X$ , its preimage  $(f^{-1})^{-1}[A] \subset Y$  is also open. But we have  $(f^{-1})^{-1}[A] = f[A]$ , so the image  $f[A] \subset Y$  is open.  $\square$

Also the converse holds.

**Exercise (✓) VII.5** (Bijections preserving openness are homeomorphisms).

Suppose that  $f: X \rightarrow Y$  is a bijection with the property that subsets  $A \subset X$  are open if and only if their images  $f[A] \subset Y$  are open. Prove that then  $f$  is a homeomorphism.

In view of the above, homeomorphisms are precisely bijective mappings between metric spaces which preserve the openness of subsets. The collection of all open subsets of a space is called the topology of the space, see Section VII.1 for a formal

<sup>5</sup>If you didn’t get enough, include numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 as well! Then the lowercase letters a, b, c, . . . (how about ä, ö?), the Greek alphabet  $\alpha, \beta, \gamma, \dots$ , simplified Chinese characters, traditional Chinese characters, . . .

definition. In view of this, homeomorphisms are precisely bijections that preserve topology.

### VII.3. Equivalence of metrics

A given set  $X$  can be equipped with various different metrics. The choice of metric  $d$  on  $X$  determines the open sets, i.e., the topology on  $X$ . It is possible for different choices of metric lead to the same notion of open sets — this leads to the following notion.

**Definition VII.8** (Topologically equivalent metrics).

Let  $X$  be a set. Let  $d_1$  and  $d_2$  be two metrics on  $X$ , and

$$\begin{aligned}\mathcal{T}_1 &= \{U_1 \subset X \mid U_1 \text{ is open with respect to } d_1\} \\ \mathcal{T}_2 &= \{U_2 \subset X \mid U_2 \text{ is open with respect to } d_2\}\end{aligned}$$

the corresponding collections of open subsets with respect to each of the two. We call the metrics  $d_1$  and  $d_2$  (**topologically**) **equivalent** if the collections of open sets determined by both are the same,  $\mathcal{T}_1 = \mathcal{T}_2$ .

The most important practical situation in which we will use this notion is metrics on product spaces, see Section VII.4. Let us nevertheless start by giving a few elementary examples.

The following simple observation shows that there are always infinitely many metrics equivalent with any given one (barring the trivial cases when the set  $X$  is either empty or a singleton, in which cases there are no nonzero distances).

**Example VII.9** (Constant multiple of the same metric).

Let  $d_1: X \times X \rightarrow [0, \infty)$  be a metric on  $X$ , and  $c > 0$  a constant. Define  $d_2: X \times X \rightarrow [0, \infty)$  by  $d_2(x, y) = c d_1(x, y)$  for all  $x, y \in X$ . Properties (M-s), (M- $\Delta$ ), and (M-0) for  $d_2$  clearly follow from the corresponding properties for  $d_1$ , so  $d_2$  is also a metric on  $X$ .

Let us denote the open balls of radius  $r > 0$  centered at  $z \in X$  with respect to  $d_1$  and  $d_2$  by

$$\mathcal{B}_r^1(z) = \{x \in X \mid d_1(z, x) < r\} \qquad \mathcal{B}_r^2(z) = \{x \in X \mid d_2(z, x) < r\}.$$

We clearly have  $\mathcal{B}_r^1(z) = \mathcal{B}_{cr}^2(z)$  for any  $r > 0$  and  $z \in X$ . In view of Definition V.18 it is therefore clear that a subset  $U \subset X$  is open with respect to  $d_1$  if and only if it is open with respect to  $d_2$ , i.e.,  $U \in \mathcal{T}_1 \Leftrightarrow U \in \mathcal{T}_2$ , where  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are the collections of open sets determined by the metrics  $d_1$  and  $d_2$ . In other words,  $\mathcal{T}_1 = \mathcal{T}_2$ , so the metrics  $d_1$  and  $d_2$  are topologically equivalent.

Thus multiplying distances by a constant  $c > 0$  can be thought of as a “change of units of distances”: if one person measures distances in miles and another in kilometres, they disagree on the numerical values of distances, but they (hopefully) agree on the “shape” of the space they both live in.

From the following we see that any metric is topologically equivalent to a bounded metric.

**Example VII.10** (Truncation of a metric).

Let  $d_1: X \times X \rightarrow [0, \infty)$  be a metric on  $X$ , and  $c > 0$  a constant. Define  $d_2: X \times X \rightarrow [0, \infty)$  by  $d_2(x, y) = \min\{d_1(x, y), c\}$  for all  $x, y \in X$ . Properties (M-s) and (M-0) for  $d_2$  clearly follow from the corresponding properties for  $d_1$ . Also the triangle inequality (M- $\Delta$ ) for  $d_2$  holds by virtue of the easy observation

$$\min\{d_1(x, y) + d_1(y, z), c\} \leq \min\{d_1(x, y), c\} + \min\{d_1(y, z), c\}$$

and the triangle inequality for  $d_1$ . Therefore  $d_2$  is also a metric on  $X$ .

Denote the open balls of radius  $r > 0$  centered at  $z \in X$  with respect to  $d_1$  and  $d_2$  by  $\mathcal{B}_r^1(z)$  and  $\mathcal{B}_r^2(z)$  again. For  $0 < r \leq c$  we have  $\mathcal{B}_r^1(z) = \mathcal{B}_r^2(z)$  for any  $z \in X$ . In Definition V.18 of open sets we could have restricted the radius of balls to be below the constant  $c > 0$  (if we found larger radius balls around points, then we certainly could use smaller ones as well). It is therefore clear that a subset  $U \subset X$  is open with respect to  $d_1$  if and only if it is open with respect to  $d_2$ . As before, this says that metrics  $d_1$  and  $d_2$  are topologically equivalent.

Thus truncating distances to at most a constant  $c > 0$  can be thought of as a “allowing teleportation at a fixed cost  $c$ ”: for any distances greater than  $c$  it is then advantageous to resort to the teleportation option. But the local geometry (at distances below  $c$ ) remains unchanged, and this is what determines the topology.

The equivalence of metrics can be phrased in terms of homeomorphisms.

**Lemma VII.11** (A characterization of equivalence of metrics).

*Let  $X$  be a set. Let  $d_1$  and  $d_2$  be two metrics on  $X$ . Consider the identity mapping of the set  $X$*

$$\text{id}_X: X \rightarrow X \qquad \text{id}_X(x) = x \quad \text{for } x \in X,$$

*as a mapping between the metric spaces  $(X, d_1)$  and  $(X, d_2)$ ; i.e., the domain  $X$  equipped with metric  $d_1$  and the codomain  $X$  with the metric  $d_2$ . Then  $\text{id}_X$  is a homeomorphism if and only if the metrics  $d_1$  and  $d_2$  are topologically equivalent.*

*Proof.* **UNDER CONSTRUCTION!**

□

**VII.4. Product spaces**

Recall that the Cartesian product of two sets  $X$  and  $Y$  is the set

$$X \times Y = \left\{ (x, y) \mid x \in X, y \in Y \right\}$$

of ordered pairs  $(x, y)$ , whose first member  $x$  belongs to the first set  $X$  and second member  $y$  belongs to the second set  $Y$ .

It is very common that we want to form Cartesian products of metric spaces.

Some examples are provided by geometric objects. The *plane*  $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$  is the Cartesian product of the real line  $\mathbb{R}$  with itself (more generally, the  $d$ -dimensional space  $\mathbb{R}^d$  is a  $d$ -fold Cartesian product of copies of  $\mathbb{R}$ ), a *rectangle*  $[a_1, b_1] \times [a_2, b_2]$  is the Cartesian product of two intervals, a (*solid*) *cylinder* is the Cartesian product  $\mathbf{D}^2 \times [a, b]$  of a disk  $\mathbf{D}^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$  with an interval  $[a, b]$ , and the *torus* is the Cartesian product  $\mathbf{S}^1 \times \mathbf{S}^1$  of a circle  $\mathbf{S}^1$  with itself.






FIGURE VII.1. Some familiar geometric objects are Cartesian products, for example rectangles  $[a_1, b_1] \times [a_2, b_2]$ , (solid) cylinders  $\mathbf{D}^2 \times [a, b]$ , and torus  $\mathbf{S}^1 \times \mathbf{S}^1$ .

Even more importantly, for a function of several variables, the domain of definition is naturally a Cartesian product<sup>6</sup>, in which the sets of allowed values of each variable are the factors of the Cartesian product. Examples we have seen already include an *inner product*  $\langle \cdot, \cdot \rangle$ , which is a real-valued function of a pair of vectors in an inner product space  $V$ , i.e., a function  $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$  defined on the Cartesian product of  $V$  with itself, and a *metric*  $d$ , which is a function of a pair of points in a metric space  $X$ , i.e., a function  $d: X \times X \rightarrow [0, \infty)$  defined on the Cartesian product of  $X$  with itself. In applications to physics, for example, one might consider a field  $\phi$  (e.g., pressure, temperature, magnetic field, ...) which depends on both spatial and temporal variables so that  $\phi(t, x)$  represents the field value at time  $t \in [0, \infty)$  and position  $x \in D \subset \mathbb{R}^3$  so that  $\phi$  is a function defined on  $[0, \infty) \times D$ .<sup>7</sup>

It therefore seems quite important that we can equip the Cartesian product  $X \times Y$  of two metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  with enough structure so that for example continuity of functions of several variables makes sense — we would like to equip  $X \times Y$  with a metric. It turns out that there are a few different choices of metric on the product space  $X \times Y$ , each choice having its own advantages and each natural in certain situations, but none of the choices generally better than the others. At first this might seem like a problem — if the choice of metric is not canonical, does it mean for example that the continuity of functions of several variables depends on what choice one makes? The answer is that while the choice of *metric* on a product space is not canonical, the choice of *topology* is! All of the different natural metrics are topologically equivalent, so for example continuity of functions (and other topological properties) does not depend on the choice!

For simplicity, we will only discuss Cartesian products of two metric spaces  $(X, d_X)$  and  $(Y, d_Y)$ . Treating Cartesian products of finitely many metric spaces works similarly, only the notation gets somewhat more tangled.<sup>8</sup>

<sup>6</sup>...or sometimes a subset of a Cartesian product...

<sup>7</sup>The precise temporal and spatial domains would of course depend on the application, this is just one quite typical example to give a concrete example.

To convince yourself about the prominence of such examples in different areas of physics, just look at the wave equation, the heat equation, Maxwell's equations, the Schrödinger equation, ....

<sup>8</sup>One could also consider Cartesian products of infinitely many metric spaces, not just as an academic curiosity; but also as something that is often relevant (for example in probability theory). The details become just slightly more complicated for countably infinite Cartesian products, but the

### Metrics on product space

Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces. As three natural metrics on the Cartesian product  $X \times Y$ , we consider

$$d_1, d_2, d_\infty : (X \times Y) \times (X \times Y) \rightarrow [0, \infty) \quad (\text{VII.1})$$

given by

$$d_1((x, y), (x', y')) = d_X(x, x') + d_Y(y, y'), \quad (\text{VII.2})$$

$$d_2((x, y), (x', y')) = \sqrt{d_X(x, x')^2 + d_Y(y, y')^2}, \quad (\text{VII.3})$$

$$d_\infty((x, y), (x', y')) = \max\{d_X(x, x'), d_Y(y, y')\}, \quad (\text{VII.4})$$

for  $(x, y) \in X \times Y$  and  $(x', y') \in X \times Y$ . Of course we need to prove that each of these three satisfies the properties (M-s), (M- $\Delta$ ), (M-0), and therefore indeed is a metric on the set  $X \times Y$ .

**Lemma VII.12** ( $d_1, d_2, d_\infty$  are metrics on the product space).

*Each of the formulas (VII.2) – (VII.4) defines a metric on  $X \times Y$ .*

*Proof.* **UNDER CONSTRUCTION!**

□

### Equivalence of the product space metrics

The important observation is that the choice of metric does not matter for any purely topological features — the metrics given by (VII.2) – (VII.4) are topologically equivalent.

**Proposition VII.13** (The natural metrics on the product space are equivalent).

*The metrics  $d_1, d_2, d_\infty$  on  $X \times Y$  given by (VII.2) – (VII.4) are topologically equivalent.*

*Proof.* **UNDER CONSTRUCTION!**

□

### Projections

On the Cartesian product  $X \times Y$  we have two functions, which pick the first and second “coordinates” of a point  $(x, y) \in X \times Y$ ,

$$\begin{aligned} \text{pr}_1 : X \times Y &\rightarrow X & \text{pr}_2 : X \times Y &\rightarrow Y \\ (x, y) &\mapsto x & (x, y) &\mapsto y. \end{aligned} \quad (\text{VII.5})$$

These are called the **projections** to  $X$  and  $Y$ . Let us first observe that the projections are continuous functions, when the product space  $X \times Y$  is equipped with any of the natural metrics.

---

conclusion remains similar: there are various natural metrics, each giving rise to the same topology. In uncountably infinite Cartesian products, however, the natural topology does not necessarily come from a metric anymore. Uncountably infinite Cartesian products are one important source of *non-metrizable topologies*, and they illustrate the advantages of considering general topological spaces instead of metric spaces. For simplicity, in this course we restrict our attention to just finite Cartesian products.

**Lemma VII.14** (Projections are continuous).

*The projections  $\text{pr}_1: X \times Y \rightarrow X$  and  $\text{pr}_2: X \times Y \rightarrow Y$  given by (VII.5) are continuous functions.*

*Proof.* Observe first that because of the topological equivalence of the natural metrics  $d_1, d_2, d_\infty$  (Proposition VII.13), continuity does not depend on which of these metrics we use on the domain  $X \times Y$ .

The cases of both projections are similar, so let us only consider  $\text{pr}_1: X \times Y \rightarrow X$ . We will prove that  $\text{pr}_1$  is 1-Lipschitz when the domain  $X \times Y$  is equipped with the metric  $d_1$  of (VII.2) and the codomain  $X$  with its own metric  $d_X$ . The continuity of  $\text{pr}_1$  will then follow from Proposition VI.6.

So let  $(x, y), (x', y') \in X \times Y$ . For the Lipschitz property, we calculate

$$\begin{aligned} d_X(\text{pr}_1(x, y), \text{pr}_1(x', y')) &= d_X(x, x') && \text{(by def. of } \text{pr}_1) \\ &\leq d_X(x, x') + d_Y(y, y') && \text{(since } d_Y(y, y') \geq 0) \\ &= d_1((x, y), (x', y')). && \text{(by def. of } d_1) \end{aligned}$$

This shows the asserted 1-Lipschitz property and in particular proves continuity of  $\text{pr}_1$ .  $\square$

### Component functions

Suppose that  $(X, d_X)$ ,  $(Y, d_Y)$ , and  $(Z, d_Z)$  are metric spaces. For a function

$$f: Z \rightarrow X \times Y$$

with values in the product space  $X \times Y$ , we can consider the **component functions** (or coordinate functions) obtained by composing the function  $f$  with one of the projections (VII.5):

$$\begin{aligned} f_1 &= \text{pr}_1 \circ f : Z \rightarrow X \\ f_2 &= \text{pr}_2 \circ f : Z \rightarrow Y, \end{aligned}$$

in terms of which the function  $f$  can be written as

$$f(z) = (f_1(z), f_2(z)) \quad \text{for } z \in Z.$$

The continuity of a function with values in a product space can be characterized in terms of its component functions.

**Proposition VII.15** (Continuity of functions with values in a product space).

*A function  $f: Z \rightarrow X \times Y$  is continuous if and only if both of its component functions  $f_1: Z \rightarrow X$  and  $f_2: Z \rightarrow Y$  are continuous.*

*Proof.* If  $f: Z \rightarrow X \times Y$  is continuous, then both of its component functions  $f_1$  and  $f_2$  are also continuous in view of the facts that the projections are continuous (Lemma VII.14), and compositions of continuous functions are continuous (Proposition VI.20).

Conversely, suppose that both  $f_1: Z \rightarrow X$  and  $f_2: Z \rightarrow Y$  are continuous. By virtue of the topological equivalence of the natural metrics  $d_1, d_2, d_\infty$  (Proposition VII.13), the continuity of  $f$  does not depend on which of these metrics we use on the codomain  $X \times Y$ , so let us use the metric  $d_1$  of (VII.2). To prove continuity, let  $z \in Z$  and let  $\varepsilon > 0$ . By the assumed continuity of  $f_1$  and  $f_2$ , there exists  $\delta_1 > 0$  and  $\delta_2 > 0$  so that

$$\begin{aligned} d_Z(z, z') < \delta_1 &\implies d_X(f_1(z), f_1(z')) < \frac{\varepsilon}{2} \\ d_Z(z, z') < \delta_2 &\implies d_Y(f_2(z), f_2(z')) < \frac{\varepsilon}{2}. \end{aligned}$$

Let  $\delta = \min \{\delta_1, \delta_2\} > 0$ . Then for  $z' \in Z$  such that  $d_Z(z, z') < \delta$  we can use both of the above, and we get

$$\begin{aligned}
 & d_1(f(z), f(z')) \\
 &= d_1\left((f_1(z), f_2(z)), (f_1(z'), f_2(z'))\right) && (f \text{ in terms of } f_1 \text{ and } f_2) \\
 &= d_X(f_1(z), f_1(z')) + d_Y(f_2(z), f_2(z')) && (\text{definition of } d_1) \\
 &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. && (d_Z(z, z') < \delta_1 \text{ and } d_Z(z, z') < \delta_2)
 \end{aligned}$$

This proves that  $f: Z \rightarrow X \times Y$  is continuous. □

## Lecture VIII

### Sequences in metric spaces

One of the straightforward but important notions that are easy to make sense of in metric spaces is sequences and their limits.

#### VIII.1. Sequences and limits in metric spaces

Throughout this section, let  $(X, d)$  be a metric space.

A **sequence** in the metric space  $(X, d)$  is an “infinite list”

$$(x_1, x_2, x_3, \dots)$$

of points  $x_1, x_2, x_3, \dots \in X$ . A precise definition again is a function from the natural numbers to this space,

$$\xi: \mathbb{N} \rightarrow X,$$

and the “list” consists of the values  $x_n = \xi(n)$  for  $n \in \mathbb{N}$ . As before, the sequence is usually denoted by

$$(x_1, x_2, x_3, \dots) \quad \text{or} \quad (x_n)_{n=1}^{\infty} \quad \text{or} \quad (x_n)_{n \in \mathbb{N}}.$$

It is straightforward to generalize the notion of a limit of a sequence to metric spaces. Indeed, the intuitive meaning in the following definition remains exactly the same as in Definition II.11: however close we want to get to the limit, we achieve this by going far enough to the tail of the sequence. [\[ILLUSTRATION\]](#)

**Definition VIII.1** (Limit of a sequence in a metric space).

A sequence  $(x_n)_{n \in \mathbb{N}}$  in a metric space  $X$  has a **limit**  $a \in X$  if for all  $\varepsilon > 0$  there exists an  $n_\varepsilon \in \mathbb{N}$  such that  $d(x_n, a) < \varepsilon$  whenever  $n \geq n_\varepsilon$ . We then denote

$$\lim_{n \rightarrow \infty} x_n = a \quad \text{or} \quad x_n \rightarrow a \text{ as } n \rightarrow \infty.$$

**Remark VIII.2** (Definition of a limit in logical symbols).

According to Definition VIII.1, the logical meaning of the statement

$$\lim_{n \rightarrow \infty} x_n = a$$

is

$$\forall \varepsilon > 0: \exists n_\varepsilon \in \mathbb{N}: \forall n \geq n_\varepsilon: d(x_n, a) < \varepsilon.$$

Another equivalent logical formulation is  $\forall \varepsilon > 0: \exists n_\varepsilon \in \mathbb{N}: n \geq n_\varepsilon \Rightarrow d(x_n, a) < \varepsilon$ .

Instead of saying that a sequence  $(x_n)_{n \in \mathbb{N}}$  has a limit  $a$ , we occasionally say that  $(x_n)_{n \in \mathbb{N}}$  **converges to**  $a$ , or that  $x_n$  **tends to**  $a$  (as  $n$  tends to  $\infty$ ). If a sequence converges to some limit (especially in situations when we do not specify the limit explicitly), we say that the sequence **converges**, or that it is **convergent**.

By comparing Definition VIII.1 with Definition II.11 it is evident that limits of real number sequences as defined in Chapter II coincide with limits of sequences in the metric space  $(\mathbb{R}, d_{\mathbb{R}})$ , the real line with its standard metric.

Note again that when a limit exists, it is unique.

**Lemma VIII.3** (Uniqueness of limits of sequences in metric spaces).

*If a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  has as its limits both  $a \in X$  and  $b \in X$ , then necessarily  $a = b$ .*

*Proof.* Suppose that both  $a \in X$  and  $b \in X$  are limits of  $(x_n)_{n \in \mathbb{N}}$ . Let  $\varepsilon > 0$ . Then since  $\frac{\varepsilon}{2} > 0$ , by definition of limits there exists  $n', n'' \in \mathbb{N}$  such that  $d(x_n, a) < \frac{\varepsilon}{2}$  for  $n \geq n'$  and  $d(x_n, b) < \frac{\varepsilon}{2}$  for  $n \geq n''$ . In particular letting  $n_0 = \max\{n', n''\}$  and applying the triangle inequality (M- $\Delta$ ) to  $a, x_{n_0}, b \in X$ , we get

$$0 \leq d(a, b) \leq \underbrace{d(a, x_{n_0})}_{< \varepsilon/2} + \underbrace{d(x_{n_0}, b)}_{< \varepsilon/2} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Since  $\varepsilon > 0$  was arbitrary, this implies  $d(a, b) = 0$ , which by (M-0) implies  $a = b$ .  $\square$

In particular, the notation  $\lim_{n \rightarrow \infty} x_n$  for a convergent sequence  $(x_n)_{n \in \mathbb{N}}$  is thus indeed unambiguous.

There are various conditions which are equivalent with the definition of a limit. Depending on the situation, it may be more convenient to verify or apply one of these instead of using Definition VIII.1 directly. The last of the equivalent conditions below refers only to the notion of an open set, and serves as the definition of a limit in general topological spaces, where we do not in general have a metric.<sup>1</sup>

**Proposition VIII.4** (Characterizations of limits in metric spaces).

*Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $X$  and  $x \in X$  a point. Then the following are equivalent:*

- (i)  $\lim_{n \rightarrow \infty} x_n = x$ ;
- (ii)  $\lim_{n \rightarrow \infty} d(x_n, x) = 0$ ;
- (iii) *for any neighborhood  $U$  of  $x$ , only finitely many members of sequence  $(x_n)_{n \in \mathbb{N}}$  are not in  $U$ .*

*Proof.* We will prove the equivalence (i)  $\Leftrightarrow$  (ii) and the implications (i)  $\Rightarrow$  (iii) and (iii)  $\Rightarrow$  (i) separately. The equivalence of all three conditions follows.

*proof of (i)  $\Leftrightarrow$  (ii):* The logical equivalence of (i) and (ii) is clear from Definitions VIII.1 and II.11, once one notices that by non-negativity of metrics the condition  $d(x_n, x) < \varepsilon$  is equivalent to  $|d(x_n, x) - 0| < \varepsilon$ .

*proof of (i)  $\Rightarrow$  (iii):* If  $U \subset X$  is a neighborhood of  $x$ , then there exists an  $\varepsilon > 0$  such that  $\mathcal{B}_{\varepsilon}(x) \subset U$ . If we assume  $\lim_{n \rightarrow \infty} x_n = x$ , then for some  $n_{\varepsilon} \in \mathbb{N}$ , and all  $n \geq n_{\varepsilon}$  we have  $x_n \in \mathcal{B}_{\varepsilon}(x) \subset U$ . Therefore only the finitely many first members  $x_1, x_2, \dots, x_{n_{\varepsilon}-1}$  are possibly not in  $U$ .

---

<sup>1</sup>Be warned, however, that in general topology the notion of a limit may not enjoy all of the good properties familiar from metric spaces. In particular in general topology, a limit may not be unique.

*proof of (iii)  $\Rightarrow$  (i):* Assume that for any neighborhood  $U$  of  $x$ , only finitely many members of sequence  $(x_n)_{n \in \mathbb{N}}$  are not in  $U$ . Let  $\varepsilon > 0$ . The open ball  $\mathcal{B}_\varepsilon(x)$ , in particular, is a neighborhood of  $x$ , and therefore by assumption for only finitely many indices  $n_1 < \cdots < n_m$  we have  $x_{n_k} \notin \mathcal{B}_\varepsilon(x)$ . For  $n \geq n' = n_m + 1$  we then have  $x_n \in \mathcal{B}_\varepsilon(x)$ , i.e.,  $d(x_n, x) < \varepsilon$ . This shows that  $\lim_{n \rightarrow \infty} x_n = x$ .  $\square$

Boundedness of sequences in a metric space  $(X, d)$  is defined as follows.

**Definition VIII.5** (Boundedness of a sequence).

A sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  is **bounded** if there exists some  $z \in X$  and  $r > 0$  such that  $x_n \in \mathcal{B}_r(z)$  for all  $n \in \mathbb{N}$ .

**Exercise VIII.1** (Boundedness of a sequence is boundedness of the set of its points).

Prove that a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  is bounded (in the sense of Definition VIII.5) if and only if the subset  $A = \{x_n \mid n \in \mathbb{N}\} \subset X$  is bounded (in the sense of Chapter V).

Hint: Recall Example V.15 and Exercise V.3.

**Exercise VIII.2** (Convergent sequences are bounded).

Show that if  $(x_n)_{n \in \mathbb{N}}$  is a sequence in  $X$  which is convergent, then  $(x_n)_{n \in \mathbb{N}}$  is bounded.

**Exercise VIII.3** (A sequence on the real line with 0/1-metric).

Consider the metric space  $(\mathbb{R}, d)$ , where the metric  $d$  is the discrete 0-1-metric. Then the sequence  $(x_n)_{n \in \mathbb{N}}$  in  $(\mathbb{R}, d)$  defined by the formula  $x_n = \frac{1}{n}$  does not have 0 as its limit.

**Example VIII.6** (A sequence converging to the zero function in  $L^1$ ).

Consider the space  $\mathcal{C}([0, 1])$  of all continuous functions  $f: [0, 1] \rightarrow \mathbb{R}$  with the  $L^1$ -norm  $\|f\|_1 = \int_0^1 |f(t)| dt$  and the metric induced by it.

For  $n \in \mathbb{N}$ , define the element  $f_n \in \mathcal{C}([0, 1])$  by

$$f_n(t) = \begin{cases} 1 - nt & \text{for } 0 \leq t \leq \frac{1}{n} \\ 0 & \text{for } \frac{1}{n} \leq t \leq 1 \end{cases}$$

(the continuity of  $f_n$  is easy to check).

We claim that the limit of the sequence  $(f_n)_{n \in \mathbb{N}}$  in  $\mathcal{C}([0, 1])$  is the zero function  $0 \in \mathcal{C}([0, 1])$ . This is done by an easy calculation:

$$d(0, f_n) = \|f_n - 0\|_1 = \int_0^1 |f(t) - 0| dt = \int_0^{1/n} (1 - nt) dt = \frac{1}{2n}.$$

Since the right hand side tends to 0 (in  $\mathbb{R}$ ) as  $n \rightarrow \infty$ , we get  $\lim_{n \rightarrow \infty} d(f_n, 0) = 0$ . It then follows from Proposition VIII.4 that the sequence  $(f_n)_{n \in \mathbb{N}}$  converges to the zero function  $0 \in \mathcal{C}([0, 1])$ .

## VIII.2. Characterization of closure and closed sets

Sequences provide intuitive and commonly used characterizations of closure and closed sets.

If  $A \subset X$ , we say that  $(a_n)_{n \in \mathbb{N}}$  is a **sequence in**  $A$  if  $a_n \in A$  for every  $n \in \mathbb{N}$ .

The following characterizations of closed sets and closure are very commonly used, particularly in calculus.

**Proposition VIII.7** (Characterization of closed sets by sequences).

*A subset  $A \subset X$  is closed if and only if for every sequence  $(a_n)_{n \in \mathbb{N}}$  in  $A$  which is convergent in  $X$ , the limit is in  $A \subset X$ .*

*Proof.* We prove “if” and “only if” separately.

*proof of “if”:* We will prove this by contrapositive. Assume that  $A \subset X$  is *not* closed, i.e., that its complement  $X \setminus A$  is not open. By Definition V.18 this means that for some point  $x \in X \setminus A$  of the complement, there exists no  $r > 0$  such that the ball  $\mathcal{B}_r(x)$  would be contained in the complement  $X \setminus A$ . In particular, for  $n \in \mathbb{N}$ , using  $r = \frac{1}{n}$  as the radius, we find that  $\mathcal{B}_{1/n}(x) \not\subset X \setminus A$ , i.e.,  $\mathcal{B}_{1/n}(x) \cap A \neq \emptyset$ . This means that for every  $n \in \mathbb{N}$  we can choose a point  $a_n \in \mathcal{B}_{1/n}(x) \cap A$ , i.e., a point  $a_n \in A$  such that  $d(x, a_n) < \frac{1}{n}$ . Such points form a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $A$ . Also  $0 \leq d(x, a_n) < \frac{1}{n}$  for all  $n \in \mathbb{N}$ , so by the squeeze theorem (Lemma II.17) we have  $\lim_{n \rightarrow \infty} d(x, a_n) = 0$ , which by Proposition VIII.4 implies  $\lim_{n \rightarrow \infty} a_n = x$ . But  $x \in X \setminus A$ , so we have found a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $A$  whose limit  $x$  is not in  $A$ . By contrapositive we have proven the “if” implication.

*proof of “only if”:* We will also prove this by contrapositive. Assume that there exists a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $A$  which is convergent, and its limit  $x = \lim_{n \rightarrow \infty} a_n \in X$  is not in  $A$ , i.e.,  $x \in X \setminus A$ . Now for any  $r > 0$ , since  $\mathcal{B}_r(x)$  is a neighborhood of  $x$ , by the characterization of limits in Proposition VIII.4 we have that all but finitely many members of the sequence  $(a_n)_{n \in \mathbb{N}}$  lie in  $\mathcal{B}_r(x)$ . In particular there exists some such member  $x_n \in \mathcal{B}_r(x)$ , and since  $a_n \in A$ , we get  $\mathcal{B}_r(x) \cap A \neq \emptyset$ , i.e.,  $\mathcal{B}_r(x) \not\subset X \setminus A$ . We have thus shown that no open ball centered at  $x$  is contained in the complement  $X \setminus A$  of  $A$ , so the complement  $X \setminus A$  is not open, and therefore  $A$  is not closed. By contrapositive we have proven the “only if” implication.  $\square$

**Proposition VIII.8** (Characterization of closure by sequences).

*For a subset  $A \subset X$  and a point  $x \in X$ , the following are equivalent:*

- (a)  $x \in \overline{A}$ ;
- (b) *there exists a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $A$  such that  $\lim_{n \rightarrow \infty} a_n = x$ .*

The proof is quite similar to that of Proposition VIII.7. We leave the details as an exercise.

**Exercise VIII.4** (Proof of Proposition VIII.7).

Prove Proposition VIII.7.

Hint: First observe that  $\overline{A} = X \setminus \text{ext}(A)$ . From this it is easy to see that  $x \in \overline{A}$  if and only if for every  $r > 0$  we have  $\mathcal{B}_r(x) \cap A \neq \emptyset$ .

The following is clear by combining Propositions VIII.8 and VIII.7.

**Corollary VIII.9** (A characterization of closed sets by closures).

*For a subset  $A \subset X$ , the following are equivalent:*

- (i)  $A \subset X$  is closed;
- (ii)  $\overline{A} \subset A$ ;
- (iii)  $\overline{A} = A$ .



## VIII.3. Characterization of continuity

We next give a characterization of continuity of functions using sequences and limits. This will show that continuity of real functions as defined in Chapter III is merely a special case of the notion of continuity of functions between metric spaces defined in Chapter VI.

**Proposition VIII.10** (Characterization of continuity at a point).

*Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces,  $f: X \rightarrow Y$  a function, and  $x \in X$  a point. Then the following are equivalent:*

- (i)  *$f$  is continuous at  $x$ ;*
- (ii) *For any sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  with limit  $\lim_{n \rightarrow \infty} x_n = x$ , the sequence  $(f(x_n))_{n \in \mathbb{N}}$  in  $Y$  has the limit  $\lim_{n \rightarrow \infty} f(x_n) = f(x)$ .*

*Proof.* We prove implications in both directions separately.

*proof of (i)  $\Rightarrow$  (ii):* Assume that  $f: X \rightarrow Y$  is continuous at  $x \in X$ . Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $X$  such that  $\lim_{n \rightarrow \infty} x_n = x$ . We must show that the sequence  $(f(x_n))_{n \in \mathbb{N}}$  in  $Y$  converges to  $f(x)$ . So let  $\varepsilon > 0$ . By continuity of  $f$  at  $x$ , there exists a  $\delta > 0$  such that when  $d_X(x, x') < \delta$ , we have  $d_Y(f(x), f(x')) < \varepsilon$ . Since  $\delta > 0$ , by definition of limits, there exists an  $n' \in \mathbb{N}$  such that for  $n \geq n'$  we have  $d_X(x, x_n) < \delta$ . Combining these, we get that for  $n \geq n'$  we have  $d_Y(f(x), f(x_n)) < \varepsilon$ . Since  $\varepsilon > 0$  was arbitrary, this shows that  $\lim_{n \rightarrow \infty} f(x_n) = f(x)$ , and the implication (i)  $\Rightarrow$  (ii) is thus proven.

*proof of (ii)  $\Rightarrow$  (i):* Let us prove this by contrapositive. Assume that  $f: X \rightarrow Y$  is *not* continuous at  $x$ . Then for some  $\varepsilon > 0$  there does not exist any  $\delta > 0$  such that  $f[\mathcal{B}_\delta(x)]$  would be contained in  $\mathcal{B}_\varepsilon(f(x))$ . In particular for each  $n \in \mathbb{N}$ , using  $\delta = \frac{1}{n}$ , we have that  $f[\mathcal{B}_{1/n}(x)] \not\subset \mathcal{B}_\varepsilon(f(x))$ , i.e., there exists some  $x_n \in \mathcal{B}_{1/n}(x)$  such that  $f(x_n) \notin \mathcal{B}_\varepsilon(f(x))$ . Now for a sequence  $(x_n)_{n \in \mathbb{N}}$  of such choices, we have  $0 \leq d_X(x, x_n) < \frac{1}{n}$ , so by the squeeze theorem we get  $\lim_{n \rightarrow \infty} d_X(x, x_n) = 0$  and therefore  $\lim_{n \rightarrow \infty} x_n = x$  (by Proposition VIII.4). On the other hand since  $f(x_n) \notin \mathcal{B}_\varepsilon(f(x))$ , we have  $d_Y(f(x), f(x_n)) \geq \varepsilon$  for all  $n \in \mathbb{N}$ . In particular there is no  $n_\varepsilon$  as required in the definition of limit for this  $\varepsilon > 0$  and candidate limit  $f(x)$ , so the sequence  $(f(x_n))_{n \in \mathbb{N}}$  does not converge to  $f(x)$ . By contrapositive, the existence of such a sequence  $(x_n)_{n \in \mathbb{N}}$  proves the implication (ii)  $\Rightarrow$  (i).  $\square$

**Corollary VIII.11** (Characterization of continuity using limits).

*Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces and  $f: X \rightarrow Y$  a function. Then the following are equivalent:*

- (i)  *$f: X \rightarrow Y$  is a continuous function.*
- (ii) *For every sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  which converges in  $X$ , we have that the sequence  $(f(x_n))_{n \in \mathbb{N}}$  converges in  $Y$  to the limit*

$$\lim_{n \rightarrow \infty} f(x_n) = f\left(\lim_{n \rightarrow \infty} x_n\right).$$

*Proof.* Since the continuity of a function  $f: X \rightarrow Y$  is defined as its continuity at every point  $x$  of its domain  $X$  (Definition VI.1), the assertion follows directly from Proposition VIII.10.  $\square$

From Proposition VIII.10 and Corollary VIII.11 it is now clear that continuity of real-valued functions defined on subsets of the real line as defined in Chapter III indeed coincides with the general metric space definition of continuity, specialized to the case of the metric space  $(\mathbb{R}, d_{\mathbb{R}})$  and its subspaces with their inherited metrics.

**Example VIII.12** (Evaluation is not continuous with respect to the  $L^1$ -norm).

Consider the space  $\mathcal{C}([0, 1])$  of all continuous functions  $f: [0, 1] \rightarrow \mathbb{R}$  with the  $L^1$ -norm  $\|f\|_1 = \int_0^1 |f(t)| dt$  and the metric induced by it. We will show that the evaluation at  $z \in [0, 1]$ ,

$$\text{ev}_z: \mathcal{C}([0, 1]) \rightarrow \mathbb{R} \qquad \text{ev}_z(f) = f(z) \quad \text{for } f \in \mathcal{C}([0, 1]),$$

is not continuous in this metric space (compare with Example VI.9, where the  $\|\cdot\|_\infty$ -norm was used instead). We will specifically treat the case  $z = 0$  below. Similar arguments could be used to show that evaluations at other points  $z \in [0, 1]$  are not continuous, either. For notational clarity we nevertheless often write  $z$  instead of 0 below, in particular to make a distinction from the constant function zero, also denoted by 0.

In Example VIII.6 we exhibited a sequence  $(f_n)_{n \in \mathbb{N}}$  in  $\mathcal{C}([0, 1])$  such that at  $z = 0$  we have  $f_n(z) = 1$  for every  $n \in \mathbb{N}$ , and  $\lim_{n \rightarrow \infty} f_n = 0 \in \mathcal{C}([0, 1])$  (the limit of the sequence is the zero function 0). Applying  $\text{ev}_z$  to this sequence and taking the limit, we find

$$\lim_{n \rightarrow \infty} \text{ev}_z(f_n) = \lim_{n \rightarrow \infty} f_n(z) = \lim_{n \rightarrow \infty} 1 = 1,$$

while on the other hand  $\text{ev}_z$  applied to the limit of the sequence gives

$$\text{ev}_z\left(\lim_{n \rightarrow \infty} f_n\right) = \text{ev}_z(0) = 0(z) = 0.$$

The two calculations together show that  $\lim_{n \rightarrow \infty} \text{ev}_z(f_n) \neq \text{ev}_z(\lim_{n \rightarrow \infty} f_n)$ , which by Corollary VIII.11 shows that  $\text{ev}_z: \mathcal{C}([0, 1]) \rightarrow \mathbb{R}$  is not continuous.

**Exercise VIII.5** (Taking the derivative is not continuous with respect to the uniform norm).

Let

$$\mathcal{C}^1([-1, 1]) = \{x: [-1, 1] \rightarrow \mathbb{R} \mid x' \text{ is continuous}\}$$

be the set of all continuously differentiable functions on the interval  $[-1, 1]$ . Interpret it as a subset  $\mathcal{C}^1([-1, 1]) \subset \mathcal{C}([-1, 1])$ , where the space  $\mathcal{C}([-1, 1])$  of continuous functions is equipped again with the metric induced by the norm  $\|\cdot\|_\infty$ . As in Exercise VI.13, consider the function  $g: \mathcal{C}^1([-1, 1]) \rightarrow \mathcal{C}([-1, 1])$  given by

$$[g(x)](t) = x'(t), \quad \text{for } x \in \mathcal{C}^1([-1, 1]) \text{ and } t \in [-1, 1].$$

Show that  $g: \mathcal{C}^1([-1, 1]) \rightarrow \mathcal{C}([-1, 1])$  is not continuous.

**Exercise VIII.6** (Coordinatewise convergence is not sufficient for convergence in  $\ell^1$ ).

Consider the space

$$\ell^1 = \left\{ x = (x_j)_{j \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}} \mid \sum_{j=1}^{\infty} |x_j| < \infty \right\}$$

of absolutely summable real sequences. We consider it known that the formula  $\|x\|_1 = \sum_{j=1}^{\infty} |x_j|$  for  $x = (x_j)_{j \in \mathbb{N}} \in \ell^1$  defines a norm on  $\ell^1$ . We equip  $\ell^1$  with the metric induced by the norm  $\|\cdot\|_1$ .

- (a) Show that if a sequence  $(x^{(n)})_{n \in \mathbb{N}}$  of elements  $x^{(n)} = (x_j^{(n)})_{j \in \mathbb{N}} \in \ell^1$  converges in  $\ell^1$  to  $x = (x_j)_{j \in \mathbb{N}}$ , then for every  $k \in \mathbb{N}$ , the sequence  $(x_k^{(n)})_{n \in \mathbb{N}}$  of the  $k$ :th coordinates of  $x^{(n)}$ 's converges to  $\lim_{n \rightarrow \infty} x_k^{(n)} = x_k$  (limit in  $\mathbb{R}$ ).

*Hint:* You can start by showing that the  $k$ :th coordinate projection function  $(x_j)_{j \in \mathbb{N}} \mapsto x_k$  is a 1-Lipschitz function  $\ell^1 \rightarrow \mathbb{R}$ .

- (b) For  $n \in \mathbb{N}$  let  $x^{(n)} = (x_j^{(n)})_{j \in \mathbb{N}} \in \ell^1$  be the element given by

$$x_j^{(n)} = \begin{cases} 1 & \text{if } j = n \\ 0 & \text{if } j \neq n. \end{cases}$$

Show that for any  $k \in \mathbb{N}$  we have  $\lim_{n \rightarrow \infty} x_k^{(n)} = 0$  but in the space  $(\ell^1, \|\cdot\|_1)$  the sequence  $(x^{(n)})_{n \in \mathbb{N}}$  does not converge.

*Hint:* If the sequence would converge in  $\ell^1$ , then part (a) together with the first calculation of (b) identifies the only possibility for a limit  $x \in \ell^1$ . Now show directly from the definition of limits that we do not have convergence to that candidate limit.

### Operations on continuous real-valued functions

Similarly to Proposition III.10, we can now show that continuity of real-valued functions on general metric spaces is preserved under taking pointwise sums, products, and quotients.

UNDER CONSTRUCTION!

### VIII.4. Sequences in product spaces

The convergence of sequences in product spaces can be characterized in terms of the “coordinate sequences” obtained by projections as follows. If  $(z_n)_{n \in \mathbb{N}}$  is a sequence in the product space  $X \times Y$  of two metric spaces  $X$  and  $Y$ , then we may, for  $n \in \mathbb{N}$ , write the member of the sequence as  $z_n = (x_n, y_n)$ , where

$$x_n = \text{pr}_1(z_n) \in X \quad \text{and} \quad y_n = \text{pr}_2(z_n) \in Y$$

are the projections. We thus obtain the “coordinate sequences”  $(x_n)_{n \in \mathbb{N}}$  in  $X$  and  $(y_n)_{n \in \mathbb{N}}$  in  $Y$  of the sequence  $(z_n)_{n \in \mathbb{N}} = ((x_n, y_n))_{n \in \mathbb{N}}$ . We will see below that the convergence and limit of the sequence  $(z_n)_{n \in \mathbb{N}}$  in the product space  $X \times Y$  can be characterized in terms of the convergence and limits of  $(x_n)_{n \in \mathbb{N}}$  in  $X$  and  $(y_n)_{n \in \mathbb{N}}$  in  $Y$ .

Notice first that convergence does not depend on which of the topologically equivalent metrics we equip the product space with.

**Corollary VIII.13** (Convergence of sequences only depends on the topology).

*If  $d_1$  and  $d_2$  are two topologically equivalent metrics on  $X$ , then a sequence  $(x_n)_{n \in \mathbb{N}}$  converges to  $x \in X$  in the metric space  $(X, d_1)$  if and only if it converges to  $x \in X$  in the metric space  $(X, d_2)$ .*

*Proof.* One of the characterizations of limits in Proposition VIII.4 only uses the notion of neighborhoods. In view of that, it is clear that if two metrics determine the same open sets, the limits of sequences are the same.  $\square$

**Proposition VIII.14** (Convergence of sequences in a product space).

*Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces, and  $X \times Y$  their product space equipped with any metric  $d$  which is topologically equivalent to those in (VII.1).*

*Consider a sequence  $((x_n, y_n))_{n \in \mathbb{N}}$  in the product space  $X \times Y$ . Then the following are equivalent:*

- (i) *The sequence  $((x_n, y_n))_{n \in \mathbb{N}}$  converges to  $(x, y) \in X \times Y$ .*
- (ii) *The coordinate sequences  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  converge in  $X$  and  $Y$ , respectively, to limits*

$$\lim_{n \rightarrow \infty} x_n = x \in X \quad \text{and} \quad \lim_{n \rightarrow \infty} y_n = y \in Y.$$

*Proof.* We must prove the implications in both directions.

*proof of (i)  $\Rightarrow$  (ii):* Assume that  $((x_n, y_n))_{n \in \mathbb{N}}$  converges to  $(x, y)$  in the product space  $X \times Y$ . The projections  $\text{pr}_1: X \times Y \rightarrow X$  and  $\text{pr}_2: X \times Y \rightarrow Y$  are continuous by Lemma VII.14.

Therefore by Proposition VIII.10 in  $X$  we have

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \text{pr}_1((x_n, y_n)) = \text{pr}_1\left(\lim_{n \rightarrow \infty} (x_n, y_n)\right) = \text{pr}_1((x, y)) = x.$$

Similarly we get  $\lim_{n \rightarrow \infty} y_n = y$  in  $Y$ .

*proof of (ii)  $\Rightarrow$  (i):* Assume that  $\lim_{n \rightarrow \infty} x_n = x \in X$  and  $\lim_{n \rightarrow \infty} y_n = y \in Y$ . We must prove that  $((x_n, y_n))_{n \in \mathbb{N}}$  converges to  $(x, y) \in X \times Y$ . Note that by Corollary VIII.13 this convergence does not depend on the specific choice of the metric, so we may assume that the metric  $d_1$  of (VII.2) is used.

Let  $\varepsilon > 0$ . Since the sequences  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  converge to  $x$  and  $y$ , respectively, there exists some  $n', n'' \in \mathbb{N}$  such that

$$\begin{aligned} d_X(x_n, x) &< \frac{\varepsilon}{2} && \text{for } n \geq n', \\ d_Y(y_n, y) &< \frac{\varepsilon}{2} && \text{for } n \geq n''. \end{aligned}$$

Then for  $n \geq n_\varepsilon = \max\{n', n''\}$ , we have

$$d_1((x_n, y_n), (x, y)) = \underbrace{d_X(x_n, x)}_{< \varepsilon/2} + \underbrace{d_Y(y_n, y)}_{< \varepsilon/2} < \varepsilon.$$

This shows the convergence  $\lim_{n \rightarrow \infty} (x_n, y_n) = (x, y)$  in  $X \times Y$ . □

## Function sequences

Let  $X$  and  $Y$  be two sets. If for every index  $n \in \mathbb{N}$  we are given a function

$$f_n: X \rightarrow Y,$$

then the sequence

$$(f_1, f_2, f_3, \dots) = (f_n)_{n \in \mathbb{N}}$$

is called a **function sequence**. Such sequences are important in various contexts, for example:

- *Function series:* In analysis one often deals with series representations of functions (for example Fourier series, power series, ...). To give a meaning to such an infinite series of functions, one must first consider, for each  $n \in \mathbb{N}$ , the partial sum of the first  $n$  terms. The sequence of the partial sums is a function sequence.
- *Numerical analysis:* In practice one often performs mathematical tasks (particularly on a computer) not exactly but numerically, to a good (but not perfect) numerical accuracy. If one is interested in the numerics of a function (for example solutions to partial differential equations, ...) and if the parameters of the numerical approximation can be tuned, then one can form a sequence of functions out of ever improving choices of the numerical accuracy parameters, and one has a sequence of functions.
- *Statistics and stochastics:* A random variable is (according to its mathematically idealized definition) a function defined on the “sample space”. In statistics and probability one often considers sequences of random variables (say time series, sequences indexed by increasing sample size, ...), which then are by definition function sequences.

We will discuss two different notions of convergence of function sequences: pointwise convergence and uniform convergence. Both of these refer to the convergence of the values of the functions in an appropriate sense, the codomain  $Y$  needs to be equipped with a metric. By contrast, on the domain  $X$  it is not necessary to have a metric or any other structure (although the domain may also naturally happen to have such structure).

### IX.1. Pointwise convergence and uniform convergence

Let  $X$  be a set and  $(Y, d_Y)$  a metric space. We consider sequences  $(f_n)_{n \in \mathbb{N}}$  of functions  $X \rightarrow Y$ . Observe that for any given  $x \in X$ , the sequence  $(f_n(x))_{n \in \mathbb{N}}$  of the values of these functions at the point  $x$  is a sequence in the metric space  $Y$ , so it is meaningful to speak of its convergence, as in Chapter VIII. The first notion of convergence of function sequences is very straightforward in view of this.

**Definition IX.1** (Pointwise convergence of a function sequence).

A sequence  $(f_n)_{n \in \mathbb{N}}$  of functions  $X \rightarrow Y$  **converges pointwise** to a function  $f: X \rightarrow Y$ , if

$$\text{for all } x \in X \text{ we have } \lim_{n \rightarrow \infty} f_n(x) = f(x) \in Y.$$

In this situation, we alternatively say that  $f_n$  **tends to  $f$  pointwise** as  $n \rightarrow \infty$ , or simply that  $f_n \rightarrow f$  pointwise as  $n \rightarrow \infty$ .

Pointwise convergence thus literally means that at every point of the domain, the values of the functions converge. If we unravel the definition of convergence of the values, we can rewrite pointwise convergence as follows.

**Remark IX.2** (Meaning of pointwise convergence as a logical statement).

In view of Definition VIII.1,  $f_n \rightarrow f$  pointwise as  $n \rightarrow \infty$  means that

$$\forall x \in X : \quad \forall \varepsilon > 0 : \quad \exists n' \in \mathbb{N} : \quad n \geq n' \implies d_Y(f_n(x), f(x)) < \varepsilon.$$

Using the characterization of limits from Proposition VIII.4,  $f_n \rightarrow f$  pointwise as  $n \rightarrow \infty$  is also equivalent to

$$\forall x \in X : \quad \lim_{n \rightarrow \infty} d_Y(f_n(x), f(x)) = 0.$$

The other notion of convergence of function sequences that we examine here is uniform convergence.

**Definition IX.3** (Uniform convergence of a function sequence).

A sequence  $(f_n)_{n \in \mathbb{N}}$  of functions  $X \rightarrow Y$  **converges uniformly** to a function  $f: X \rightarrow Y$ , if we have

$$\lim_{n \rightarrow \infty} \sup \left\{ d_Y(f_n(x), f(x)) \mid x \in X \right\} = 0.$$

In this situation, we alternatively say that  $f_n$  **tends to  $f$  uniformly** as  $n \rightarrow \infty$ , or simply that  $f_n \rightarrow f$  uniformly as  $n \rightarrow \infty$ .

In this course (and also very commonly otherwise) we most often consider real valued functions, i.e., we take the codomain to be  $Y = \mathbb{R}$  with its standard metric. In that case the condition for uniform convergence reads

$$\lim_{n \rightarrow \infty} \sup \left\{ |f_n(x) - f(x)| \mid x \in X \right\} = 0.$$

**Exercise IX.1** (Meaning of uniform convergence as a logical statement).

Prove that the statement  $f_n \rightarrow f$  uniformly as  $n \rightarrow \infty$  is logically equivalent to

$$\forall \varepsilon > 0 : \quad \exists n' \in \mathbb{N} : \quad \forall x \in X : \quad n \geq n' \implies d_Y(f_n(x), f(x)) < \varepsilon.$$

Compare Remark IX.2 and Exercise IX.1, and note the difference!

In pointwise convergence (Remark IX.2), the  $n'$  is allowed to depend on the point  $x \in X$  as well, whereas in uniform convergence (Exercise IX.1) the *same* choice of  $n'$  must work simultaneously for all  $x \in X$  (i.e., a *uniform* choice can be made for the whole space  $X$  at once). Once one internalizes this key difference between the two definitions, the following becomes obvious.

**Lemma IX.4** (Uniform convergence implies pointwise convergence).

If  $f_n \rightarrow f$  uniformly as  $n \rightarrow \infty$ , then we also have that  $f_n \rightarrow f$  pointwise as  $n \rightarrow \infty$ .

*Proof.* Suppose that  $f_n \rightarrow f$  uniformly as  $n \rightarrow \infty$ . To prove pointwise convergence, first fix a point  $x \in X$ . We must then show that  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ .

For the distance  $d_Y(f_n(x), f(x))$ , observe the obvious inequalities

$$0 \leq d_Y(f_n(x), f(x)) \leq \sup \left\{ d_Y(f_n(x'), f(x')) \mid x' \in X \right\} \xrightarrow{n \rightarrow \infty} 0,$$

where the right hand side tends to 0 as  $n \rightarrow \infty$  by virtue of the assumed uniform convergence  $f_n \rightarrow f$ . From these inequalities and the squeeze theorem (Lemma II.17), we get that  $\lim_{n \rightarrow \infty} d_Y(f_n(x), f(x)) = 0$ . By the characterization of limits in Proposition VIII.4, this implies  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ , as desired. Pointwise convergence  $f_n \rightarrow f$  as  $n \rightarrow \infty$  follows.  $\square$

In other words, Lemma IX.4 says that uniform convergence of a function sequence is “stronger” (a more stringent requirement) than pointwise convergence. One practical use of this is that if we want to show uniform convergence of a function sequence  $(f_n)_{n \in \mathbb{N}}$  to some limit function  $f$ , this limit function must also be the pointwise limit function. Therefore in order to find the right candidate for the limit function, we first simply calculate the pointwise limits  $\lim_{n \rightarrow \infty} f_n(x)$  at all points  $x$ . Examples IX.5, IX.6, and IX.7 below illustrate this use.

In fact strictly speaking Lemma IX.4 only says that uniform convergence is “at least as strong” as pointwise convergence — to conclude that it is a “strictly stronger” notion, we should show that the converse implication does not hold. That will, indeed, be seen in the following example.

**Example IX.5** (An example of pointwise but not uniform convergence).

Consider the following case, where the domain of the functions is taken to be the interval  $X = [0, 1]$ , and the codomain is taken to be the real axis  $Y = \mathbb{R}$  with its standard metric.

For  $n \in \mathbb{N}$ , define the function

$$f_n: [0, 1] \rightarrow \mathbb{R} \quad \text{by} \quad f_n(x) = x^n \quad \text{for } x \in \mathbb{N},$$

see Figure IX.1. Let us verify that this sequence converges pointwise. We handle the cases  $x = 1$  and  $x < 1$  separately.

Consider the case  $x = 1$ . The function values at this point are simply  $f_n(1) = 1^n = 1$  for any  $n \in \mathbb{N}$ . Obviously this constant sequence of values converges:

$$\lim_{n \rightarrow \infty} f_n(1) = \lim_{n \rightarrow \infty} 1^n = 1.$$

A pointwise limit function  $f$  must therefore assume the value  $f(1) = 1$  at the point  $x = 1$ .

Consider then the case  $x \in [0, 1)$ , i.e.,  $0 \leq x < 1$ . The function values at such a point are  $f_n(x) = x^n$  for  $n \in \mathbb{N}$ , and the calculation of the limit is easy

$$\lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} x^n = 0.$$

A pointwise limit function  $f$  must therefore assume the value  $f(x) = 0$  at any point  $x \in [0, 1)$ .

By the above, we have narrowed down the only possible candidate for a pointwise limit function as the function

$$f: [0, 1] \rightarrow \mathbb{R} \quad \text{given by} \quad f(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 \\ 1 & \text{if } x = 1. \end{cases} \quad (\text{IX.1})$$

In the two cases above we in fact verified that  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for any  $x \in [0, 1]$ , so indeed we have  $f_n \rightarrow f$  pointwise as  $n \rightarrow \infty$ .




FIGURE IX.1. The functions  $f_n: [0, 1] \rightarrow \mathbb{R}$  given by  $f_n(x) = x^n$ .

But does the function sequence  $(f_n)_{n \in \mathbb{N}}$  converge uniformly? If it does converge uniformly to some function  $\tilde{f}: [0, 1] \rightarrow \mathbb{R}$ , then by Lemma IX.4 it also converges pointwise to  $\tilde{f}$ . But since we showed that  $(f_n)_{n \in \mathbb{N}}$  converges pointwise to the function  $f$  given in (IX.1), so in this case we must have<sup>1</sup>  $\tilde{f} = f$ . Therefore the remaining question is: do we have  $f_n \rightarrow f$  uniformly as  $n \rightarrow \infty$ ?

We will show that  $(f_n)_{n \in \mathbb{N}}$  does *not* converge uniformly to  $f$ . According to Definition IX.3, we should show that the supremum

$$\sup \left\{ d_Y(f_n(x), f(x)) \mid x \in [0, 1] \right\}$$

is not tending to zero as  $n \rightarrow \infty$ . To get a lower bound for the supremum for a given  $n \in \mathbb{N}$ , it is sufficient to find some point at which  $d_Y(f_n(x), f(x))$  is not small. Consider for example the point

$$x_n = \sqrt[n]{1/2} = 2^{-1/n}.$$

Clearly  $0 < x_n < 1$ , so the value of the pointwise limit function (IX.1) at this point is  $f(x_n) = 0$ . On the other hand, the value of the  $n$ :th member  $f_n$  of the function sequence at this point is  $f_n(x_n) = (2^{-1/n})^n = 2^{-1} = 1/2$ . We get

$$\sup \left\{ d_Y(f_n(x), f(x)) \mid x \in [0, 1] \right\} \geq \frac{1}{2}.$$

(In fact a little bit more careful thinking would show that the supremum here is exactly 1, for any  $n \in \mathbb{N}$  — you should think about this as an exercise).

Since this lower bound was obtained for an arbitrary  $n \in \mathbb{N}$ , the supremum here cannot be tending to 0 as  $n \rightarrow \infty$ . Therefore the sequence  $(f_n)_{n \in \mathbb{N}}$  does not converge uniformly.

**Example IX.6** (An example of uniform convergence).

Consider then the following case, where the domain of the functions is taken to be the

<sup>1</sup>Here we are using the fact that a pointwise limit function is unique if it exists. This is an easy consequence of the uniqueness of limits of sequences in metric spaces (Lemma VIII.3), as the careful reader will quickly verify.



interval  $X = [-\frac{6}{7}, \frac{6}{7}]$ , and the codomain is taken to be the real axis  $Y = \mathbb{R}$  with its standard metric.

For  $n \in \mathbb{N}$ , define the function

$$f_n: \left[-\frac{6}{7}, \frac{6}{7}\right] \rightarrow \mathbb{R} \quad \text{by} \quad f_n(x) = x^n \quad \text{for } x \in \mathbb{N}.$$

The formula for  $f_n$  is the same as in Example IX.5, but the domain is different! In this case, we will verify that this sequence converges uniformly.

First fix  $n \in \mathbb{N}$ . Then observe that for any  $x \in [-\frac{6}{7}, \frac{6}{7}]$  we have the bound

$$|f_n(x)| = |x^n| = |x|^n \leq \left(\frac{6}{7}\right)^n.$$

Since this holds for any  $x$  in the domain, we get

$$\sup \left\{ |f_n(x)| \mid x \in \left[-\frac{6}{7}, \frac{6}{7}\right] \right\} \leq \left(\frac{6}{7}\right)^n$$

Let us define the zero function  $f(x) = 0$  for all  $x \in [-\frac{6}{7}, \frac{6}{7}]$ . Subtracting zero has no effect, so  $|f_n(x) - f(x)| = |f_n(x)|$  above, and we have obtained

$$\sup \left\{ |f_n(x) - f(x)| \mid x \in \left[-\frac{6}{7}, \frac{6}{7}\right] \right\} \leq \left(\frac{6}{7}\right)^n \xrightarrow{n \rightarrow \infty} 0,$$

where the right hand side is tending to zero as  $n \rightarrow \infty$ , because  $\frac{6}{7} < 1$ . This shows that we have uniform convergence  $f_n \rightarrow f$  as  $n \rightarrow \infty$ . By Lemma IX.4 uniform convergence implies also pointwise convergence  $f_n \rightarrow f$  as  $n \rightarrow \infty$ , to the same limit function  $f$  (the zero function).

**Exercise IX.2** (Visualizing the uniform convergence in an example).

Plot the fifteen or twenty first functions  $f_n$  in Example IX.6 to visualize the uniformly converging function sequence  $(f_n)_{n \in \mathbb{N}}$ . Compare with Example IX.5 and Figure IX.1.

**Example IX.7** (An example of neither type of convergence).

Consider then the following case, where the domain of the functions is taken to be the interval  $X = [-1, 1]$ , and the codomain is taken to be the real axis  $Y = \mathbb{R}$  with its standard metric.

For  $n \in \mathbb{N}$ , define the function

$$f_n: [-1, 1] \rightarrow \mathbb{R} \quad \text{by} \quad f_n(x) = x^n \quad \text{for } x \in \mathbb{N}.$$

The formula for  $f_n$  is the same as in Examples IX.5 and IX.6, but the domain is again different! In this case, we will verify that this sequence converges neither pointwise nor uniformly.

Indeed, consider the point  $x = -1$  in the domain  $[-1, 1]$ . The value of the  $n$ :th function at this point is  $f_n(-1) = (-1)^n$ , so the sequence  $(f_n(-1))_{n \in \mathbb{N}} = ((-1)^n)_{n \in \mathbb{N}}$  of values does not converge. Then we cannot have pointwise convergence  $f_n \rightarrow f$  to any function  $f$ . And by Lemma IX.4 it is consequently also impossible to have uniform convergence.

**Exercise IX.3** (A function sequence).

Define  $f_n: [0, 1] \rightarrow \mathbb{R}$  by setting

$$f_n(x) = n \sin(x/n), \quad \text{for } x \in [0, 1] \text{ and } n \in \mathbb{N}.$$

(a) Determine the limit function  $f: [0, 1] \rightarrow \mathbb{R}$  of the sequence  $(f_n)$ ,

$$f(x) = \lim_{n \rightarrow \infty} f_n(x), \quad x \in [0, 1].$$

(b) Does the sequence  $(f_n)$  converge uniformly on the interval  $[0, 1]$  to the function  $f$ ?

Hint: Either the 1:st order Taylor polynomial of the function  $\sin$  or L'Hospital's rule can be helpful.

**Exercise IX.4** (A different function sequence).

Define  $f_n: \mathbb{R} \rightarrow \mathbb{R}$  by setting

$$f_n(x) = n \sin(x/n), \quad \text{for } x \in \mathbb{R} \text{ and } n \in \mathbb{N}$$

(the formula is the same as in Exercise IX.3, but the domain is different!). Does this function sequence  $(f_n)_{n \in \mathbb{N}}$  converge pointwise? Does it converge uniformly?

## IX.2. Preservation of continuity in uniform limits

In Example IX.5 we got a pointwise limit function  $f$  that was discontinuous, although each member  $f_n$  of the sequence was continuous. In particular we see that, in general, continuity is not preserved under pointwise convergence! The situation with uniform convergence is better, and this is one of the main reasons why uniform convergence is so useful.

Of course, to talk about continuity of the functions  $X \rightarrow Y$ , also the domain  $X$  needs to have a metric<sup>2</sup>. In the following we therefore assume that  $(X, d_X)$  and  $(Y, d_Y)$  are metric spaces.

**Theorem IX.8** (Uniform limit of continuous functions is continuous).

*Suppose that  $(f_n)_{n \in \mathbb{N}}$  is a sequence of continuous functions  $f_n: X \rightarrow Y$  which converges uniformly to a limit function  $f: X \rightarrow Y$ . Then the limit function  $f$  is also continuous.*

*Proof.* We are assuming the hypotheses: that each  $f_n$  is continuous, and that  $f_n \rightarrow f$  uniformly.

We must prove the continuity of the limit function  $f: X \rightarrow Y$ . For this, let  $z \in X$  and let  $\varepsilon > 0$ . We must find a  $\delta > 0$  so that whenever  $d_X(z, x) < \delta$  we have  $d_Y(f(z), f(x)) < \varepsilon$ . By the assumed uniform convergence, since  $\frac{\varepsilon}{4} > 0$ , there exists an  $n_0 \in \mathbb{N}$  such that

$$\sup \{d_Y(f_n(x), f(x)) \mid x \in X\} < \frac{\varepsilon}{4} \quad \text{for all } n \geq n_0. \quad (\text{IX.2})$$

Also since  $f_{n_0}: X \rightarrow Y$  is (by assumption) continuous at  $z$  and  $\frac{\varepsilon}{2} > 0$ , there exists a  $\delta > 0$  such that

$$d_Y(f_{n_0}(z), f_{n_0}(x)) < \frac{\varepsilon}{2} \quad \text{whenever } d_X(z, x) < \delta. \quad (\text{IX.3})$$

We claim that this  $\delta$  will work for us. To estimate the distance from  $f(z)$  to  $f(x)$  in  $Y$ , we use triangle inequality twice for the four points  $f(z), f_{n_0}(z), f_{n_0}(x), f(x) \in Y$ . For any  $x \in X$  such that  $d_X(z, x) < \delta$  we thus get

$$\begin{aligned} & d_Y(f(z), f(x)) \\ & \leq \underbrace{d_Y(f(z), f_{n_0}(z))}_{< \varepsilon/4 \text{ by (IX.2)}} + \underbrace{d_Y(f_{n_0}(z), f_{n_0}(x))}_{< \varepsilon/2 \text{ by (IX.3)}} + \underbrace{d_Y(f_{n_0}(x), f(x))}_{< \varepsilon/4 \text{ by (IX.2)}} < \frac{\varepsilon}{4} + \frac{\varepsilon}{2} + \frac{\varepsilon}{4} = \varepsilon. \end{aligned}$$

Since for an arbitrary  $\varepsilon > 0$  we found a such  $\delta > 0$ , we obtain continuity of  $f$  at  $z$ . Since  $z \in X$  was arbitrary, this shows continuity of the function  $f: X \rightarrow Y$ .  $\square$

<sup>2</sup>... or at least a topology.

### IX.3. Function series

Let us now focus on the case of real-valued functions, i.e., taking the codomain to be  $Y = \mathbb{R}$  (with its standard metric). The domain can be an arbitrary set  $X$ . The addition, multiplication, and division in  $\mathbb{R}$  allow us to define the pointwise sums, products, and quotients of functions  $X \rightarrow \mathbb{R}$  (for quotients we only need the denominator to be non-zero).

Let  $(g_k)_{k \in \mathbb{N}}$  be a sequence of functions  $g_k: X \rightarrow \mathbb{R}$ . Pointwise addition of the values allows us to define, for any  $n \in \mathbb{N}$ ,

$$f_n: X \rightarrow \mathbb{R} \qquad f_n(x) = \sum_{k=1}^n g_k(x) \quad \text{for } x \in X \qquad (\text{IX.4})$$

This is the  $n$ :th partial sum of the **function series**  $\sum_{k=1}^{\infty} g_k$ . We say that this function series **converges** to  $\varsigma: X \rightarrow \mathbb{R}$  **pointwise** or **uniformly** if the sequence  $(f_n)_{n \in \mathbb{N}}$  of its partial sums converges to  $\varsigma$  pointwise or uniformly, respectively.

The following result, known as **Weierstrass'  $M$  test** is an often practical criterion by which one can verify uniform convergence of a function series, and to prove the continuity of the function represented by the series.

**Exercise IX.5** (Weierstrass'  $M$  test).

Let  $I \subset \mathbb{R}$  be an interval and  $g_k: I \rightarrow \mathbb{R}$  continuous functions, for  $k \in \mathbb{N}$ . Assume that  $|g_k(x)| \leq M_k$  for all  $x \in I$  and that the series  $\sum_{k=1}^{\infty} M_k$  is convergent. Prove that the infinite series

$$f(x) = \sum_{k=1}^{\infty} g_k(x),$$

defines a continuous function  $f: I \rightarrow \mathbb{R}$ .

*Hint: Uniform convergence of the partial sums and a result concerning uniform convergence, or a direct argument of the same type. The continuity of the partial sums (finitely many terms) is considered known. The remainder of a convergent series tends to zero as the starting index increases.*

Here is a typical application of the Weierstrass'  $M$ -test.

**Exercise IX.6** (A Fourier series application of the Weierstrass  $M$ -test).

Using Exercise IX.5, prove that the Fourier series

$$\sum_{k=1}^{\infty} \frac{4(-1)^k \cos(kx)}{k^2}$$

defines a continuous function on  $\mathbb{R}$ .

*Remark:* Apart from an additive constant, this is the Fourier series of the  $2\pi$ -periodic extension of  $f(x) = x^2$ ,  $-\pi \leq x \leq \pi$ , see Figure IX.2. This can be used to calculate the sum of the series  $\sum_{k=1}^{\infty} \frac{(-1)^k}{k^2}$  (by first determining the additive constant and then substituting  $x = 0$ ).

♥ **Power series**

UNDER CONSTRUCTION!




FIGURE IX.2. The continuous function represented by the Fourier series  $\sum_{k=1}^{\infty} \frac{4(-1)^k \cos(kx)}{k^2}$  of Exercise IX.6 is a  $2\pi$ -periodic extension of a parabola.

### ♥ Fourier series

UNDER CONSTRUCTION!

### ♥ Weierstrass' function

The function  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by

$$f(x) = \sum_{k=1}^{\infty} 2^{-k} \sin(4^k x) \quad \text{for } x \in \mathbb{R}$$

is known as the Weierstrass' function, see Figure IX.3 for a plot. Since  $|\sin(\theta)| \leq 1$  for any  $\theta \in \mathbb{R}$ , it is easy to apply the Weierstrass'  $M$ -test (Exercise IX.5) with the convergent geometric series  $\sum_{k=1}^{\infty} 2^{-k} = 1 < \infty$  to show that the series is uniformly convergent and determines a continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$ . Furthermore, it is possible to show that the function is nowhere differentiable, i.e., it does not have a derivative  $f'(x)$  at any point  $x \in \mathbb{R}$ .<sup>3</sup>

**Exercise (✓) IX.7** (Continuity of the Weierstrass' function).

Apply the Weierstrass'  $M$ -test (Exercise IX.5) to show that the Weierstrass' function given by  $f(x) = \sum_{k=1}^{\infty} 2^{-k} \sin(4^k x)$  is continuous  $\mathbb{R} \rightarrow \mathbb{R}$ .

**Exercise IX.8** (Non-differentiability of the Weierstrass' function at the origin).

Consider the Weierstrass' function given by  $f(x) = \sum_{k=1}^{\infty} 2^{-k} \sin(4^k x)$  for  $x \in \mathbb{R}$ .

- Calculate  $f(0)$ .
- For  $n \in \mathbb{N}$ , let  $x_n = \pi 4^{-n}$ . Show that  $f(x_n) \geq \frac{1}{\sqrt{2}} 2^{-n}$ .
- By considering the difference quotients  $\frac{f(x_n) - f(0)}{x_n - 0}$ , show that  $f$  has no derivative at 0.

**Exercise (✂) IX.9** (Nowhere differentiability of the Weierstrass' function).

Prove that the Weierstrass' function is not differentiable at any point  $x \in \mathbb{R}$ .

<sup>3</sup>Besides just looking at the plot in Figure IX.3, you can start to see why the function might be non-differentiable by observing the following. If you differentiate the series term by term, you formally obtain  $f'(x) = \sum_{k=1}^{\infty} 2^k \cos(4^k x)$ . This series does not converge at any point  $x \in \mathbb{R}$  (the terms do not even tend to zero). While this is not yet a precise proof of non-differentiability, it gives an idea of the problem. Exercise IX.8 indicates how to get started with the precise proof.




FIGURE IX.3. The continuous function represented by the Fourier series  $\sum_{k=1}^{\infty} 2^{-k} \sin(4^k x)$ . This function is continuous, but nowhere differentiable.

#### IX.4. ♥ Uniform convergence and integration

Another major advantage of uniform convergence is that it commutes with integration in the following sense.

**Theorem IX.9** (Integrals of uniform limits).

Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence of Riemann-integrable functions  $f_n: [a, b] \rightarrow \mathbb{R}$  which converges uniformly to a Riemann-integrable function  $f: [a, b] \rightarrow \mathbb{R}$ . Then we have

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) \, dx = \int_a^b \left( \lim_{n \rightarrow \infty} f_n(x) \right) \, dx.$$

*Sketch of proof.* Denote

$$s_n = \sup_{x \in [a, b]} |f_n(x) - f(x)|.$$

By uniform convergence  $f_n \rightarrow f$ , we have  $s_n \rightarrow 0$  as  $n \rightarrow \infty$ . For any  $x \in [a, b]$  and  $n \in \mathbb{N}$ , with the triangle inequality we get

$$f(x) - s_n \leq f_n(x) \leq f(x) + s_n.$$

Integrating over  $x$  and using monotonicity of integrals, we then get

$$\int_a^b f(x) \, dx - s_n(b-a) \leq \int_a^b f_n(x) \, dx \leq \int_a^b f(x) \, dx + s_n(b-a).$$

Here  $s_n(b-a) \rightarrow 0$  as  $n \rightarrow \infty$ , so by the squeeze theorem we obtain

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) \, dx = \int_a^b f(x) \, dx,$$

which is the asserted result.  $\square$

The same does not hold for pointwise convergence.

**Exercise IX.10** (A function sequence).

Consider the sequence  $(f_n)$  of functions  $f_n: [0, \infty) \rightarrow \mathbb{R}$  given by

$$f_n(x) = \begin{cases} x - n + 1, & \text{for } n - 1 \leq x \leq n, \\ n + 1 - x, & \text{for } n \leq x \leq n + 1, \\ 0, & \text{otherwise.} \end{cases}$$

- (a) Sketch the graphs of a few of the functions. Determine the pointwise limit function  $f$  of the function sequence  $(f_n)_{n \in \mathbb{N}}$ . (After guessing the limit with the help of the sketches, prove pointwise convergence from the definition).
- (b) Does the function sequence  $(f_n)_{n \in \mathbb{N}}$  converge uniformly on  $[0, \infty)$  to the limit function  $f$ ?
- (c) Do we have

$$\lim_{n \rightarrow \infty} \int_0^\infty f_n(x) \, dx = \int_0^\infty f(x) \, dx ?$$

- (d) By a minor modification one obtains a similar sequence  $(\tilde{f}_n)_{n \in \mathbb{N}}$  of functions which has the same limit function  $f$ , but the integrals of  $\tilde{f}_n$  tend to  $+\infty$  as  $n \rightarrow \infty$ . Find such a modification (an illustration suffices).

## Lecture X

# Completeness

### X.1. Cauchy sequences

Let  $(X, d)$  be a metric space.

Recall, from Definition VIII.1, that a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  is said to converge to a limit  $x \in X$  if for all  $\varepsilon > 0$  there exists a  $n_\varepsilon \in \mathbb{N}$  such that for  $n \geq n_\varepsilon$  we have  $d(x_n, x) < \varepsilon$ . Now note that according to this definition, *before* we can check this criterion of the convergence of the sequence  $(x_n)_{n \in \mathbb{N}}$ , we *already* have to know what the limit  $x \in X$  should be! Without a doubt, it would often be more convenient, if we could decide about the convergence of the sequence by looking only at the members of the sequence themselves. Cauchy sequences provide such a criterion, and to some extent it answers questions about convergence already.

**Definition X.1** (Cauchy sequence).

A sequence  $(x_n)_{n \in \mathbb{N}}$  in a metric space  $(X, d)$  is said to be **Cauchy** if for any  $\varepsilon > 0$  there exists an  $n_\varepsilon \in \mathbb{N}$  such that

$$d(x_k, x_\ell) < \varepsilon \quad \text{for all } k, \ell \geq n_\varepsilon. \quad (\text{X.1})$$

In other words, a sequence is Cauchy, if we can make the mutual distances among the members of the sequence arbitrarily small, by going far enough to the tail of the sequence.

#### *Cauchy sequences and convergent sequences*

Roughly, the idea of Definition X.1 is that a “Cauchy sequence looks (internally) like a convergent sequence”. As a first observation in support of such an intuitive interpretation we show that a convergent sequence must be Cauchy (so “actually convergent sequences also internally look convergent”).

**Lemma X.2** (Every convergent sequence is Cauchy).

*If a sequence  $(x_n)_{n \in \mathbb{N}}$  in a metric space  $X$  is convergent, then it is Cauchy.*

*Proof.* Suppose that  $(x_n)_{n \in \mathbb{N}}$  converges in  $X$ , and denote its limit by  $x = \lim_{n \rightarrow \infty} x_n \in X$ . Let  $\varepsilon > 0$ . By the assumed convergence, there exists  $n' \in \mathbb{N}$  such that

$$d(x_n, x) < \frac{\varepsilon}{2} \quad \text{for all } n \geq n'.$$

Then if  $k, \ell \geq n'$ , we have by triangle inequality

$$d(x_k, x_\ell) \leq \underbrace{d(x_k, x)}_{< \varepsilon/2} + \underbrace{d(x, x_\ell)}_{< \varepsilon/2} < \varepsilon.$$

This shows that the sequence  $(x_n)_{n \in \mathbb{N}}$  is Cauchy. □

The converge implication does not hold generally. Phrased intuitively again: there are metric spaces with sequences which “have the (internal) appearance of being convergent but which fail to actually converge within the space”.

**Example X.3** (Not every Cauchy sequence is convergent).

Consider the punctured real line  $X = \mathbb{R} \setminus \{0\} \subset \mathbb{R}$  with the metric  $d$  that it inherits from the whole real line  $\mathbb{R}$ . Consider the sequence  $(x_n)_{n \in \mathbb{N}}$  in  $\mathbb{R} \setminus \{0\}$  given by  $x_n = \frac{1}{n}$  for  $n \in \mathbb{N}$ . For any  $\varepsilon > 0$  we may set  $n_\varepsilon = \lceil 1/\varepsilon \rceil$ , and then for any  $k, \ell \geq n_\varepsilon$  we have  $d(x_k, x_\ell) = \left| \frac{1}{k} - \frac{1}{\ell} \right| < \frac{1}{\min\{k, \ell\}} \leq \frac{1}{n_\varepsilon} \leq \varepsilon$ . This shows that the sequence  $(x_n)_{n \in \mathbb{N}}$  in  $\mathbb{R} \setminus \{0\}$  is Cauchy. However, it is clear that it does not converge in  $\mathbb{R} \setminus \{0\}$  (why?).

### *Cauchy sequences and bounded sequences*

Above we saw that in general a Cauchy sequence may fail to be actually convergent, although it “appears (internally) convergent”. Next we check that at least a Cauchy sequence must be bounded.

**Lemma X.4** (Every Cauchy sequence is bounded).

*If a sequence  $(x_n)_{n \in \mathbb{N}}$  in a metric space  $X$  is Cauchy, then it is bounded.*

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $X$ . To show that  $(x_n)_{n \in \mathbb{N}}$  is bounded, we will find an  $R > 0$  such that  $x_n \in \mathcal{B}_R(x_1) \subset X$  for all  $n \in \mathbb{N}$ .

Let us fix  $\varepsilon = 1$ .<sup>1</sup> Using the Cauchy property (X.1) with this  $\varepsilon = 1 > 0$ , we get that there exists an  $n_\varepsilon \in \mathbb{N}$  such that  $d(x_k, x_\ell) < \varepsilon$  for all  $k, \ell \geq n_\varepsilon$ . Among the finitely many first indices  $n \leq n_\varepsilon$ , we can choose the one which maximizes distance to  $x_1$  — we now set

$$R' = \max \{d(x_1, x_n) \mid n \leq n_\varepsilon\} \quad \text{and} \quad R = R' + \varepsilon.$$

Because of the maximum above, for any  $n \leq n_\varepsilon$  we clearly have  $d(x_1, x_n) \leq R' < R$ , so  $x_n \in \mathcal{B}_R(x_1)$ . It remains to consider  $n > n_\varepsilon$ . But for  $n > n_\varepsilon$  we may use the triangle inequality and the choices of  $n_\varepsilon$  and  $R'$  to get

$$d(x_1, x_n) \leq \underbrace{d(x_1, x_{n_\varepsilon})}_{\leq R'} + \underbrace{d(x_{n_\varepsilon}, x_n)}_{< \varepsilon} < R' + \varepsilon = R$$

We have thus showed that  $x_n \in \mathcal{B}_R(x_1)$  for all  $n \in \mathbb{N}$ , so that the sequence  $(x_n)_{n \in \mathbb{N}}$  is indeed bounded (recall Lemma V.16).  $\square$

In this case it is hardly surprising that the converse implication does not hold.

**Exercise X.1** (Not every bounded sequence is Cauchy).


Find an example of a sequence in some metric space, which is bounded but is not Cauchy.

<sup>1</sup>Any positive number  $\varepsilon$  would work just as well — what is important is that one choice is fixed. For the convenience of the reader who prefers to choose  $\varepsilon = 42$  or  $\varepsilon = \frac{1}{361}$  or some other value instead, we keep referring to this *fixed* value simply by  $\varepsilon$  in the remaining proof (this should also help keep track of the origin of the different terms).



*Logical relationships among three different properties of sequences*

We may summarize general logical implications (and lack thereof) between three properties of sequences in metric spaces by the following diagram:

*A criterion for the convergence of a Cauchy sequence*

A Cauchy sequence “appears convergent”, and it turns out that if just any subsequence has a limit, then the whole sequence is in fact convergent.

**Lemma X.5** (Cauchy sequence with convergent subsequence is convergent).

Let  $(x_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $X$ . If there exists a subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  which converges, then the whole sequence  $(x_n)_{n \in \mathbb{N}}$  also converges to the same limit,

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} x_{\varphi(n)}.$$

*Proof.* Suppose that  $(x_n)_{n \in \mathbb{N}}$  is Cauchy and  $(x_{\varphi(n)})_{n \in \mathbb{N}}$  is a convergent subsequence. Denote the limit of the subsequence by  $\xi = \lim_{n \rightarrow \infty} x_{\varphi(n)} \in X$ .

We must prove that  $(x_n)_{n \in \mathbb{N}}$  also converges to  $\xi$ . So let  $\varepsilon > 0$ . By the Cauchy property of  $(x_n)_{n \in \mathbb{N}}$ , we know that there exists some  $n' \in \mathbb{N}$  such that

$$d(x_k, x_\ell) < \frac{\varepsilon}{2} \quad \text{for } k, \ell \geq n'. \quad (\text{X.2})$$

On the other hand, by the convergence of the subsequence  $(x_{\varphi(n)})_{n \in \mathbb{N}}$ , we know that there exists an  $m' \in \mathbb{N}$  such that

$$d(x_{\varphi(m)}, \xi) < \frac{\varepsilon}{2} \quad \text{for } m \geq m'. \quad (\text{X.3})$$

Let us select some  $m_0 \geq m'$  such that also  $\varphi(m_0) \geq n'$  (possible, since there are arbitrarily large subsequence indices), and denote  $n_0 = \varphi(m_0)$ . With this choice we have  $d(x_{n_0}, \xi) < \varepsilon/2$  by (X.3). Since  $n_0 \geq n'$ , by (X.2) we have for any  $n \geq n'$  that  $d(x_n, x_{n_0}) < \varepsilon/2$ . By triangle inequality, for  $n \geq n'$  we get

$$d(x_n, \xi) \leq \underbrace{d(x_n, x_{n_0})}_{< \varepsilon/2} + \underbrace{d(x_{n_0}, \xi)}_{< \varepsilon/2} < \varepsilon.$$

This shows the convergence of  $(x_n)_{n \in \mathbb{N}}$  to  $\xi$ . □

*Preservation of Cauchy sequences under Lipschitz functions*

Lipschitz images (and more generally uniformly continuous images) of Cauchy sequences are Cauchy.

**Lemma X.6** (Lipschitz maps preserve the Cauchy property).

Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces and  $f: X \rightarrow Y$  a Lipschitz function. Then if  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence in  $X$ , we have that  $(f(x_n))_{n \in \mathbb{N}}$  is a Cauchy sequence in  $Y$ .

*Proof.* Let  $K > 0$  be a Lipschitz constant for the Lipschitz function  $f: X \rightarrow Y$ . Suppose that  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence in  $X$ , and consider the image sequence  $(f(x_n))_{n \in \mathbb{N}}$  in  $Y$ .

Let  $\varepsilon > 0$ . Since  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy, there exists an  $n' \in \mathbb{N}$  such that  $d_X(x_k, x_\ell) < \frac{\varepsilon}{K}$  whenever  $k, \ell \geq n'$ . By the  $K$ -Lipchitz property, we then get for  $k, \ell \geq n'$  that

$$d_Y(f(x_k), f(x_\ell)) \leq K \underbrace{d_X(x_k, x_\ell)}_{< \varepsilon/K} < \varepsilon.$$

This shows that  $(f(x_n))_{n \in \mathbb{N}}$  is Cauchy. □

## X.2. Completeness

### X.2.1. Definition of completeness

**Definition X.7** (Complete metric space).

A metric space  $X$  is **complete** if every Cauchy sequence in  $X$  converges in  $X$ .

We can readily give an example of a space that is *not* complete.

**Example X.8** (The punctured real axis is not complete).

In Example X.3 we saw that the punctured real axis  $X = \mathbb{R} \setminus \{0\}$  (equipped with the metric inherited from the real axis) has a Cauchy sequence which is not convergent. Therefore  $\mathbb{R} \setminus \{0\}$  is not a complete metric space.

A slightly more interesting counterexample is the following.

**Example X.9** (The set of rational numbers is not complete).

Consider the set  $\mathbb{Q}$  of rational numbers, with the metric  $d_{\mathbb{Q}}$  it inherits from the real line as a subset  $\mathbb{Q} \subset \mathbb{R}$ , i.e.  $d_{\mathbb{Q}}(q_1, q_2) = |q_2 - q_1|$  for  $q_1, q_2 \in \mathbb{Q}$ . We claim that the metric space  $(\mathbb{Q}, d_{\mathbb{Q}})$  is not complete. To see this, it suffices to find a Cauchy sequence in  $\mathbb{Q}$  which does not converge in  $\mathbb{Q}$ .

Let us form a sequence  $(q_n)_{n \in \mathbb{N}}$  whose  $n$ :th member  $q_n$  is the truncated decimal expansion of  $\sqrt{2}$  with  $n$  decimals included, i.e.,

$$q_1 = 1.4, \quad q_2 = 1.41, \quad q_3 = 1.414, \quad q_4 = 1.4142, \quad q_5 = 1.41421, \quad \dots,$$

or more formally  $q_n = \lfloor 10^n \sqrt{2} \rfloor 10^{-n}$  for  $n \in \mathbb{N}$ . Each of these truncated decimal expansions is a rational number,  $q_n \in \mathbb{Q}$ , so  $(q_n)_{n \in \mathbb{N}}$  is a sequence in  $\mathbb{Q}$ . Let us check directly that this sequence is Cauchy.<sup>2</sup> For  $k, \ell \geq n$  we have that  $q_k$  and  $q_\ell$  may only differ in decimals after the  $n$ :th one, so as in Example II.33 we get  $|q_k - q_\ell| \leq \sum_{j=n+1}^{\infty} 9 \times 10^{-j} = 10^{-n}$ . Therefore for a given  $\varepsilon > 0$  it suffices to set  $n_\varepsilon = \lceil \log_{10}(1/\varepsilon) \rceil$  to ensure that for all  $k, \ell \geq n_\varepsilon$  we have  $|q_k - q_\ell| < \varepsilon$ .

Viewed as a sequence in  $\mathbb{R}$ , the sequence  $(q_n)_{n \in \mathbb{N}}$  converges to  $\sqrt{2}$ . It is therefore impossible that  $(q_n)_{n \in \mathbb{N}}$  converges in  $\mathbb{Q} \subset \mathbb{R}$ , because if it would, its limit  $q \in \mathbb{Q}$  would by the uniqueness of limits (Lemma VIII.3) in  $\mathbb{R}$  be  $q = \sqrt{2} \notin \mathbb{Q}$ ; a contradiction. Therefore the Cauchy sequence  $(q_n)_{n \in \mathbb{N}}$  in  $\mathbb{Q}$  does not converge in  $\mathbb{Q}$ , and the metric space  $(\mathbb{Q}, d_{\mathbb{Q}})$  is not complete.

---

<sup>2</sup>Instead of the direct proof, it is possible to note the following. Since  $\mathbb{Q} \subset \mathbb{R}$ , the sequence  $(q_n)_{n \in \mathbb{N}}$  may also be viewed as a sequence in  $\mathbb{R}$ . It is known to converge to  $\sqrt{2}$  in  $\mathbb{R}$  (recall Example II.33), so by Lemma ?? it has to be a Cauchy sequence in  $\mathbb{R}$ . But since the metric  $d_{\mathbb{Q}}$  we are using in  $\mathbb{Q}$  is the one inherited from  $\mathbb{R}$ , it is therefore also Cauchy in  $\mathbb{Q}$ .

### X.2.2. Completeness of the real line

The real axis is our simplest (and in some sense the most important) example of a complete metric space. It is worth noting that the completeness of  $\mathbb{R}$  as a metric space is in fact intimately related to the completeness axiom of  $\mathbb{R}$  (think about which part of the the proof below relies on the completeness axiom of  $\mathbb{R}$ ).

**Theorem X.10** (The real line is a complete metric space).

*The real line with its standard metric, i.e.  $(\mathbb{R}, d_{\mathbb{R}})$ , is a complete metric space.*

*Proof.* According to Definition X.7, we must show that every Cauchy sequence in  $\mathbb{R}$  converges.

So let  $(x_n)_{n \in \mathbb{N}}$  be a Cauchy sequence. By Lemma X.4, the sequence  $(x_n)_{n \in \mathbb{N}}$  must then be bounded. By Theorem III.6, the bounded real number sequence  $(x_n)_{n \in \mathbb{N}}$  has a convergent subsequence. By Lemma X.5, the Cauchy sequence  $(x_n)_{n \in \mathbb{N}}$  which has a convergent subsequence must itself be convergent. This finishes the proof.  $\square$

Completeness is *not* a purely topological property — it genuinely depends on the metric. This is seen by the following.

**Exercise X.2** (Completeness depends on metric).

Consider the open interval  $(-\frac{\pi}{2}, \frac{\pi}{2}) \subset \mathbb{R}$  with the metric it inherits from the real line. Recall that the function

$$\tan: \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$$

is a homeomorphism, and therefore on  $\mathbb{R}$  the metric defined by

$$\tilde{d}_{\mathbb{R}}(x, y) = |\tan^{-1}(y) - \tan^{-1}(x)| \quad \text{for } x, y \in \mathbb{R}$$

is equivalent to the usual metric (i.e., it induces the same topology).

Show that the metric space  $(\mathbb{R}, \tilde{d}_{\mathbb{R}})$  is not complete, although it is homeomorphic to the complete metric space  $(\mathbb{R}, d_{\mathbb{R}})$ .

### X.2.3. Important complete spaces

Let us now look at a few other important complete metric spaces. Note that each of the proofs of completeness below ultimately uses the completeness of  $\mathbb{R}$ .

*Euclidean spaces are complete*

**Theorem X.11** (The Euclidean spaces  $\mathbb{R}^d$  are complete).

*For any  $d \in \mathbb{N}$ , the Euclidean space  $\mathbb{R}^d$  equipped with its usual metric is a complete metric space.*

*Proof.* Let us only prove the case  $d = 2$  and leave the general case as an exercise.

Let  $(v_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $\mathbb{R}^2$  (equipped with the Euclidean metric). For each member  $v_n \in \mathbb{R}^2$  of the sequence, let us denote the projections to the two coordinates by  $x_n = \text{pr}_1(v_n) \in \mathbb{R}$  and  $y_n = \text{pr}_2(v_n) \in \mathbb{R}$ , so that

$$v_n = (x_n, y_n) \in \mathbb{R}^2.$$

Then  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  are sequences in  $\mathbb{R}$ . Since the projections  $\text{pr}_1, \text{pr}_2: \mathbb{R}^2 \rightarrow \mathbb{R}$  are Lipschitz, it follows from Lemma X.6 that  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  are Cauchy sequences.

Since  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  are Cauchy sequences in the complete metric space  $\mathbb{R}$ , they converge to some limits

$$x = \lim_{n \rightarrow \infty} x_n, \quad y = \lim_{n \rightarrow \infty} y_n.$$

By Proposition VIII.14, then, the original sequence  $(v_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}^2$  converges to the point  $v = (x, y) \in \mathbb{R}$  whose coordinates are the above limits of the coordinate sequences.  $\square$

**Exercise X.3** (The remaining part of the proof of Theorem X.11).

Complete the proof of Theorem X.11 by adapting the argument above so that it works in  $d > 2$  as well.

**Exercise X.4** (Completeness is preserved under bilipschitz-equivalence).

Show that ... **UNDER CONSTRUCTION!**

**Exercise X.5** (The space  $\mathbb{R}^d$  is complete with any norm induced metric).

Use Exercise X.4 to show that ... **UNDER CONSTRUCTION!**

### *A complete space of continuous functions*

The following complete metric space is very important in many applications.

**Theorem X.12** (A complete space of continuous functions).

*Consider the space*

$$\mathcal{C}([a, b]) = \left\{ f: [a, b] \rightarrow \mathbb{R} \mid f \text{ is continuous} \right\}$$

*of continuous functions on a closed interval  $[a, b] \subset \mathbb{R}$ , equipped with the metric  $d_\infty$  induced by the supremum-norm  $\|\cdot\|_\infty$  (see Example IV.27). Then  $(\mathcal{C}([a, b]), d_\infty)$  is a complete metric space.*

Let us immediately point out that the conclusion of Theorem X.12 depends crucially on the fact that we equip it with the supremum norm. We have seen also two other norms on  $\mathcal{C}([a, b])$ : the norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$  of Examples IV.21 and IV.15. The space  $\mathcal{C}([a, b])$  equipped with the metrics induced by these are *not* complete!

**Exercise X.6** (Incomplete spaces of continuous functions).

- (a) Prove that the space  $\mathcal{C}([a, b])$  equipped with the metric  $d_1$  induced by the norm  $\|\cdot\|_1$  of Example IV.21 is not complete.
- (b) Prove that the space  $\mathcal{C}([a, b])$  equipped with the metric  $d_2$  induced by the norm  $\|\cdot\|_2$  of Example IV.15 is not complete.

*Proof of Theorem X.12.* Let  $(f_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $\mathcal{C}([a, b])$  equipped with the norm

$$\|f\|_\infty = \sup \{|f(x)| \mid x \in [a, b]\}$$

and the corresponding metric

$$d_\infty(f, g) = \|g - f\|_\infty.$$

First recall from Example VI.9 that for any point  $x \in [a, b]$ , the evaluation mapping  $f \mapsto f(x) = \text{ev}_x(f)$  is 1-Lipschitz  $\mathcal{C}([a, b]) \rightarrow \mathbb{R}$ . Therefore by Lemma X.6, the real number sequence  $(\text{ev}_x(f_n))_{n \in \mathbb{N}} = (f_n(x))_{n \in \mathbb{N}}$  is also Cauchy. By completeness of  $\mathbb{R}$  (Theorem X.10), the Cauchy sequence  $(f_n(x))_{n \in \mathbb{N}}$  on  $\mathbb{R}$  converges. Let us denote its limit by

$$f(x) := \lim_{n \rightarrow \infty} f_n(x).$$

The numbers  $f(x) \in \mathbb{R}$  thus obtained for different  $x \in [a, b]$  define a function  $f: [a, b] \rightarrow \mathbb{R}$ . From above, we see that the sequence  $(f_n)_{n \in \mathbb{N}}$  of functions converges (at least) pointwise to the function  $f$ . We will show that the convergence is in fact uniform.

Let  $\varepsilon > 0$ . By the Cauchy property of the sequence  $(f_n)_{n \in \mathbb{N}}$ , there exists an  $n' \in \mathbb{N}$  such that

$$\|f_k - f_\ell\|_\infty < \frac{\varepsilon}{2} \quad \text{whenever} \quad k, \ell \geq n'.$$

This implies that for any  $x \in [a, b]$  and any  $k, \ell \geq n'$ , we have

$$|f_k(x) - f_\ell(x)| \leq \|f_k - f_\ell\|_\infty < \frac{\varepsilon}{2}. \quad (\text{X.4})$$

Now note that  $z \mapsto |f_k(x) - z|$  is continuous  $\mathbb{R} \rightarrow \mathbb{R}$  (in fact 1-Lipschitz) and  $\lim_{\ell \rightarrow \infty} f_\ell(x) = f(x)$ . Therefore we may fix  $k \geq n'$  and take the limit as  $\ell \rightarrow \infty$  in (X.4) and conclude

$$|f_k(x) - f(x)| \leq \frac{\varepsilon}{2} \quad \text{for any } x \in [a, b].$$

As this holds for all  $x \in [a, b]$ , we in fact get

$$\sup \left\{ |f_k(x) - f(x)| \mid x \in [a, b] \right\} \leq \frac{\varepsilon}{2} < \varepsilon \quad (\text{X.5})$$

when  $k \geq n'$ . This shows that the function sequence  $(f_n)_{n \in \mathbb{N}}$  converges uniformly to  $f$ . As a uniform limit of continuous functions, the function  $f$  is also continuous (Theorem IX.8), i.e., we have  $f \in \mathcal{C}([a, b])$ . Moreover, (X.5) can then be read as saying  $\|f_k - f\|_\infty < \varepsilon$  when  $k \geq n'$ , so we conclude that  $\|f_k - f\|_\infty \rightarrow 0$  as  $k \rightarrow \infty$ , and thus  $\lim_{k \rightarrow \infty} f_k = f$  in the metric space  $(\mathcal{C}([a, b]), d_\infty)$ . This proves that an arbitrary Cauchy sequence  $(f_n)_{n \in \mathbb{N}}$  in  $(\mathcal{C}([a, b]), d_\infty)$  converges, which amounts to the asserted completeness.  $\square$

### X.2.4. Completeness for subspaces

In general, subsets of a complete metric space need not be complete. Indeed, the subsets  $\mathbb{R} \setminus \{0\} \subset \mathbb{R}$  and  $\mathbb{Q} \subset \mathbb{R}$  of the complete metric space  $\mathbb{R}$  (see Theorem X.10) were seen to be not complete in Examples X.8 and X.9, respectively. The following result fully answers which subsets of complete metric spaces *are* complete as subspaces.

**Theorem X.13** (Subspaces of a complete metric space).

*Let  $(X, d)$  be a complete metric space. Then a subset  $A \subset X$  is complete as a subspace<sup>3</sup> if and only if  $A \subset X$  is closed.*

**Exercise X.7** (Proof of Theorem X.13).

Prove Theorem X.13.

Hint: Recall Proposition VIII.7.

## X.3. Banach's fixed point theorem

### Iterating a self-map

A **self map** of a given space  $X$  is a function whose domain and codomain are both the same space  $X$ , i.e.,

$$f: X \rightarrow X. \quad (\text{X.6})$$

---

<sup>3</sup>This means that the subset is equipped with the metric inherited from the whole space  $X$ .

Given a self map  $f$  of  $X$  and an initial point  $x_0 \in X$ , we may define a sequence  $(x_n)_{n \in \mathbb{N}_0}$  in  $X$  by

$$\begin{aligned} x_1 &= f(x_0) \\ x_2 &= f(x_1) = f(f(x_0)) \\ x_3 &= f(x_2) = f(f(x_1)) = f(f(f(x_0))) \\ &\vdots \end{aligned}$$

i.e., recursively using

$$x_n = f(x_{n-1}) \in X \quad \text{for } n \in \mathbb{N}. \quad (\text{X.7})$$

The sequence  $(x_n)_{n \in \mathbb{N}_0}$  thus obtained is called the **sequence of successive iterates** of  $x_0$  under  $f$ . The  $n$ :th member of this sequence is the image of  $x_0$  under the function  $X \rightarrow X$  obtained by composing  $f$  with itself  $n$  times; it can be denoted by

$$\underbrace{(f \circ \cdots \circ f)}_{n \text{ times}}(x_0) =: f^{\circ n}(x_0). \quad (\text{X.8})$$

The sequence of successive iterates can then be succinctly denoted  $(f^{\circ n}(x_0))_{n \in \mathbb{N}_0}$ .

**Remark X.14** (Discrete time dynamical systems).

The sequence of successive iterates is basically the mathematical formulation of a “discrete time dynamical system”. In this interpretation,  $X$  is thought of as the “state space” of the system. The self map  $f: X \rightarrow X$  of the “state space” is thought of as “dynamics” in the following way. If the system at some given time instant  $n$  is in “state”  $x_n \in X$ , then at the next time instant  $n + 1$  the system will have evolved to some new state which is a deterministic function of the previous state,  $x_{n+1} = f(x_n)$ . Letting such a deterministic time evolution play out indefinitely in discrete time amounts exactly to considering the sequence of successive iterates of the “initial state”  $x_0 \in X$  of the system at time zero. In this interpretation, the alternative term “orbit” of  $x_0$  is often used for the sequence of successive iterates.

The study of such dynamical systems originates from physics, but is nowadays used in almost any area of science. A traditional example from physics could be the solar system, where the state space parametrizes the positions and velocities of all planets, and the dynamics is determined by Newton’s laws.<sup>4</sup> An interesting application in population biology can be obtained even by simply modeling a population size in a given year as a deterministic function of the population size of the previous year. Chaos theory, in particular, emerged from the study of such dynamical systems.<sup>5</sup>

**Definition X.15** (Fixed point).

Let  $f: X \rightarrow X$  be a function. A point  $x^* \in X$  is called a **fixed point** of  $f$  if  $f(x^*) = x^*$ .

If a fixed point  $x^*$  of  $f$  is used as the initial point  $x_0 = x^*$  of a sequence of successive iterates, then one simply obtains the constant sequence  $(x^*, x^*, x^*, \dots)$ .

<sup>4</sup>Newton’s laws are physics in continuous time, from them one can also of course determine the future state after some fixed amount of time (say 1 year) — or perhaps more interestingly yet, at the next future time when some given planet is in a given plane.

<sup>5</sup>Chaos theory is of course beyond the scope of the present course, but for the reader interested in such scientific questions, it is worth noting that one of the two properties characterizing “chaos” is topological (one is sensitive dependence on initial conditions, and the other is topological transitivity).

## The contraction mapping principle

### Definition X.16 (Contraction).

Let  $(X, d)$  be a metric space. A function  $f: X \rightarrow X$  is called a **contraction** on  $X$  if it is  $K$ -Lipschitz for some  $K < 1$ .

In other words, a contraction is a self-map of a metric space which does not stretch distances and instead contracts them by a nontrivial factor.

The following result is known as the **contraction mapping principle** or alternatively **Banach's fixed point theorem**. The scope of its applications is vast! We will look at a few example applications soon.

### Theorem X.17 (Banach's fixed point theorem).

*Let  $(X, d)$  be a non-empty complete metric space and  $f: X \rightarrow X$  a contraction. Then there exists a unique fixed point  $x^* \in X$  of  $f$ . Moreover, for any  $x_0 \in X$ , the sequence  $(f^{on}(x_0))_{n \in \mathbb{N}}$  of successive iterates of  $x_0$  converges to  $x^*$ .*

*Proof.* Let us fix a  $K < 1$  such that the contraction  $f: X \rightarrow X$  is  $K$ -Lipschitz.

The uniqueness of the fixed point is deduced from the contraction property as follows. If  $x^*, \tilde{x}^* \in X$  are both fixed points, then using first the fixed point equations  $f(x^*) = x^*$  and  $f(\tilde{x}^*) = \tilde{x}^*$ , and then the  $K$ -Lipschitzness of  $f$ , we find

$$d(x^*, \tilde{x}^*) = d(f(x^*), f(\tilde{x}^*)) \leq K d(x^*, \tilde{x}^*).$$

Moving the term on the right to the left hand side this implies that  $(1 - K)d(x^*, \tilde{x}^*) \leq 0$ , which is only possible if  $d(x^*, \tilde{x}^*) = 0$ . By axiom (??), this means  $x^* = \tilde{x}^*$ , so two different points could not have been fixed points.

Let us then use the contraction property to show that the sequence  $(x_n)_{n \in \mathbb{N}_0} = (f^{on}(x_0))_{n \in \mathbb{N}_0}$  of successive iterates of any point  $x_0 \in X$  is Cauchy. Denote  $r = d(x_0, x_1) = d(x_0, f(x_0))$ . Now note that by the  $K$ -Lipschitz property, for any  $n \in \mathbb{N}$  we have

$$d(x_n, x_{n+1}) = d(f(x_{n-1}), f(x_n)) \leq K d(x_{n-1}, x_n).$$

By an easy induction starting from  $d(x_0, x_1) = r$  this implies that  $d(x_n, x_{n+1}) \leq rK^n$  for any  $n \in \mathbb{N}_0$ . Let  $k, \ell \in \mathbb{N}$  with  $k < \ell$ . To estimate the distance between members  $x_k$  and  $x_\ell$  of the sequence, we apply the triangle inequality (iterated finitely many times) and get

$$\begin{aligned} d(x_k, x_\ell) &\leq d(x_k, x_{k+1}) + d(x_{k+1}, x_{k+2}) + \cdots + d(x_{\ell-1}, x_\ell) \\ &\leq rK^k + rK^{k+1} + \cdots + rK^{\ell-1} \\ &= r \sum_{j=k}^{\ell-1} K^j = rK^k \frac{1 - K^\ell}{1 - K} \leq \frac{r}{1 - K} K^k. \end{aligned}$$

Now let  $\varepsilon > 0$ . Since  $\frac{r}{1-K} K^n \rightarrow 0$  as  $n \rightarrow \infty$ , there exists some  $n_\varepsilon$  so that  $\frac{r}{1-K} K^n < \varepsilon$  for any  $n \geq n_\varepsilon$ . The above estimate then shows that for any  $k, \ell \geq n_\varepsilon$  we have  $d(x_k, x_\ell) < \varepsilon$ . This shows that the sequence  $(x_n)_{n \in \mathbb{N}_0}$  is Cauchy.

Let us then note that by completeness of  $X$ , the sequence  $(x_n)_{n \in \mathbb{N}_0} = (f^{on}(x_0))_{n \in \mathbb{N}_0}$  of successive iterates of any point  $x_0 \in X$  converges, since we saw above that it is Cauchy. Let  $x' = \lim_{n \rightarrow \infty} x_n$  be the limit of such a sequence. We claim that  $x'$  is a fixed point of  $f$ . To see this, consider both the sequence  $(x_n)_{n \in \mathbb{N}_0}$  which converges to  $x'$  and the sequence  $(f(x_n))_{n \in \mathbb{N}_0}$  of its images under  $f$ . The contraction  $f$  is continuous (in fact Lipschitz continuous), so by Corollary VIII.11 the limit of the sequence  $(f(x_n))_{n \in \mathbb{N}_0}$  is

$$\lim_{n \rightarrow \infty} f(x_n) = f(x').$$

On the other hand,  $f(x_n) = x_{n+1}$ , so the sequence of images is simply a shifted version of the original sequence

$$(f(x_0), f(x_1), f(x_2), \dots) = (x_1, x_2, x_3, \dots),$$

and it in particular converges to the same limit  $x'$ . Since we thus have

$$\lim_{n \rightarrow \infty} f(x_n) = x',$$

a comparison with the previously obtained limit gives

$$x' = f(x'),$$

i.e., the limit  $x'$  is a fixed point of  $f$ .

Since we have already shown that there can only exist one fixed point, we conclude from the above argument that the sequences of iterates with all initial values  $x_0 \in X$  converge to the same fixed point  $x^* \in X$  of  $f$ . We finish by noting that since we assumed that  $X \neq \emptyset$ , it is possible to choose at least some initial value  $x_0 \in X$  and form its sequence of iterates, and thus obtain a limit which is a fixed point. This shows the existence of a fixed point  $x^* \in X$ . The proof is complete.  $\square$

## Applications

Banach's fixed point theorem, Theorem X.17, can be used to show the existence and uniqueness of solutions and the form an iterative approximation scheme for the solution. Let us look at a few examples.

**Example X.18** (A simple application of Banach's fixed point theorem).

Consider the equation

$$\cos(x) = x \quad \text{on the interval } x \in [0, 1]. \quad (\text{X.9})$$

By plotting the left and right sides of the equation, it is easy to convince oneself that there is a unique solution to this equation (what is the easiest way to prove this rigorously?). However, familiar methods do not give a closed form solution to (X.9). Here we will not only show the existence and uniqueness of a solution, but also find a method of producing increasingly good approximations of it, with rigorous bounds on the error of the approximations.

Since  $\cos(x) \in [0, 1]$  for  $x \in [0, 1]$ , the formula

$$f(x) = \cos(x) \quad \text{defines a function} \quad f: [0, 1] \rightarrow [0, 1].$$

Solutions to (X.9) are exactly the fixed points of  $f$ .

Now let us first note that by Theorems X.13 and X.10, the closed interval  $[0, 1]$  is a complete metric space, since it is a closed subset of the complete metric space  $\mathbb{R}$ .

To show that the function  $f$  above is a contraction, we find a bound for the absolute value of its derivative that is below one. Indeed, the derivative is  $f'(x) = -\sin(x)$ , and it satisfies the bound  $|f'(x)| \leq \sin(1)$  for all  $x \in [0, 1]$  (the derivative is negative and decreasing, so attains its minimum at  $x = 1$ ). Therefore denoting  $K = \sin(1)$  we obtain as in Example VI.7 that  $f$  is  $K$ -Lipschitz, and since  $K = \sin(1) < 1$ , the function  $f$  is a contraction.

Since  $[0, 1]$  is complete and non-empty, and  $f$  is a contractive self-map of it, we may apply Banach's fixed point theorem. The first part of Banach's fixed point theorem guarantees that (X.9) indeed has a unique solution  $x = x^* \in [0, 1]$  (the unique fixed point  $x^*$  of  $f$ ). Moreover, by the second part, given any  $x_0 \in [0, 1]$ , we know that the sequence  $(x_n)_{n \in \mathbb{N}_0} = (f^{cn}(x_0))_{n \in \mathbb{N}_0}$  converges to the solution  $x^*$ . This allows us to form increasingly



good approximate solutions. For example by letting  $x_0 = 0$ , the successive iterates are<sup>6</sup>

$$\begin{aligned}x_0 &= 0 \\x_1 &= \cos(x_0) = 1 \\x_2 &= \cos(x_1) = \cos(1) \approx 0.540302 \\x_3 &= \cos(x_2) = \cos(\cos(1)) \approx 0.857553 \\x_4 &= \cos(x_3) = \cos(\cos(\cos(1))) \approx 0.65429 \\x_5 &= \cos(x_4) = \cos(\cos(\cos(\cos(1)))) \approx 0.79348 \\x_6 &= \cos(x_5) = \cos(\cos(\cos(\cos(\cos(1)))))) \approx 0.701369 \\&\vdots\end{aligned}$$

A good numerical value of the true solution  $x^*$  is

$$x^* \approx 0.73908513321$$

and for example the 25th and 50th members of the iterative approximation scheme started from  $x_0 = 0$  yield quite reasonable approximations<sup>7</sup>

$$x_{25} \approx 0.73910557193$$

$$x_{50} \approx 0.73908513217.$$

We can also control the error made when using the iterates  $x_n$  as approximations of the true solution  $x^*$ : from the proof of Theorem X.17 we see that  $|x_n - x^*| \leq \frac{|x_1 - x_0|}{1-K} K^n$ . Plugging in values  $x_1 = 1$ ,  $x_0 = 0$ , and  $K = \sin(1) \approx 0.841471$  (in the form  $0.841 < K < 0.842$ ) we find that

$$|x_n - x^*| \leq \frac{|x_1 - x_0|}{1-K} K^n < \frac{1}{1-0.842} (0.841)^n = \frac{(0.841)^n}{0.158}.$$

The error estimates obtained here are a bit conservative: they yield for example

$$|x_{25} - x^*| < 0.0835 \quad \text{and} \quad |x_{50} - x^*| < 0.0011;$$

correctly indeed, but the reality is even better... The point is that such upper bounds for errors are fully rigorous!

The previous example was about solving an equation on the real line  $\mathbb{R}$ . Perhaps you already knew other methods that could have been used to numerically solve the equation (X.9).<sup>8</sup> But Banach's fixed point theorem applies in much more general spaces! Solving the following differential equation by other methods would be more difficult, wouldn't it?

**Example X.19** (Solving a differential equation).

Consider the differential equation

$$y'(s) = \sin(s y(s)) \quad \text{for } s \in [0, 1], \quad (\text{X.10})$$

with the initial condition

$$y(0) = 1. \quad (\text{X.11})$$

Finding an explicit formula for a solution to (X.10)–(X.11) seems hopeless (even with the help of computer algebra systems). We will nevertheless again show the existence and uniqueness of a solution, and find a method of producing increasingly good approximations of the true solution.

<sup>6</sup>With an old fashioned calculator, you can quickly obtain the iterates by entering the initial value 0, and then repeatedly pressing the cos-button.

<sup>7</sup>See for yourself, by repeatedly hitting the cos-button of an old-fashioned calculator, how the approximations converge to the solution.

<sup>8</sup>For example, write (X.9) in the form  $\cos(x) - x = 0$ , and use Newton's iteration to find a zero of the left hand side.

Observe first that the differential equation (X.10) is equivalent with its integrated form

$$\underbrace{\int_0^t y'(s) \, ds}_{y(t)-y(0)} = \int_0^t \sin(s y(s)) \, ds \quad \text{for all } t \in [0, 1],$$

and the initial value problem (X.10)–(X.11) is therefore equivalent to the integral equation

$$y(t) = 1 + \int_0^t \sin(s y(s)) \, ds \quad \text{for all } t \in [0, 1]. \quad (\text{X.12})$$

Let us now define a mapping

$$f: \mathcal{C}([0, 1]) \rightarrow \mathcal{C}([0, 1])$$

by the formula<sup>9</sup>

$$[f(x)](t) = 1 + \int_0^t \sin(s x(s)) \, ds \quad \text{for } t \in [0, 1].$$

The integral equation (X.12) is nothing but the fixed point equation  $y = f(y)$  for  $f$ . Therefore this fixed point equation is in fact equivalent to the initial value problem (X.10)–(X.11).

We equip the space  $\mathcal{C}([0, 1])$  with the metric induced by the norm  $\|\cdot\|_\infty$ , and recall that  $\mathcal{C}([0, 1])$  is then a complete metric space by Theorem X.12. We will show that the function  $f$  above is a contraction of  $\mathcal{C}([0, 1])$ . So, suppose that  $x_1, x_2 \in \mathcal{C}([0, 1])$ . In order to estimate the distance between  $f(x_1)$  and  $f(x_2)$ , first bound the absolute value of the function  $f(x_2) - f(x_1) \in \mathcal{C}([0, 1])$  at a point  $t \in [0, 1]$ :

$$\begin{aligned} |[f(x_2) - f(x_1)](t)| &= |[f(x_2)](t) - [f(x_1)](t)| \\ &= \left| 1 + \int_0^t \sin(s x_2(s)) \, ds - 1 - \int_0^t \sin(s x_1(s)) \, ds \right| \\ &= \left| \int_0^t (\sin(s x_2(s)) - \sin(s x_1(s))) \, ds \right| \\ &\leq \int_0^t |\sin(s x_2(s)) - \sin(s x_1(s))| \, ds. \end{aligned}$$

Using  $|\sin(\theta_2) - \sin(\theta_1)| \leq |\theta_2 - \theta_1|$  (the function  $\sin$  has derivative  $|\sin'(\theta)| = |\cos(\theta)| \leq 1$  bounded by 1, and is therefore 1-Lipchitz by Example VI.7), we see that the integrand in the last expression is bounded by

$$\begin{aligned} |\sin(s x_2(s)) - \sin(s x_1(s))| &\leq |s x_2(s) - s x_1(s)| = |s| |x_2(s) - x_1(s)| \\ &\leq s \|x_2 - x_1\|_\infty. \end{aligned}$$

Plugging this estimate into the previous expression, we find

$$|[f(x_2) - f(x_1)](t)| \leq \|x_2 - x_1\|_\infty \int_0^t s \, ds = \|x_2 - x_1\|_\infty \frac{t^2}{2} \leq \frac{1}{2} \|x_2 - x_1\|_\infty.$$

Since this holds for any  $t \in [0, 1]$ , by taking the supremum over  $t$  we get

$$\|f(x_2) - f(x_1)\|_\infty \leq \frac{1}{2} \|x_2 - x_1\|_\infty,$$

which shows that  $f: \mathcal{C}([0, 1]) \rightarrow \mathcal{C}([0, 1])$  is  $\frac{1}{2}$ -Lipchitz

Since  $f$  thus is a contraction of the non-empty complete metric space  $\mathcal{C}([0, 1])$ , Banach's fixed point theorem applies. It guarantees first of all that there exists a unique solution  $y \in \mathcal{C}([0, 1])$  to the initial value problem (X.10)–(X.11) (namely the unique fixed point  $y$  of  $f$ ). This already is an important conclusion.

Moreover, Banach's fixed point theorem gives the following method of approximating the solution  $y$ : choose any  $x_0 \in \mathcal{C}([0, 1])$ , and form the sequence  $(x_n)_{n \in \mathbb{N}_0} = (f^{on}(x_0))_{n \in \mathbb{N}_0}$  of successive iterates. Then we have  $\lim_{n \rightarrow \infty} f^{on}(x_0) = y$  in  $\mathcal{C}([0, 1])$  with respect to the

<sup>9</sup>Note that the argument of the function  $f: \mathcal{C}([0, 1]) \rightarrow \mathcal{C}([0, 1])$  is itself a function  $x \in \mathcal{C}([0, 1])$ , and the value is also,  $f(x) \in \mathcal{C}([0, 1])$ .

norm  $\|\cdot\|_\infty$ , i.e., uniformly. For example, we may take  $x_0: [0, 1] \rightarrow \mathbb{R}$  to be the constant function  $x_0(s) = 1$  for all  $s \in [0, 1]$ . The first few iterates are plotted in Figure ??.

The idea in Example X.19 works very generally with ordinary differential equations: the Pickard–Lindelöf theorem guarantees the existence and uniqueness of solutions, and gives an iterative approximation scheme for them. Its proof is a very straightforward modification of the above particular case.

## X.4. ♥ Hilbert spaces and Banach spaces

Very powerful theory can be developed about normed spaces and inner product spaces which are complete as metric spaces with the induced metric. Due to the prominence of the theory and scope of the applications, such spaces have been named as follows.

**Definition X.20** (Hilbert space).

An inner product space  $(V, \langle \cdot, \cdot \rangle)$ , which is complete as a metric space with respect to the metric induced by the inner product, is called a **Hilbert space**.

**Definition X.21** (Banach space).

A normed space  $(V, \|\cdot\|)$ , which is complete as a metric space with respect to the metric induced by the norm, is called a **Banach space**.

**Remark X.22** (Hilbert spaces are Banach spaces).

Every Hilbert space has a norm induced by its inner product. The metric induced by the inner product is (by definition) the same as the metric induced by the norm, so it is complete in the latter as well. Therefore every Hilbert space is in particular a Banach space.

The converse is not true: there are Banach spaces which are not Hilbert spaces; see Example X.24 below.

Our first example is familiar.

**Example X.23** (Euclidean spaces are (finite-dimensional) Hilbert spaces).

By Theorem X.11, the  $d$ -dimensional vector space  $\mathbb{R}^d$  with its usual inner product is complete, and therefore a Hilbert space. By Remark X.22 it is also a Banach space.

The theory of Hilbert spaces and Banach spaces becomes more important in the infinite-dimensional setting, however. We have seen one important example of an infinite-dimensional Banach space already.

**Example X.24** (A Banach space of continuous functions).

By Theorem X.12, the space  $\mathcal{C}([a, b])$  of continuous functions on a closed interval  $[a, b]$ , equipped with the supremum norm  $\|\cdot\|_\infty$ , is complete. Therefore it is a Banach space.

The following is an example of an infinite-dimensional Hilbert space.

**Example X.25** (An infinite-dimensional Hilbert space).

The space

$$\ell^2 = \left\{ (a_n)_{n \in \mathbb{N}} \mid \sum_{n=1}^{\infty} a_n^2 < \infty \right\}$$

of square summable sequences has an inner product by Example IV.19.

This space is complete by Exercise X.8 below. Therefore  $\ell^2$  is a Hilbert space.

**Exercise (§) X.8** (Proof of completeness of  $\ell^2$ ).

Prove that  $\ell^2$  is complete.

**Remark X.26** (Complex Hilbert spaces and complex Banach spaces).

In this course we have only considered real vector spaces. For complex vector spaces, there are notions of norm and inner product as well. Just like their real counterparts, they also induce metrics. By complex Hilbert spaces and complex Banach spaces, one means complex inner product spaces and complex normed spaces which are complete as metric spaces with the induced metric. In fact the theory of complex Hilbert spaces and Banach spaces is perhaps slightly simpler (not more complex, eh?) than that of real Hilbert spaces and Banach spaces. Complex Hilbert spaces and Banach spaces are also very common in applications.

Complex Hilbert spaces in particular have an indispensable role in the foundations of quantum mechanics and quantum field theory.

Both Hilbert spaces and Banach spaces are used very widely in mathematics and its applications. You will learn more about them in your subsequent studies!

## Appendix A

### Further topics in set theory

#### A.1. ♥ Cardinalities of sets

##### An observation about finite sets

Let us denote the number of elements of a set  $A$  by  $\#A$ .

**Example A.1** (Number of elements in a few example sets).

- For the empty set, we have  $\#\emptyset = 0$ .
- For a singleton, we have  $\#\{a\} = 1$ .
- We have  $\#\{0, 1, 2, \dots, 8, 9\} = 10$ .
- Let  $c_0, c_1, \dots, c_{d-1}, c_d \in \mathbb{R}$  with  $c_d \neq 0$ , so that  $p(x) = c_d x^d + c_{d-1} x^{d-1} + \dots + c_1 x + c_0$  is a polynomial of degree  $d \in \mathbb{N}_0$ . Then for the set of real zeroes of  $p$  we have

$$\#\{x \in \mathbb{R} \mid p(x) = 0\} \leq d.$$

If a set  $A$  has infinitely many different elements, we denote  $\#A = \infty$ .

**Example A.2** (Some infinite sets).

The sets of natural numbers, integers, rational numbers, and real numbers are infinite

$$\#\mathbb{N} = \infty, \quad \#\mathbb{Z} = \infty, \quad \#\mathbb{Q} = \infty, \quad \#\mathbb{R} = \infty.$$

Also nontrivial intervals are infinite, for example  $\#(\pi, \pi] = \infty$ .

Moreover, as we will verify later, a non-empty open interval  $(a, b) \subset \mathbb{R}$  contains infinitely many rational numbers and infinitely many irrational numbers

$$\#\left((a, b) \cap \mathbb{Q}\right) = \infty, \quad \#\left((a, b) \setminus \mathbb{Q}\right) = \infty.$$

A set  $A$  is called **finite** if  $\#A \in \{0, 1, 2, \dots\}$ , and **infinite** if  $\#A = \infty$ . It turns out that among infinite sets, some are nevertheless larger than others. The notion of *cardinality* captures this. To motivate the definition, let us nevertheless begin by some observations about sizes of finite sets.

**Observation A.3** (Comparing sizes of sets using surjective functions).

If  $A, B$  are two finite sets and  $B \neq \emptyset$ , then the following are equivalent:<sup>1</sup>

- $\#A \geq \#B$
- there exists a surjective function  $A \rightarrow B$ .<sup>2</sup>

<sup>1</sup>For the empty set  $B = \emptyset$ , the comparison of sizes has to be handled separately (the empty set is smaller than any other set), since from a nonempty set  $A \neq \emptyset$  there does not exist any functions  $A \rightarrow \emptyset$  — let alone surjective functions.

<sup>2</sup>The idea is the following. Suppose that  $A$  is the set of all students of this course, and  $B$  is the set of all exercise groups. Every student is assigned to exactly one exercise group, so that the assignment defines a function  $A \rightarrow B$ . The function is surjective if every exercise group has at least one student. The gist of this observation is that in such a case we can conclude that there are

## Cardinality comparison and equal cardinalities

In view of the above observation, it appears meaningful to consider a set  $A$  at least as large as a set  $B \neq \emptyset$  if there exists a surjective function  $A \rightarrow B$ . In this case we denote  $A \succeq B$  — or interchangeably  $B \preceq A$ . This is the comparison of **cardinalities** of sets. We may observe the following properties, which match with our intuition about sizes of sets:

- If  $A \succeq B$  and  $B \succeq C$ , then  $A \succeq C$ .<sup>3</sup>
- If  $B \subset A$  is a nonempty subset, then  $A \succeq B$ .<sup>4</sup>

**Example A.4** (Cardinality comparisons of some infinite sets).

Because of the subset relations  $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$ , we have the cardinality comparisons  $\mathbb{N} \preceq \mathbb{Z} \preceq \mathbb{Q} \preceq \mathbb{R}$ .

We say that two nonempty sets  $A$  and  $B$  have **equal cardinalities** if  $A \succeq B$  and  $A \preceq B$ . By Observation A.3, (nonempty) finite sets have equal cardinalities if and only if they have the same number of elements. Let us then look at some examples with infinite sets.

**Example A.5** (Equal cardinality for a set and its proper subsets).

Consider the set  $\mathbb{N}$  of natural numbers, and the set  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$  of nonnegative integers,

$$\begin{aligned}\mathbb{N} &= \{1, 2, 3, 4, \dots\} \\ \mathbb{N}_0 &= \{0, 1, 2, 3, 4, \dots\}.\end{aligned}$$

Because of the subset relation  $\mathbb{N} \subset \mathbb{N}_0$ , we have  $\mathbb{N} \preceq \mathbb{N}_0$ . On the other hand, we also have a surjective function  $f: \mathbb{N} \rightarrow \mathbb{N}_0$  given by  $f(n) = n - 1$  for  $n \in \mathbb{N}$ , so that  $\mathbb{N} \succeq \mathbb{N}_0$  holds, too. Therefore the sets  $\mathbb{N}$  and  $\mathbb{N}_0$  have equal cardinalities.

In particular an infinite set can have a proper subset with equal cardinality: the set  $\mathbb{N}_0$  has one extra element compared to  $\mathbb{N}$  (namely zero), but this does not affect the “size” of the set; these two infinite sets have equal cardinalities.

**Example A.6** (Another equal cardinality for a set and its proper subsets).

Consider the set  $\mathbb{N}$  of natural numbers and the set  $A = \{2, 4, 6, \dots\}$  of even natural numbers. Because of the subset relation  $A \subset \mathbb{N}$ , we have  $A \preceq \mathbb{N}$ . On the other hand, we also have a surjective function  $f: A \rightarrow \mathbb{N}$  given by  $f(m) = \frac{1}{2}m$  for  $m \in A$ , so that  $A \succeq \mathbb{N}$  holds, too. Therefore the sets  $\mathbb{N}$  and  $A$  have equal cardinalities.

The set  $\mathbb{N}_0$  has infinitely many extra elements compared to  $A$  (namely the odd natural numbers), but this does not affect the “size” of the set; these two infinite sets have equal cardinalities.

---

at least as many students as there are exercise classes (and conversely: an assignment that leaves no exercise group empty exists if there are at least as many students as there are exercise classes).

<sup>3</sup>Recall Exercise I.9: if  $f: A \rightarrow B$  and  $g: B \rightarrow C$  are surjective, then also the composition  $g \circ f: A \rightarrow C$  is surjective.

<sup>4</sup>Indeed, choose some element  $b_0 \in B$  of the nonempty set  $B$ , and define a function  $f$  from  $A$  to the subset  $B \subset A$  by

$$f(a) = \begin{cases} a & \text{if } a \in B \subset A \\ b_0 & \text{if } a \in A \setminus B. \end{cases}$$

Then  $f$  is surjective.

**Exercise A.1** (The sets of natural numbers and integers have equal cardinalities).

Prove that we have  $\mathbb{N} \preceq \mathbb{Z}$  and  $\mathbb{N} \succeq \mathbb{Z}$ .

Hint: One direction is obvious; for the other, construct some surjective function  $\mathbb{N} \rightarrow \mathbb{Z}$ .

## Countable and uncountable infinite sets

These examples show that many common infinite sets have the same cardinality as the set of natural numbers. This motivates the following definition.

**Definition A.7** (Countable and uncountable infinite sets).

An infinite set  $A$  is **countably infinite** if  $A \preceq \mathbb{N}$ , i.e., if there exists a surjective function  $f: \mathbb{N} \rightarrow A$ ; otherwise  $A$  is **uncountably infinite**.

**Remark A.8** (Enumeration of elements of a countably infinite set).

If  $A$  is a countably infinite set, then it is possible to list or enumerate all elements of  $A$  as follows. If  $f: \mathbb{N} \rightarrow A$  is a surjective function, then we may form a sequence

$$(a_1, a_2, a_3, \dots)$$

with  $a_n = f(n)$  for  $n \in \mathbb{N}$ . By surjectivity, each element of  $A$  appears at least once in this sequence.

It is not possible to enumerate elements of uncountably infinite sets in this way!

In Examples A.5 and A.6 and Exercise A.1 we have seen some examples of countably infinite sets. The most important example of an uncountably infinite set is the set of real numbers.

**Theorem A.9** (The set of real numbers is uncountably infinite).

*The set  $\mathbb{R}$  of real numbers is uncountably infinite.*

The proof is given in the next subsection.

By contrast, and perhaps surprisingly, the set of rational numbers is in fact countable. In this sense the set  $\mathbb{Q}$  of rational numbers is much smaller than the set  $\mathbb{R}$  of real numbers.

**Theorem A.10** (The set of rational numbers is countably infinite).

*The set  $\mathbb{Q}$  of rational numbers is countably infinite.*

### Countability of the set of rational numbers

The key to the proof of Theorem A.10 is the following frequently useful lemma.

**Lemma A.11** (The Cartesian product of two countable sets is countable).

*Suppose that  $A$  and  $B$  are two countably infinite sets. Then their Cartesian product  $A \times B$  is also countably infinite.*

It is still clearest to prove this lemma by first explicitly addressing the following special case.

**Lemma A.12** (The set of pairs of natural numbers is countable).

*The set  $\mathbb{N} \times \mathbb{N} = \{(n, m) \mid n \in \mathbb{N}, m \in \mathbb{N}\}$  of pairs of natural numbers is countably infinite.*

*Proof.* We must show that there exists a surjective function  $t: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ . This can be done in various ways; the enumeration illustrated in Figure A.1 is easy to visualize, but let us choose a different construction. Namely, the function  $t: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$  given by

$$t(n) = \begin{cases} (k, \ell) & \text{if } n = 2^{k-1} 3^{\ell-1} \text{ for } k, \ell \in \mathbb{N} \\ (1, 1) & \text{if } n \text{ contains other prime factors besides 2 and 3} \end{cases}$$

is well-defined because of the unique prime factorization of natural numbers, and it is surjective, since any  $(k, \ell) \in \mathbb{N} \times \mathbb{N}$  is obtained as a value at  $n = 2^{k-1} 3^{\ell-1} \in \mathbb{N}$ .  $\square$

*Proof of Lemma A.11.* Assume that  $A$  and  $B$  are countable. To prove countability of  $A \times B$ , we must exhibit a surjective function  $\mathbb{N} \rightarrow A \times B$ .

By countability of  $A$  and  $B$ , there exists surjective functions

$$g_1: \mathbb{N} \rightarrow A \qquad g_2: \mathbb{N} \rightarrow B.$$

We may combine them into a function  $g: \mathbb{N} \times \mathbb{N} \rightarrow A \times B$  defined by

$$g(n, m) = (g_1(n), g_2(m)),$$

which is clearly surjective.<sup>5</sup> From Lemma A.12 we get the existence of a surjective function  $t: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ . Now the composition  $g \circ t: \mathbb{N} \rightarrow A \times B$ ,

$$\mathbb{N} \xrightarrow{t} \mathbb{N} \times \mathbb{N} \xrightarrow{g} A \times B, \\ \qquad \qquad \qquad \searrow \scriptstyle{g \circ t}$$

is surjective. Countability of  $A \times B$  follows.  $\square$

*Proof of Theorem A.10.* To show countability of the set  $\mathbb{Q}$  of rational numbers, we must exhibit a surjective function  $\mathbb{N} \rightarrow \mathbb{Q}$ .

By definition, rational numbers  $q \in \mathbb{Q}$  are of the form  $q = \frac{m}{n}$ , with  $m \in \mathbb{Z}$  and  $n \in \mathbb{N}$ . In other words, the function

$$r: \mathbb{Z} \times \mathbb{N} \rightarrow \mathbb{Q} \qquad \text{given by} \qquad r(m, n) = \frac{m}{n} \qquad \text{for } (m, n) \in \mathbb{Z} \times \mathbb{N}$$

is surjective.

The set  $\mathbb{Z}$  of integers is countable by Exercise A.1, and the set  $\mathbb{N}$  of natural numbers is obviously countable. By Lemma A.11 the Cartesian product  $\mathbb{Z} \times \mathbb{N}$  is therefore countable, so there exists a surjective function

$$s: \mathbb{N} \rightarrow \mathbb{Z} \times \mathbb{N}.$$

Now the composition  $r \circ s: \mathbb{N} \rightarrow \mathbb{Q}$ ,

$$\mathbb{N} \xrightarrow{s} \mathbb{Z} \times \mathbb{N} \xrightarrow{r} \mathbb{Q}, \\ \qquad \qquad \qquad \searrow \scriptstyle{r \circ s}$$

is surjective. Countability of  $\mathbb{Q}$  follows.  $\square$

<sup>5</sup>Indeed, suppose that  $(a, b) \in A \times B$ . By surjectivity of  $g_1: \mathbb{N} \rightarrow A$  there exists an  $n \in \mathbb{N}$  such that  $g_1(n) = a$  and by surjectivity of  $g_2: \mathbb{N} \rightarrow B$  there exists an  $m \in \mathbb{N}$  such that  $g_2(m) = b$ . Therefore we have  $g(n, m) = (g_1(n), g_2(m)) = (a, b)$ , showing surjectivity of  $g$ .





FIGURE A.1. The arrows indicate an enumeration of  $\mathbb{N} \times \mathbb{N}$ :

$((1, 1), (2, 1), (1, 2), (3, 1), (2, 2), (1, 3), (4, 1), (3, 2), (2, 3), (1, 4), \dots)$ .

The idea is to observe that for pairs  $(n, m) \in \mathbb{N} \times \mathbb{N}$ , the sum  $n + m$  takes values  $2, 3, 4, \dots$ , and for each value of the sum it is straightforward to list the finitely many pairs  $(n, m)$  in increasing order of the corresponding  $m$  (for example). This enumeration gives another construction of a surjective function  $\mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ .

### ♥ Uncountability of the set of real numbers

Let us now provide the proof that the set  $\mathbb{R}$  of real numbers is uncountably infinite. The argument is known as Cantor's diagonal argument. Here we use it to show the impossibility of enumerating the real numbers (even on the unit interval), but variants of the diagonal argument are also used frequently constructively; we will even see some examples later in this course.

*Proof of Theorem A.9.* In order to prove that  $\mathbb{R}$  is not countable, it suffices to prove that the subset  $[0, 1) \subset \mathbb{R}$  is not countable.

We do the proof by contradiction: we assume that  $[0, 1)$  is countable, and derive a contradiction from this assumption. If  $[0, 1)$  were countable, then there would exist a surjective function  $f: \mathbb{N} \rightarrow [0, 1)$ . For each  $n \in \mathbb{N}$ , we could write the decimal expansion of the number  $f(n) \in [0, 1)$ :

$$f(n) = 0.d_1^{(n)}d_2^{(n)}d_3^{(n)}\dots \quad \text{or more precisely} \quad f(n) = \sum_{j=1}^{\infty} d_j^{(n)} 10^{-j},$$

where  $d_j^{(n)} \in \{0, 1, 2, \dots, 8, 9\}$  for every  $j \in \mathbb{N}$ . In order to show that  $f$  could not have been surjective, we now examine these decimal expansions

$$\begin{aligned} f(1) &= 0.\underline{d}_1^{(1)} d_2^{(1)} \underline{d}_3^{(1)} d_4^{(1)} \dots \\ f(2) &= 0.d_1^{(2)} \underline{d}_2^{(2)} d_3^{(2)} \underline{d}_4^{(2)} \dots \\ f(3) &= 0.d_1^{(3)} d_2^{(3)} \underline{d}_3^{(3)} d_4^{(3)} \dots \\ f(4) &= 0.d_1^{(4)} d_2^{(4)} d_3^{(4)} \underline{d}_4^{(4)} \dots \\ &\vdots = \quad \ddots \end{aligned}$$

in particular along the diagonal marked by the underlined digits  $\underline{d}_k^{(k)}$ ,  $k \in \mathbb{N}$ . Let us construct a number  $x \in [0, 1)$  whose first decimal digit is different from the first digit of  $f(1)$ , second digit is different from the second digit of  $f(2)$ , etc. Specifically, we choose the  $k$ :th digit as

$$d_k := \begin{cases} 8 & \text{if } d_k^{(k)} = 3, \\ 3 & \text{if } d_k^{(k)} \neq 3, \end{cases}$$

which guarantees that  $d_k \neq d_k^{(k)}$  for each  $k \in \mathbb{N}$ . Now consider the number

$$x = 0.d_1 d_2 d_3 \dots \quad \text{or more precisely} \quad x = \sum_{j=1}^{\infty} d_j 10^{-j}.$$

We have  $0 \leq x < 1$ , (the strict second inequality uses the fact that 9 does not appear as a digit in  $x$ , so  $x$  is not  $0.9999\dots = 1$ ). It is straightforward to check that for every  $k \in \mathbb{N}$ , we have  $x \neq f(k)$ , since the  $k$ :th digits of  $x$  and  $f(k)$  are different. But the existence of such an  $x \in [0, 1)$  contradicts the surjectivity of  $f$ , and finishes the proof.  $\square$

## Appendix B

### Further topics about real numbers

#### B.1. ♥ Field axioms and order axioms of the real numbers

Here we take a somewhat closer look into the *field axioms* and *order axioms* of the real numbers. The properties addressed in these axioms are undoubtedly more familiar than those in the completeness axiom. A closer look at nevertheless offers an instructive perspective especially into the similarities and differences of the real numbers to other fields used in mathematics (rational numbers, complex numbers, algebraic number fields, finite fields, ...).

##### ♥ Field axioms

The field axioms concern two operations of calculation with numbers: addition and multiplication, and we of course use the standard notational convention that multiplication takes precedence over addition.

The field axioms of real numbers state that the set  $\mathbb{R}$  equipped with the operations  $+$  and  $\cdot$  is a field:

Commutativity of addition:

$$\forall x, y \in \mathbb{R} : x + y = y + x \quad (\text{B.1})$$

Associativity of addition:

$$\forall x, y, z \in \mathbb{R} : x + (y + z) = (x + y) + z \quad (\text{B.2})$$

Neutral element for addition:

$$\exists 0 \in \mathbb{R} : \forall x \in \mathbb{R} : x + 0 = x \quad (\text{B.3})$$

Opposite elements:

$$\forall x \in \mathbb{R} : \exists (-x) \in \mathbb{R} : x + (-x) = 0 \quad (\text{B.4})$$

Commutativity of multiplication:

$$\forall x, y \in \mathbb{R} : x \cdot y = y \cdot x \quad (\text{B.5})$$

Associativity of multiplication:

$$\forall x, y, z \in \mathbb{R} : x \cdot (y \cdot z) = (x \cdot y) \cdot z \quad (\text{B.6})$$

Neutral element of multiplication:

$$\exists 1 \in \mathbb{R} \setminus \{0\} : \forall x \in \mathbb{R} : x \cdot 1 = x \quad (\text{B.7})$$

Inverse elements:

$$\forall x \in \mathbb{R} \setminus \{0\} : \exists x^{-1} \in \mathbb{R} : x \cdot x^{-1} = 1 \quad (\text{B.8})$$

Distributivity of addition over multiplication:

$$\forall x, y, z \in \mathbb{R} : x \cdot (y + z) = x \cdot y + x \cdot z \quad (\text{B.9})$$

Hopefully the reader finds these uncontroversial enough to be admitted as axioms.

Note that the axioms explicitly state the existence of two real numbers, 0 (“zero”, the neutral element for addition) and 1 (“one”, the neutral element for multiplication), and they require that these two are different,  $1 \neq 0$  (since  $1 \in \mathbb{R} \setminus \{0\}$ ). Moreover, zero and one are uniquely determined by the properties required of them in the axioms.

**Example B.1** (Uniqueness of zero).

*Claim:* If both  $0 \in \mathbb{R}$  and  $0' \in \mathbb{R}$  satisfy the property of being neutral elements for addition, then we have  $0 = 0'$ .

*Proof:* Suppose that  $x + 0 = x$  and  $x + 0' = x$  for all  $x \in \mathbb{R}$ . Then we get

$$0 \stackrel{(\text{B.3})'}{=} 0 + 0' \stackrel{(\text{B.1})}{=} 0' + 0 \stackrel{(\text{B.3})}{=} 0' ,$$

where we first used the neutral element property of  $0'$ , then commutativity of addition, and finally the neutral element property of  $0$ .  $\square$

**Exercise (✓) B.1** (Uniqueness of one).

Prove from the field axioms of real numbers that if both  $1 \in \mathbb{R}$  and  $1' \in \mathbb{R}$  satisfy the property of being neutral elements for multiplication, then we have  $1 = 1'$ .

**Exercise B.2** (Uniqueness of opposite elements and inverse elements).

Prove that for any  $x \in \mathbb{R}$ , the opposite element  $(-x)$  is unique, and that for any  $x \in \mathbb{R} \setminus \{0\}$ , the inverse element  $x^{-1}$  is unique.

The following example indicates how yet a few other familiar facts about the real numbers are proved starting from the field axioms.

**Example B.2** (Natural numbers from the field axioms).

Since we know that there exists a real number  $1 \in \mathbb{R}$ , we can use addition to define a new number  $2 := 1 + 1 \in \mathbb{R}$ . Similarly we define  $3 := 2 + 1$ ,  $4 := 3 + 1$ ,  $5 := 4 + 1$ ,  $6 := 5 + 1$ , etc. These satisfy the usual properties. As an example, let us verify the following.

*Claim:* We have  $2 + 2 = 4$ .

*Proof:* Calculate, using the definition of 2, the associativity of addition, the definition of 3, and the definition of 4:

$$2 + 2 \stackrel{(\text{def of } 2)}{=} 2 + (1 + 1) \stackrel{(\text{B.2})}{=} (2 + 1) + 1 \stackrel{(\text{def of } 3)}{=} 3 + 1 \stackrel{(\text{def of } 4)}{=} 4 .$$

This proves the claim.  $\square$

**Exercise B.3** (Other exciting properties of natural numbers).

Prove some other exciting properties of these numbers, for example  $2 + 3 = 5$  and  $2 \cdot 3 = 6$ .

**Example B.3** (Yet another consequence of the field axioms).

We next verify a familiar property of real numbers which involves this number  $2 := 1 + 1$  and an arbitrary real number  $x$ .

*Claim:* For any  $x \in \mathbb{R}$  we have  $x + x = 2x$ .

*Proof:* Using the neutral element for multiplication (twice), distributivity, and definition of 2, we get the following expression for  $x + x$ :

$$x + x \stackrel{(\text{B.7})}{=} 1 \cdot x + 1 \cdot x \stackrel{(\text{B.9})}{=} (1 + 1) \cdot x \stackrel{(\text{def of } 2)}{=} 2 \cdot x .$$

The claim follows.  $\square$

**Exercise B.4** (Multiplication by zero).

Prove that for any  $x \in \mathbb{R}$  we have  $0 \cdot x = 0$ .

Ok, you get the point! The familiar facts about the real numbers that you knew since kindergarten can be deduced from the axioms. In order that we get somewhere in this course, we will not actually require you to provide detailed proofs of totally commonplace statements such as  $0 \cdot x = 0$  or  $2 \cdot 2 = 4$  or  $(-1)^2 = 1$  or  $(x - y)^2 = x^2 - 2xy + y^2$  — as long as you realize that *in principle* they should be obtained as logical consequences of the axioms (and familiar notational conventions and definitions).<sup>1</sup>

*Other fields*

There are many other fields besides  $\mathbb{R}$ , and they all satisfy the field axioms. In this course we do not use other fields than  $\mathbb{R}$  and  $\mathbb{Q}$ , but for perspective we mention a few prominent examples of fields:

- the field of rational numbers  $\mathbb{Q}$ ;
- the field  $\mathbb{Q}(\sqrt{5})$  of rational numbers adjoined with a square root of 5;
- the field of complex numbers  $\mathbb{C}$ ;
- the finite field  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  of  $p$  elements, where  $p$  is a prime number;
- the finite field  $\mathbb{F}_{p^k}$  of  $p^k$  elements, where  $p$  is a prime number and  $k \in \mathbb{N}$ ;
- the field  $\mathbb{Q}_p$  of  $p$ -adic numbers, where  $p$  is a prime number;
- the field  $\mathbb{K}(q)$  of rational functions in a single variable  $q$  over another field  $\mathbb{K}$ ;
- ...

In particular all (familiar) consequences of the field axioms hold for these as well: for example if  $z, w \in \mathbb{C}$  are complex numbers, then  $(z + w)^2 = z^2 + 2zw + w^2$ , where  $2 := 1 + 1 \in \mathbb{C}$ . Note, however, that for instance in the two element field  $\mathbb{F}_2$ , we have  $2 = 0$ . So you should not try to derive the property  $2 \neq 0$  of real numbers from the field axioms (you would fail, it is not a logical consequence of them!).

♥ **Order axioms**

Since there are other fields besides just the real numbers, the real numbers must have some further specific properties that distinguish them. Having an order relation  $<$  turns out to be quite special already. To work with the order relation  $<$ , we adopt the usual additional notations such as:  $y > x$  means  $x < y$ ,  $x \leq y$  means that either  $x < y$  or  $x = y$ , etc.

---

<sup>1</sup>Nowadays there are, by the way, formally verified proofs of a vast amount of basic statements about real numbers and even metric space topology; certainly including all statements in this section. What this means is that the statements have been phrased in an entirely formal language, proofs have also been given in the formal language, and a computer has verified that the proofs adhere to the rules of logic and actually demonstrate the statements. See, for example, <https://leanprover-community.github.io/>. I slightly doubt, however, that all of that makes for much more pleasant reading than these lecture notes... (If you think otherwise, I will accept your formally verified solutions to the exam problems!) Moreover, producing the fully formal proofs is so tedious that already in some parts of this undergraduate course we actually get deeper into mathematics than where hundreds of mathematicians and computer scientists have gotten in the past decades with formally verified mathematics. A noble pursuit, nevertheless!

The order axioms of the real number field are:

Alternatives:

$$\forall x, y \in \mathbb{R} : \text{ one and only one of the three relations } x < y, x = y, x > y \text{ is true} \quad (\text{B.10})$$

Transitivity:

$$\forall x, y, z \in \mathbb{R} : x < y \text{ and } y < z \implies x < z \quad (\text{B.11})$$

Compatibility with addition:

$$\forall x, y, z \in \mathbb{R} : x < y \implies x + z < y + z \quad (\text{B.12})$$

Compatibility with multiplication:

$$\forall x, y \in \mathbb{R} : x > 0 \text{ and } y > 0 \implies x \cdot y > 0 \quad (\text{B.13})$$

From these and the field axioms together, one can derive consequences.

**Example B.4** (Some consequences of the order axioms).

*Claim:* We have  $0 < 1$ .

*Proof:* Since we know that  $0 \neq 1$ , there are two mutually exclusive alternatives left:  $0 < 1$  or  $0 > 1$ . Let us prove that the second one is impossible. So assume that  $0 > 1$ . By adding  $-1$  and using compatibility with addition, we get (after simplifying by neutral element of addition and opposite element properties) that  $-1 > 0$ . Then by compatibility with multiplication we get  $(-1)(-1) > 0$ . But from the field axioms one can show that  $(-1)(-1) = 1$ , so this simplifies to  $1 > 0$ . In this case we have both  $1 > 0$  and  $0 > 1$ , which is not allowed by the alternatives. So we have to discard the possibility that  $0 > 1$ , and we conclude that  $0 < 1$  holds.  $\square$

As the example of the field of two elements shows, the property  $2 \neq 0$  of real numbers cannot be derived from the field axioms alone. Having also the order axioms, we can now prove it.

*Claim:* We have  $2 \neq 0$  in  $\mathbb{R}$ .

*Proof:* We already know that  $1 > 0$ . Adding 1 and using compatibility with addition, we find  $2 > 1$ . Transitivity then implies  $2 > 0$ . This rules out the possibility that  $2 = 0$  by the axiom that exactly one of the alternatives  $2 > 0$ ,  $2 = 0$ , and  $2 < 0$ , is true.  $\square$

**Exercise B.5** (Product of two negative numbers).

Prove that the product of any two negative real numbers is positive.

Ok, you get the point again. All the familiar properties you already knew in kindergarten are logical consequences of the axioms.

Note that the rational number field  $\mathbb{Q}$  also satisfies the order axioms. Therefore if  $\mathbb{R} \neq \mathbb{Q}$ , there must be yet something else that distinguishes the real numbers. It is going to be the most profound of the axioms of real numbers.

## B.2. Equivalence of the formulations of the completeness axiom

Recall that in Lecture II, any of the following statements was called the **completeness axiom** of  $\mathbb{R}$ :

**(C1):** Every non-empty subset  $A \subset \mathbb{R}$  which is bounded from above has a least upper bound  $\sup A \in \mathbb{R}$ .

- (C2):** Every increasing real number sequence  $(a_n)_{n \in \mathbb{N}}$  which is bounded from above has a limit  $\lim_{n \rightarrow \infty} a_n \in \mathbb{R}$ .
- (C3):** Every collection  $(I_n)_{n \in \mathbb{N}}$  of closed intervals  $I_n \subset \mathbb{R}$ , which is nested in the sense that  $I_{n+1} \subset I_n$  for every  $n \in \mathbb{N}$ , has a nonempty intersection

$$\bigcap_{n \in \mathbb{N}} I_n \neq \emptyset.$$

Let us now prove the equivalence of (C1), (C2), and (C3). We prove separately the implications

$$(C1) \implies (C2), \quad (C2) \implies (C3), \quad (C3) \implies (C1).$$

The equivalence of all three follows by combining these implications.

*Proof of  $(C1) \Rightarrow (C2)$ .* Assume (C1). Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence of real numbers which is increasing and bounded from above. We must prove that this sequence converges.

Since the sequence  $(a_n)_{n \in \mathbb{N}}$  is bounded from above, the set

$$A = \{a_n \mid n \in \mathbb{N}\}$$

of its values is a subset in  $\mathbb{R}$  which is bounded from above. By assumption (C1), then, this subset has a least upper bound  $t_0 := \sup \{a_n \mid n \in \mathbb{N}\} \in \mathbb{R}$ . We will prove that the sequence  $(a_n)_{n \in \mathbb{N}}$  converges to  $t_0$ .

Let  $\varepsilon > 0$ . Since  $t_0 - \varepsilon$  is not an upper bound for  $\{a_n \mid n \in \mathbb{N}\}$ , there exists some  $n_\varepsilon \in \mathbb{N}$  such that  $a_{n_\varepsilon} > t_0 - \varepsilon$ . Since the sequence is increasing, for all  $n \geq n_\varepsilon$  we must then also have  $a_n > t_0 - \varepsilon$ . On the other hand, since  $t_0$  is an upper bound for  $\{a_n \mid n \in \mathbb{N}\}$ , we have  $a_n \leq t_0$  for all  $n \in \mathbb{N}$ . For  $n \geq n_\varepsilon$  we have thus obtained

$$t_0 - \varepsilon < a_n \leq t_0 < t_0 + \varepsilon,$$

which implies  $|a_n - t_0| < \varepsilon$ . Since such an  $n_\varepsilon \in \mathbb{N}$  was found for an arbitrary  $\varepsilon > 0$ , we have by definition of limits shown that  $\lim_{n \rightarrow \infty} a_n = t_0$ .

Property (C2) is thus established.  $\square$

*Proof of  $(C2) \Rightarrow (C3)$ .* Assume (C2). Let  $(I_n)_{n \in \mathbb{N}}$  be a collection of closed intervals  $I_n = [a_n, b_n] \subset \mathbb{R}$  such that  $I_{n+1} \subset I_n$  for every  $n \in \mathbb{N}$ . We must prove that  $\bigcap_{n \in \mathbb{N}} I_n \neq \emptyset$ .

The nesting condition

$$[a_{n+1}, b_{n+1}] = I_{n+1} \subset I_n = [a_n, b_n]$$

simply amounts to the following inequalities among the endpoints of the intervals:

$$a_n \leq a_{n+1} \leq b_{n+1} \leq b_n.$$

Therefore the sequence  $(a_n)_{n \in \mathbb{N}}$  of the left endpoints of the intervals is increasing. This sequence is also bounded from above, because for any  $n \in \mathbb{N}$  we have  $a_n \leq b_n \leq b_1$  (the second inequality follows from the fact that the sequence of right endpoints is decreasing). By assumption (C2), the sequence  $(a_n)_{n \in \mathbb{N}}$  therefore has a limit  $\alpha = \lim_{n \rightarrow \infty} a_n$ . We will show that  $\alpha \in \bigcap_{n \in \mathbb{N}} I_n$ , and it will follow that the intersection is nonempty.

To show that  $\alpha \in \bigcap_{m \in \mathbb{N}} I_m$ , by definition of intersection we must show that  $\alpha \in I_m$  for all  $m \in \mathbb{N}$ . Since  $I_m = [a_m, b_m]$ , this amounts to proving that

$$a_m \leq \alpha \leq b_m.$$

The first inequality above is clear: for  $n \geq m$  we have  $a_n \geq a_m$  (the left endpoints form an increasing sequence), so by preservation of constant inequalities (Corollary II.15) we have  $\alpha = \lim_{n \rightarrow \infty} a_n \geq a_m$ . To prove the other inequality, we argue by contradiction. Suppose that it is not true, i.e., that  $\alpha > b_m$ . By definition of the limit  $\alpha = \lim_{n \rightarrow \infty} a_n$ , for the positive number  $\varepsilon = \alpha - b_m > 0$ , there exists an  $n_\varepsilon \in \mathbb{N}$  so that for  $n \geq n_\varepsilon$  we have  $|a_n - \alpha| < \varepsilon = \alpha - b_m$ , which implies

$$a_n > \alpha - \varepsilon = \alpha - (\alpha - b_m) = b_m.$$

In particular for  $n \geq \max\{n_\varepsilon, m\}$ , since the sequence of right endpoints is decreasing, we would get  $a_n > b_m \geq b_n$ . This, however, is a contradiction with the ordering  $a_n \leq b_n$  of the left and right endpoints of  $I_n = [a_n, b_n]$ . With this contradiction, we have also established the other claimed inequality. We have thus concluded  $\alpha \in I_m$  for all  $m \in \mathbb{N}$ , which shows that the intersection  $\bigcap_{m \in \mathbb{N}} I_m$  is not empty, establishing (C3).  $\square$

*Proof of (C3)  $\Rightarrow$  (C1).* Assume (C3). Let  $A \subset \mathbb{R}$  be a nonempty subset which is bounded from above. We must prove that  $A$  has a least upper bound.

Since  $A$  is bounded from above, it has an upper bound, and since  $A$  is moreover nonempty, it is possible to choose an upper bound  $b_0 \in \mathbb{R}$  so that the number  $a_0 = b_0 - 1$  is not an upper bound for  $A$ .

Consider the midpoint  $c_0 = \frac{a_0 + b_0}{2}$ . If  $c_0$  is an upper bound for  $A$ , we set  $a_1 = a_0$  and  $b_1 = c_0$ . Otherwise we set  $a_1 = c_0$  and  $b_1 = b_0$ . With such choices, we are guaranteed that  $b_1$  is an upper bound for  $A$ , while  $a_1$  is not.

Continue inductively. When  $b_n$  is an upper bound for  $A$  and  $a_n$  is not, consider the midpoint  $c_n = \frac{a_n + b_n}{2}$ . Depending on whether  $c_n$  is an upper bound for  $A$  or not, set  $a_{n+1} = a_n$  and  $b_{n+1} = c_n$ , or  $a_{n+1} = c_n$  and  $b_{n+1} = b_n$ . By construction we get a decreasing sequence  $(b_n)_{n \in \mathbb{N}}$  of upper bounds for  $A$ , and an increasing sequence  $(a_n)_{n \in \mathbb{N}}$  of numbers that are not upper bounds for  $A$ , and moreover  $a_n < b_n$  for all  $n \in \mathbb{N}$ . The distances are halved at every step, so  $b_n - a_n = 2^{-n}$ .

From the above properties we get that the closed intervals  $I_n = [a_n, b_n]$  for  $n \in \mathbb{N}$  are nested,  $I_{n+1} \subset I_n$  for all  $n \in \mathbb{N}$ . By the assumed property (C3), the intersection  $\bigcap_{n \in \mathbb{N}} I_n$  is then nonempty. On the other hand, the intersection cannot contain two distinct points: if  $x, y \in \bigcap_{n \in \mathbb{N}} I_n$ , then for all  $n \in \mathbb{N}$  we have  $x, y \in I_n = [a_n, b_n]$ , which implies that  $|x - y| \leq b_n - a_n = 2^{-n} \rightarrow 0$ , so  $|x - y| = 0$  and  $x = y$ . We conclude that the intersection contains exactly one point  $\xi$ , i.e., it is a singleton  $\bigcap_{n \in \mathbb{N}} I_n = \{\xi\}$ . From the same estimate we also get that

$$\xi - 2^{-n} \leq a_n \leq \xi \leq b_n \leq \xi + 2^{-n}.$$

By the squeeze theorem (Lemma II.17), this implies  $\lim_{n \rightarrow \infty} a_n = \xi$  and  $\lim_{n \rightarrow \infty} b_n = \xi$ .

Our goal was to show that the set  $A$  has a least upper bound. We will show that  $\xi$  is it.

Let us first verify that  $\xi$  is an upper bound for  $A$ . So let  $x \in A$ . By construction each  $b_n$  is an upper bound for  $A$ , so  $b_n \geq x$  for all  $n \in \mathbb{N}$ . Since we have  $\xi = \lim_{n \rightarrow \infty} b_n$ , from the preservation of bounds (Corollary II.15) we get that  $\xi \geq x$ . This was true for an arbitrary element  $x \in A$ , so indeed  $\xi$  is an upper bound for  $A$ .

Let us then show that no smaller number  $\xi' < \xi$  is an upper bound for  $A$ . Since  $\lim_{n \rightarrow \infty} a_n = \xi$ , for any  $\xi' < \xi$  we have  $a_n > \xi'$  for large enough  $n$  (take  $\varepsilon = \xi - \xi' > 0$  in the definition of limit), and since  $a_n$  was not an upper bound for  $A$ , the smaller number  $\xi' < a_n$  can not be either.

This proves that  $\sup A = \xi$  exists, establishing (C1).  $\square$

### B.3. ♥ Cantor set

The Cantor set, or Cantor's  $\frac{1}{3}$ -set, is a subset  $C \subset [0, 1]$ , which has interesting properties from a topological, set theoretic, and measure theoretic points of view.

Informally, the Cantor set  $C$  is constructed by taking the unit interval  $[0, 1]$ , removing its middle third, then removing the middle third of both of the remaining thirds, and successively always removing the middle third of each remaining interval. Figure B.1 illustrates this iterative construction.





FIGURE B.1. The Cantor set  $C = \bigcap_{n \in \mathbb{N}} C_n \subset [0, 1]$  is constructed iteratively so that from stage  $n$  approximation  $C_n$  one removes the middle thirds of the remaining intervals to obtain stage  $n + 1$  approximation  $C_{n+1}$ .

More formally, the Cantor set  $C$  is defined, using a sequence  $(C_n)_{n \in \mathbb{N}}$  of nested approximating sets, as the intersection

$$C = \bigcap_{n \in \mathbb{N}} C_n, \quad (\text{B.14})$$

where the sets  $C_n$ ,  $n \in \mathbb{N}$ , are as follows. It makes sense to think of the unit interval  $C_0 = [0, 1]$  as the zeroth stage approximation. The first stage approximation is the set  $C_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ ; a union of 2 closed intervals of length  $\frac{1}{3}$ . The second stage approximation is  $C_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$ ; a union of 4 closed intervals of length  $\frac{1}{9}$ . The  $n$ :th stage approximation is the union of  $2^n$  closed intervals of length  $3^{-n}$ , specifically

$$C_n = \bigcup_{b_1, \dots, b_n \in \{0, 1\}} I_{b_1, \dots, b_n}^{(n)} \quad (\text{B.15})$$

where

$$I_{b_1, \dots, b_n}^{(n)} = \left[ 2 \sum_{j=1}^n b_j 3^{-j}, 2 \sum_{j=1}^n b_j 3^{-j} + 3^{-n} \right]. \quad (\text{B.16})$$

The interval (B.16) is exactly obtained by keeping the left or right third, corresponding to  $b_n = 0$  and  $b_n = 1$ , respectively, of the  $n - 1$ :st stage interval  $I_{b_1, \dots, b_{n-1}}^{(n-1)}$ . The middle third of  $I_{b_1, \dots, b_{n-1}}^{(n-1)}$  has been removed in stage  $n$ . In particular, we have the nesting property  $C_n \subset C_{n-1}$  for all  $n \in \mathbb{N}$ , i.e.,

$$C_0 \supset C_1 \supset C_2 \supset C_3 \supset \dots$$

Let us now make some observations regarding the set  $C$  defined as the intersection (B.14).

*The Cantor set is nonempty*

The first observation is that the set  $C$  is non-empty,

$$C \neq \emptyset.$$

There are various ways to see this; the easiest is to just note that  $0 \in C_n$  and  $1 \in C_n$  for every  $n \in \mathbb{N}$ , so in the intersection we at least have  $0 \in C$  and  $1 \in C$ . In other words, at least the two endpoints of the unit interval which constitutes the zeroth stage approximation  $C_0$  belong to the Cantor set  $C$ . By an essentially similar reasoning one can see that all of the  $2^{n+1}$  endpoints of the  $2^n$  intervals of stage  $n$  remain in the intersection — we leave the detailed verification of this as an exercise. When all different stages  $n \in \mathbb{N}$  are considered, such approximating interval endpoints give countably infinitely many different points in  $C$  — we again leave the details as an exercise, because it turns out that there are vastly more points in  $C$ , and they admit a unified description.

*The Cantor set is uncountably infinite*

The unified description of points of  $C$  is by infinite binary sequences  $(b_1, b_2, b_3, \dots)$ .

First, suppose that an infinite binary sequence  $(b_n)_{n \in \mathbb{N}}$  is given, with  $b_n \in \{0, 1\}$  for  $n \in \mathbb{N}$ . Note the nesting of the intervals,

$$I_{b_1}^{(1)} \supset I_{b_1, b_2}^{(2)} \supset I_{b_1, b_2, b_3}^{(3)} \supset \dots \supset I_{b_1, \dots, b_n}^{(n)} \supset I_{b_1, \dots, b_n, b_{n+1}}^{(n+1)} \supset \dots$$

By the formulation (C3) of the completeness axiom, the intersection of these nested closed intervals is nonempty, and since the lengths  $3^{-n}$  of the intervals are tending to 0 as  $n \rightarrow \infty$ , it is not difficult to see that the intersection cannot contain more than one point. Therefore there exists some  $x \in \mathbb{R}$  (which of course depends on the chosen binary sequence  $(b_n)_{n \in \mathbb{N}}$ ) such that

$$\bigcap_{n \in \mathbb{N}} I_{b_1, \dots, b_n}^{(n)} = \{x\}.$$

Since  $x \in I_{b_1, \dots, b_n}^{(n)} \subset C_n$  for every  $n \in \mathbb{N}$ , we then have also

$$x \in \bigcap_{n \in \mathbb{N}} C_n = C.$$

This shows that to any infinite binary sequence  $(b_n)_{n \in \mathbb{N}}$ , we can associate a point  $x$  in the Cantor set  $C$ . We will observe also that if  $(b_n)_{n \in \mathbb{N}}$  and  $(b'_n)_{n \in \mathbb{N}}$  are two different binary sequences, then the corresponding points  $x$  and  $x'$  are different. Indeed then for some  $k \in \mathbb{N}$  we have  $b_k \neq b'_k$ , and the corresponding  $k$ :th stage intervals are disjoint,  $I_{b_1, \dots, b_k}^{(k)} \cap I_{b'_1, \dots, b'_k}^{(k)} = \emptyset$ . But since the corresponding points belong to these disjoint intervals,  $x \in I_{b_1, \dots, b_k}^{(k)}$  and  $x' \in I_{b'_1, \dots, b'_k}^{(k)}$ , we must have  $x \neq x'$ .

**Exercise (#) B.6** (The point in Cantor set corresponding to a binary sequence).

Show that the point  $x$ , corresponding to the binary sequence  $(b_n)_{n \in \mathbb{N}}$ , is given by the formula

$$x = 2 \sum_{j=1}^{\infty} b_j 3^{-j}. \quad (\text{B.17})$$

Use this to give a different proof that different binary sequences yield different points in the Cantor set.

**Remark B.5** (Describing the Cantor set by ternary numbers).

Consider the analogue of decimal expansions  $\sum_{j=1}^{\infty} d_j 10^{-j}$  where  $d_1, d_2, \dots \in \{0, 1, \dots, 8, 9\}$  (see Example II.33), but in base three instead of base ten. These are numbers of the form  $\sum_{j=1}^{\infty} t_j 3^{-j}$  where the “digits” are  $t_1, t_2, \dots \in \{0, 1, 2\}$ .

From the expression (B.17) we see that the points in the Cantor set correspond to base three “decimal expansions” (ternary expansions) which only ever use digits 0 and 2; never 1.

Conversely, we will show that each point of the Cantor set corresponds to some binary sequence as above. Suppose that  $x \in C$ . By the definition of intersection, we then have  $x \in C_n$  for every  $n \in \mathbb{N}$ . Since  $C_n$  is the union of disjoint intervals (B.15), there must exist exactly one choice of  $b_1^{(n)}, \dots, b_n^{(n)} \in \{0, 1\}$  such that  $x \in I_{b_1^{(n)}, \dots, b_n^{(n)}}^{(n)}$ .

Let us now define an infinite sequence  $(b_n)_{n \in \mathbb{N}}$  by setting  $b_n = b_n^{(n)}$  where  $b_n^{(n)}$  came from the above choice of the  $n$ :th stage interval. But in fact  $b_n^{(m)}$  will not depend on which stage  $m \geq n$  we were considering<sup>2</sup>; we have  $b_n^{(m)} = b_n$  for all  $m \geq n$ . Therefore we have  $x \in I_{b_1^{(n)}, \dots, b_n^{(n)}}^{(n)}$  for all  $n \in \mathbb{N}$ , i.e.,  $x \in \bigcap_{n \in \mathbb{N}} I_{b_1^{(n)}, \dots, b_n^{(n)}}^{(n)}$ . But this intersection of nested intervals is exactly the singleton that we used to define the point in  $C$  which corresponds to the binary sequence  $(b_n)_{n \in \mathbb{N}}$ , so we have now shown that any point  $x \in C$  is of this form.

### *On measure theoretic properties of the Cantor set*

While this is not a course in measure theory, it is interesting to note that the Cantor set  $C$ , which is set theoretically large in that it is uncountably infinite, is nevertheless measure theoretically as small as possible: it has *Lebesgue measure zero*. The Lebesgue measure on  $\mathbb{R}$  is the good mathematical generalization of length. It is not difficult to convince oneself that the “length” of the Cantor set  $C$  should indeed be zero, by the following argument. For any  $n \in \mathbb{N}$ , we have  $C \subset C_n$ , i.e., The Cantor set  $C$  is contained in the union of  $2^n$  intervals of length  $3^{-n}$  each. Therefore it is reasonable to say that its length is at most  $2^n 3^{-n} = \left(\frac{2}{3}\right)^n$ . But  $n \in \mathbb{N}$  was arbitrary here, and as  $n \rightarrow \infty$ , this upper bound on the length is tending to 0.

Another measure theoretic point of view to the Cantor set is its fractal dimension. In arbitrary metric spaces one can define Hausdorff measures of arbitrary dimensions  $d \in [0, +\infty)$ , and the Hausdorff dimension of a set is defined as the infimum of dimensions  $d$  such that the  $d$ -dimensional Hausdorff measure vanishes. The set  $C$  has Hausdorff dimension  $\frac{\log 2}{\log 3} \approx 0.6309$ , which is strictly less than the dimension 1 of any nonempty open subset of the real line, but strictly larger than the dimension 0 of countable sets such as  $\mathbb{Q}$ .

---

<sup>2</sup>Suppose  $m \geq n$ . The numbers  $b_1^{(n)}, \dots, b_n^{(n)} \in \{0, 1\}$  are chosen so that  $x \in I_{b_1^{(n)}, \dots, b_n^{(n)}}^{(n)}$ . Likewise, the numbers  $b_1^{(m)}, \dots, b_m^{(m)} \in \{0, 1\}$  are chosen so that  $x \in I_{b_1^{(m)}, \dots, b_m^{(m)}}^{(m)}$ . But we have the nesting property: an  $m$ :th stage interval is contained in exactly one  $n$ :th stage interval

$$I_{b_1^{(m)}, \dots, b_m^{(m)}}^{(m)} \subset I_{b_1^{(n)}, \dots, b_n^{(n)}}^{(n)}.$$

By disjointness of the  $n$ :th stage intervals, we must have  $b_1^{(m)} = b_1^{(n)}, \dots, b_n^{(m)} = b_n^{(n)}$ .

### *On topological properties of the Cantor set*

We mention without entering details a few interesting topological properties of the Cantor set  $C \subset \mathbb{R}$ . These require notions from later lectures, and are perhaps best treated as additional exercises at the suitable point in the course.

After Lecture XI you can do the following.

**Exercise B.7** (The Cantor set is compact).

Prove that the Cantor set  $C$  is compact.

After the definitions in Lecture V, it is possible to do the following, with concepts from Lecture XI this probably becomes easier.

**Exercise (#) B.8** (The Cantor set is nowhere dense).

Prove that the Cantor set  $C$  has the property that any nonempty open set  $V \subset \mathbb{R}$  contains a nonempty open set  $U \subset V \subset \mathbb{R}$  such that  $U \cap C = \emptyset$ .

The set  $C$  is in this topological sense small; smaller even than the countable but dense set  $\mathbb{Q}$  of rational numbers, which it intersects every nonempty open set  $U \subset \mathbb{R}$  (which can be seen from Theorem II.21).

After Lecture XII you can do the following.

**Exercise (#) B.9** (The Cantor set is totally disconnected).

Prove that all connected components of the Cantor set  $C$  are singletons.

## **B.4. ♥ Applications of supremum and infimum**

The notions of supremum and infimum may seem a bit abstract. Indirectly, via a different formulation of the completeness axiom, we saw that supremum is important for instance for the existence of real numbers with given decimal expansions (Example II.33). But it is good to realize that supremum is also quite directly used in many common constructions in analysis. We describe a few examples here, but the intention is not to elaborate on the details in full.

### ♥ Arc length of a curve

One relatively simple, intuitive, and commonly used application of supremum is the length of a curve; also called *arc length*. Here we describe the definition in the context of curves in a metric space  $(X, d)$ , which is introduced Lecture V. Also the notion of continuity from Lecture VI is briefly invoked.

It is also possible to treat this as an example without the general definitions, however. The reader who so prefers may consider curves in the familiar  $d$ -dimensional space  $X = \mathbb{R}^d$ , and interpret the distances between two points  $\vec{x} = (x_1, \dots, x_d)$  and  $\vec{y} = (y_1, \dots, y_d)$  in  $\mathbb{R}^d$  as given by the usual formula

$$d(\vec{x}, \vec{y}) = \sqrt{(y_1 - x_1)^2 + \dots + (y_d - x_d)^2}.$$

A (parametrized) curve in  $X$  is a continuous function from the unit interval  $[0, 1]$  to the space  $X$ ,

$$\gamma: [0, 1] \rightarrow X.$$

For a given subdivision

$$0 = t_0 < t_1 < t_2 < \cdots < t_n = 1$$

of the unit interval, it is natural to interpret the expression

$$\sum_{j=1}^n d(\gamma(t_{j-1}), \gamma(t_j)) = d(\gamma(t_0), \gamma(t_1)) + \cdots + d(\gamma(t_{n-1}), \gamma(t_n))$$

as an approximation of the length of the curve  $\gamma$ . Under refinement of the subdivision, the approximate length never decreases. Therefore the natural notion of the length of the curve, as obtained in the limit of refinements of subdivisions, is the supremum of all the approximating expressions: the **length** of the curve  $\gamma$  is *defined* as

$$L(\gamma) := \sup \left\{ \sum_{j=1}^n d(\gamma(t_{j-1}), \gamma(t_j)) \mid n \in \mathbb{N}, 0 = t_0 < t_1 < \cdots < t_n = 1 \right\}. \quad (\text{B.18})$$

In  $\mathbb{R}^d$ , for differentiable (or at least piecewise differentiable) curves  $\gamma$ , one can show that this definition of length coincides with the integral  $L(\gamma) = \int_0^1 \|\gamma'(t)\| dt$ , but the integral is in fact more complicated to define precisely (see below) and it exists less generally than (B.18): it cannot be defined in general metric spaces, and even in  $\mathbb{R}^d$  it requires differentiability assumptions, which depend on the chosen parametrization, etc. The straightforward definition (B.18) using supremum is clearly good!

### ♥ Infinite sums with non-negative terms

If we have an arbitrary collection  $(s_i)_{i \in I}$  of nonnegative terms  $s_i \geq 0$ , then there is a quick and rather simple way to give a meaning to the sum

$$\sum_{i \in I} s_i.$$

This is rather remarkable, since we can allow the index set  $I$  to be arbitrarily large, meaning that the number of terms in the sum can be finite, countably infinite, or even uncountably infinite.

Of course a partial sum that includes only finitely many terms  $s_{i_1}, \dots, s_{i_k}$ , with  $i_1, \dots, i_k \in I$  distinct, is trivially defined as

$$s_{i_1} + \cdots + s_{i_k}.$$

The idea is that the other terms omitted are non-negative, so the “true sum” must be at least as large. In other words the “true sum” should be an upper bound for the set of all finite partial sums. The definition

$$\sum_{i \in I} s_i := \sup \left\{ s_{i_1} + \cdots + s_{i_k} \mid k \in \mathbb{N}_0, i_1, \dots, i_k \in I \text{ distinct} \right\}$$

declares the “true sum” as the least upper bound for the set of all finite partial sums — i.e., it requires that the gap in between the finite partial sums and the “true sum” can be made arbitrarily small by including enough terms.

The set of all finite partial sums is nonempty (we always have the possibility of including  $k = 0$  terms in the partial sum), but it may or may not be bounded, so we obviously have to allow the possibility  $\sum_{i \in I} s_i = +\infty$  (in keeping with the convention of supremum of unbounded sets) if arbitrarily large finite partial sums exist.

This definition also has the advantage that it is clear that the sum does not depend on the “order of elements” (in fact we did not even require the index set  $I$  to be ordered!).

But to be fair, this definition crucially relies on the terms being nonnegative. It does not admit easy generalizations to infinite sums with terms of both signs.<sup>3</sup>

**Exercise B.10** (♥ Uncountably many strictly positive terms yields infinite sum).

Let  $(s_i)_{i \in I}$  be a collection of nonnegative terms  $s_i \geq 0$ . Consider the sum  $\sum_{i \in I} s_i$ .

- (a) Suppose that  $\sum_{i \in I} s_i < +\infty$ . Show that for any  $m \in \mathbb{N}$ , there can only exist finitely many indices  $i \in I$  such that  $s_i \geq \frac{1}{m}$ .
- (b) Suppose again that  $\sum_{i \in I} s_i < +\infty$ . Show that there can only exist countably many indices  $i \in I$  such that  $s_i > 0$ .

*Hint: Consider all different  $m \in \mathbb{N}$  in (a). Recall that countable unions of finite sets are countable.*

- (c) Now prove that if there are uncountably infinitely many indices  $i$  corresponding to a strictly positive term  $s_i > 0$ , then we have  $\sum_{i \in I} s_i = +\infty$ .

*Hint: Contrapositive of (b).*

## ♥ Riemann integration

Let

$$f: [a, b] \rightarrow \mathbb{R}$$

be a real valued function defined on a closed interval  $[a, b]$ . Assume also that  $f$  is bounded in the sense that there exists some  $M \in \mathbb{R}$  such that  $|f(x)| \leq M$  for all  $x \in [a, b]$ .

Consider a finite subdivision of the interval  $[a, b]$ , consisting of points

$$a = x_0 < x_1 < x_2 < \cdots < x_n = b.$$

On the subdivision interval  $[x_{j-1}, x_j]$  for  $j \in \{1, \dots, n\}$ , the values of the function have the greatest lower bound and least upper bound

$$s_j = \inf \left\{ f(x) \mid x_{j-1} \leq x \leq x_j \right\} \quad \text{and} \quad t_j = \sup \left\{ f(x) \mid x_{j-1} \leq x \leq x_j \right\}.$$

Using these, we define the lower and upper Riemann sums

$$S = \sum_{j=1}^n s_j (x_j - x_{j-1}) \quad \text{and} \quad T = \sum_{j=1}^n t_j (x_j - x_{j-1})$$

associated with the subdivision  $a = x_0 < x_1 < x_2 < \cdots < x_n = b$ .

It is not particularly difficult to show that the lower and upper Riemann sums satisfy the following:

---

<sup>3</sup>Somewhat more general sums (and integrals) are studied in measure theory.

- If a subdivision  $a = x'_0 < x'_1 < x'_2 < \dots < x'_m = b$  is a refinement of the above subdivision (i.e.  $\{x_0, x_1, \dots, x_n\} \subset \{x'_0, x'_1, \dots, x'_m\}$ ) and  $S'$  and  $T'$  are the associated lower and upper Riemann sums, then  $S \leq S'$  and  $T' \leq T$ .
- If  $\tilde{S}$  and  $\tilde{T}$  are the lower and upper Riemann sums associated with any subdivision, then  $\tilde{S} \leq T$  (and symmetrically  $S \leq \tilde{T}$ ).

**Exercise B.11** (♥ Proof of the properties of upper and lower Riemann sums).  
Prove the two claims above.

Since increasingly fine subdivisions lead to larger lower Riemann sums, it is natural to define the lower integral  $I_-(f)$  as the least upper bound for the set of all lower Riemann sums for all subdivisions; or explicitly

$$I_-(f) := \sup \left\{ \sum_{j=1}^n \left( \inf_{x \in [x_{j-1}, x_j]} f(x) \right) (x_j - x_{j-1}) \mid a = x_0 < x_1 < \dots < x_n = b \right\}.$$

Similarly, since increasingly fine subdivisions lead to smaller upper Riemann sums, it is natural to define the upper integral  $I_+(f)$  as the greatest lower bound for the set of all upper Riemann sums for all subdivisions; or explicitly

$$I_+(f) := \inf \left\{ \sum_{j=1}^n \left( \sup_{x \in [x_{j-1}, x_j]} f(x) \right) (x_j - x_{j-1}) \mid a = x_0 < x_1 < \dots < x_n = b \right\}.$$

From the earlier observations one straightforwardly gets  $I_-(f) \leq I_+(f)$ .

**Definition B.6** (Riemann integral).

A function  $f: [a, b] \rightarrow \mathbb{R}$  is said to be **(Riemann) integrable**, if  $I_-(f) = I_+(f)$ .  
In this case we define its **(Riemann) integral** as

$$\int_a^b f(x) \, dx := I_-(f) = I_+(f).$$

From characterizations of supremum and infimum, it is quite straightforward to prove that

**Theorem B.7** (Characterization of Riemann integrability).

*A function  $f: [a, b] \rightarrow \mathbb{R}$  is Riemann integrable if and only if for every  $\varepsilon > 0$  there exists a subdivision such that the associated upper and lower Riemann sums satisfy  $T - S < \varepsilon$ .*

**Exercise B.12** (♥ Proof of Theorem B.7).  
Prove the above theorem.

To properly define the familiar integral, therefore, we end up using the notions of supremum and infimum — in fact many times.





## Index

- 0/1-metric, 80
- absolute value, 25
- angle, 69
- Archimedean property, 22
- associativity, 7
- Banach space, 147
- Banach's fixed point theorem, 143
- bijective, 12
- bilinearity
  - of inner product, 64
- bold, 1
- boundary, 90
- boundary point, 90
- bounded
  - sequence in a metric space, 119
  - set in a metric space, 83
- bounded from above
  - set of numbers, 38
- bounded from above (sequence), 28
- bounded from below
  - set of numbers, 38
- bounded from below (sequence), 28
- bounded function, 54
- bounded sequence, 28
- cardinality, 157, 158
  - equal, 158
- Cartesian product, 9, 10
- Cauchy sequence, 135
- challenging, v
- clopen set, 89
- closed ball, 82
- closed interval, 6
- closed set, 88
- closure, 91
- codomain, 10
- commutativity, 7
- complement, 7
- complete metric space, 138
- completeness, 138
- completeness axiom, 40, 166
- component function, 115
- composition, 13
- continuity at a point, 94
- continuity of a function, 48
- continuous function, 94
- contraction, 143
- contraction mapping principle, 143
- contrapositive, 21
- converge, 117
- converge to, 117
- convergence
  - of a real number sequence, 29
  - of function series, 131
- convergent, 117
- countable, 159
- countably infinite, 159
- counterexample, 22
- De Morgan's laws, 7
- decimal number, 41
- decreasing sequence, 28
- diagonal argument, 161
- diameter
  - of a set in a metric space, 83
- discrete topology, 87
- distance
  - between subsets, 84
- distance on the real line, 26
- distributivity, 7
- domain, 10
- element, 1, 2
- empty set, 3
- equivalence
  - logic, 20
- Euclidean space, 65, 79
- evaluation, 97
- existential quantifier, 18
- exterior point, 90
- finite set, 157
- fixed point, 142
- free variable, 17
- function, 10
- function sequence, 125
- function series, 131
- greatest lower bound, 39
- Hilbert space, 147
- homeomorphism, 107
- homogeneity

- of norm, 71
- identity function, 11
- if, 20
- if . . . , then, 19
- if and only if, 20
- image, 14
- implication, 19
- increasing sequence, 28
- infimum, 39
- infinite set, 157
- inherited metric, 80
- injective, 12, 21
- inner product, 64
- inner product space, 65
- integrable
  - in the sense of Riemann, 175
- integral
  - in the sense of Riemann, 175
- interior point, 90
- intersection, 7, 8, 18
- interval, 6
- inverse function, 12, 13
- isolated point, 87
- iterate, 142
  
- $\ell^1$  norm, 73
- $\ell^\infty$  norm, 72
- $\ell^p$  norm, 74
- $L^1$ -norm, 75
- largest element
  - of a set of real numbers, 39
- least upper bound, 39
- lemma of two policemen, 32
- length, 173
- lengthy, v
- limit, 117
  - of a real number sequence, 29
- linearity
  - of inner product, 64
- Lipschitz, 95
- Lipschitz constant, 95
- lower bound, 28
  - for a set of real numbers, 38
  
- mapping, 10
- maximum
  - of a set of real numbers, 39
- maximum of a function, 54
- metric, 77
  - induced by a norm, 79
- metric space, 78
- minimum
  - of a set of real numbers, 39
- minimum of a function, 54
- monotone sequence, 28
  
- n-tuple, 10
- negation, 20
  
- neighborhood, 85
- non-degeneracy
  - of inner product, 64
  - of norm, 72
- norm, 67, 71
  - induced by inner product, 67
- normed space, 72
  
- only if, 19
- open ball, 82
- open interval, 6, 7
- open set, 85
- optional, v
- ordered pairs, 9
- orthogonal
  - collection, 69, 70
  - vectors, 69
  
- pair, 10
- pointwise addition
  - of functions, 62
- pointwise convergence, 126
  - of function series, 131
- pointwise product of functions, 51
- pointwise quotient of functions, 51
- pointwise scalar multiplication, 62
- pointwise sum
  - of functions, 62
- pointwise sum of functions, 51
- positive definite, 66
- positive semi-definite, 66
- positive semi-definiteness
  - of inner product, 64
- preimage, 15
- projection, 114
- punctured space, 86
  
- quadruple, 10
- quintuple, 10
  
- range, 10
- range of a function, 54
- real vector space, 60
- restriction, 14
- Riemann integrable, 175
- Riemann integral, 175
- routine, v
  
- sandwich principle, 32
- scalar, 59
- self map, 141
- sentence, 17
- sequence, 117
  - of real numbers, 27
- sequence in a subset, 119
- set, 1, 2
- set difference, 7
- sextuple, 10
- singleton, 3

- smallest element
  - of a set of real numbers, 39
- squeeze theorem, 32
- standard metric on  $\mathbb{R}$ , 78
- strictly decreasing sequence, 28
- strictly increasing sequence, 28
- subsequence, 46
- subset, 4, 18, 19
- successive iterate, 142
- supremum, 39
- supremum norm, 75
- surjective, 12
- symmetricity
  - of inner product, 64
  
- tends to, 117
- topologically equivalent metrics, 111
- transitive, 5
- transitivity, 166
- triangle inequality, 26
  - of norm, 71
- triple, 10
- tuple, 10
- two policemen
  - lemma of, 32
  
- unbounded interval, 6
- uncountable, 159
- uncountably infinite, 159
- uniform convergence, 126
  - of function series, 131
- uniform norm, 75
- union, 7, 8, 18
- universal quantifier, 18
- upper bound, 28
  - for a set of real numbers, 38
  
- vector, 59
- vector space over  $\mathbb{R}$ , 60
- vector subspace, 62
  
- Weierstrass'  $M$  test, 131



## References

### Bibliography

- [Car00] N. L. Carothers. *Real analysis*. Cambridge University Press, 2000.
- [Hal74] Paul R. Halmos. *Naïve set theory*. Undergraduate Texts in Mathematics. Springer, 1974.
- [Rud76] W. Rudin. *Principles of mathematical analysis*. McGraw-Hill, 1976.
- [Väi99] Jussi Väisälä. *Topologia I*. Limes ry., 1999.