

Supplemental Information

Within-session learning in DishBrain is likely explained by asymmetric feedback rather than by unpredictability

Konstantin Kladko

Document S1. Methods of the re-analysis and statistics, derivation of the stationary density, and the closed-loop model.

Data and code

Data: per-rally gameplay and Rest data of Kagan et al. (2022), Open Science Framework, <https://doi.org/10.17605/osf.io/5u6qv>, file `in_vitro_cells_sentience_2.pkl`. Code: <https://github.com/kladkogex/20-watt-computer>, folder `papers/dishbrain-scrambler:reanalysis/kagan2022_reanalysis.py` (all statistics), `reanalysis/figure_data.py` (Figure 1A), `scrambler.py` (closed-loop model, Figure 1B; standard-library Python).

Re-analysis

Each row of the data file is one rally, with chip, session, time in session, number of hits and condition. Groups 0, 1 and 2 are Stimulus, Silent (tagged “no_feedback”) and No feedback (tagged “open_loop”); each chip’s Rest session, without stimulation, is marked `control = 1`. T1 and T2 are the first 5 and the remaining 15 minutes of gameplay (column `half`, as in the authors’ analysis notebook). The unit of analysis is the chip (culture). For each chip, hits per rally is the mean over all its rallies in T1, in T2 and at Rest; chips without both T1 and T2 are excluded, leaving 24 Stimulus, 17 Silent and 15 No feedback cultures. Cell line is read from the recording tag: Stimulus 12 human GFP-iPSC, 7 human NGN2 and 5 primary mouse cultures; Silent 11 and 7 (one chip has recordings under both human tags, so these sum to 18 for 17 chips) and no mouse cultures; No feedback 2, 6 and 7. Mean within-session gains by cell line (hits per rally): Stimulus +0.123, +0.127, +0.257; Silent −0.055, +0.005; No feedback +0.005, −0.014, +0.007.

Statistics

Within-condition changes T2–T1 were tested with two-sided Wilcoxon signed-rank tests over cultures; differences between conditions with Kruskal–Wallis tests: T2 levels over the 56 cultures above, Rest baselines over every chip with a Rest session. Gameplay relative to Rest (T2 minus Rest) is +0.173, +0.040 and −0.012 hits per rally for Stimulus, Silent and No feedback. Values in Figure 1A are mean $\pm$ SEM over cultures. No correction for multiple comparisons was applied; the three within-condition tests give $p = 0.0002$, 0.35 and 1.

Stationary density

Let ψ be the network’s setting in a bounded region, $P(\psi)$ its miss probability, and let each approach move ψ by a random, symmetric step of variance σ_m^2 after a miss and σ_h^2 after a hit, the step being set by the outcome at the point of departure. For small steps the density obeys $\partial_t \rho = \nabla^2(D\rho)$ with $D = \frac{1}{2}[\sigma_m^2 P + \sigma_h^2(1 - P)]$ (Schnitzer, 1993). With a reflecting boundary the stationary flux $-\nabla(D\rho)$

vanishes, so $\rho \propto 1/D \propto 1/[\kappa + (1 - \kappa)P]$ with $\kappa = \sigma_h^2/\sigma_m^2$. Writing $w = 1/[\kappa + (1 - \kappa)P]$ and $\mathbb{E}_U$ for the uniform average, the long-run miss rate is $\mathbb{E}_U[Pw]/\mathbb{E}_U[w]$. For $\kappa < 1$, w decreases with P and Chebyshev's association inequality gives $\mathbb{E}_U[Pw] \leq \mathbb{E}_U[P] \mathbb{E}_U[w]$: the miss rate is below the uniform average $\mathbb{E}_U[P]$, the miss rate of a random setting. For $\kappa > 1$ the inequality reverses, and for $\kappa = 1$ the density is uniform. Since $D = \frac{1}{2}\sigma_m^2[\kappa + (1 - \kappa)P]$, changing σ_m at fixed κ rescales time only: it changes the rate of convergence, not the stationary state. A Schrödinger and a Boltzmann form of these dynamics are given in the companion preprint.

Closed-loop model

Court 1×1 ; paddle half-length 0.1 on the right wall; steps of 20 ms. Each serve starts at the far wall with horizontal speed 1 and vertical speed uniform in $[-0.8, 0.8]$; the ball reflects from the other walls. The ball's height relative to the paddle selects one of 8 channels, $k = \lfloor 8(y - p + \frac{1}{2}) \rfloor$, and its position x sets the rate $s = (4 + 36x)/40$, as in DishBrain's place and rate code. Two motor rates are $[W_{\uparrow, \downarrow}[k]s + \xi]_+$ with Gaussian noise (SD 0.35 per step), and the paddle moves by $1.5(r_{\uparrow} - r_{\downarrow})$ court heights per second. The 16 weights start uniform in $[-1, 1]$ and reflect at ± 1 . After every approach each weight receives a Gaussian kick of SD 0.01, then one of SD σ_h after a hit or $\sigma_m = 0.08$ after a miss; $\kappa = (\sigma_h/0.08)^2$. Each of 400 simulated cultures per value of κ plays one session of 2000 approaches; performance is the hit fraction over approaches 501–2000, averaged over cultures. The information-free level is the performance at $\kappa = 1$.

Spiking-network model

The network model referred to in the main text (leaky integrate-and-fire neurons on the 3.85×2.10 mm electrode array with the published electrode layout, stimulation programs, decoder and session timing, calibrated to eight statistics of the 168 Rest recordings of the human cultures) and the measurement of κ under activity-gated and spike-timing-dependent plasticity are described in the companion preprint; code in folder `network`. The 3000-neuron model (375 neurons per mm^2) and re-calibrated versions with 10^5 , 3×10^5 and 10^6 neurons on the same array (up to 1.2×10^5 per mm^2) all lack a sensory-to-motor path: single pulses at the sensory electrodes excite the motor regions in at most 3% of cases and suppress them in up to 97%, and 500-ms place-code trains at the rate of play give no consistent up-minus-down drive ($|z| > 3$ in 3.5–6.2% of cases; the sign of an electrode's effect agrees across cultures in 22–32% of cases). The small bias that does appear is reproducible across independent topologies and would steer the paddle away from the ball. The model therefore measures κ under the published stimulation programs and session timing, but closed-loop learning cannot be tested in it. The model perturbs synapses isotropically; real stimulation need not, since a patterned 100 Hz burst can induce pathway-specific potentiation and depression in cultured networks (Jimbo et al., 1999). The displacement analysis proposed in the main text measures such directed effects directly.

Supplemental references

Jimbo, Y., Tateno, T., and Robinson, H.P.C. (1999). Simultaneous induction of pathway-specific potentiation and depression in networks of cortical neurons. *Biophys. J.* 76, 670–678.

Schnitzer, M.J. (1993). Theory of continuum random walks and application to chemotaxis. *Phys. Rev. E* 48, 2553–2568.