

# Gradients in Logistic Regression and Perceptrons Supplement

CS114B Lab 4


Kenneth Lai

February 10, 2023

# Gradients in Logistic Regression

- ▶ Cross-entropy loss  $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$

# Graphical Representation of a Linear Classifier (1)



# Gradients in Logistic Regression

- ▶ Cross-entropy loss  $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph:  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$

# Gradients in Logistic Regression

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$$

# Gradients in Logistic Regression

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} x_j$$

$$\blacktriangleright \frac{\partial z}{\partial w_j} = x_j$$

# Gradients in Logistic Regression

- ▶  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \hat{y}(1 - \hat{y}) x_j$ 
  - ▶  $\frac{\partial z}{\partial w_j} = x_j$
  - ▶ For the logistic function:  $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$

# Gradients in Logistic Regression

- ▶  $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j$ 
  - ▶  $\frac{\partial z}{\partial w_j} = x_j$
  - ▶ For the logistic function:  $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
  - ▶ For the cross-entropy loss:  $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$



# Gradients in Logistic Regression

- ▶  $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j = (\hat{y} - y) x_j$ 
  - ▶  $\frac{\partial z}{\partial w_j} = x_j$
  - ▶ For the logistic function:  $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
  - ▶ For the cross-entropy loss:  $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$

# Gradients in Logistic Regression

- ▶  $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j = (\hat{y} - y) x_j$ 
  - ▶  $\frac{\partial z}{\partial w_j} = x_j$
  - ▶ For the logistic function:  $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
  - ▶ For the cross-entropy loss:  $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$
- ▶  $\frac{\partial L}{\partial b} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y})(1) = \hat{y} - y$ 
  - ▶  $\frac{\partial z}{\partial b} = 1$
  - ▶ ...

# Gradients in Perceptrons

- ▶ Perceptron loss  $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph:  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$

# Gradients in Perceptrons

- ▶ Perceptron loss  $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph:  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$ 
  - ▶ Problem:  $\frac{\partial \hat{y}}{\partial z} = 0$  almost everywhere
    - ▶ A change in  $z$  does not result in a change in  $\hat{y}$  unless you are at the threshold

# Gradients in Perceptrons

- ▶ Perceptron loss  $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph:  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$ 
  - ▶ Problem:  $\frac{\partial \hat{y}}{\partial z} = 0$  almost everywhere
    - ▶ A change in  $z$  does not result in a change in  $\hat{y}$  unless you are at the threshold
  - ▶ Solution: consider  $\frac{\partial L}{\partial z}$  directly

# Gradients in Perceptrons

- ▶ Perceptron loss  $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute  $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus:  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph:  $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial w_j}$

# Gradients in Perceptrons

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial w_j}$$

# Gradients in Perceptrons

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} x_j$$

$$\blacktriangleright \frac{\partial z}{\partial w_j} = x_j$$



# Gradients in Perceptrons

- ▶  $\frac{\partial L}{\partial w_j} = (\hat{y} - y)x_j$

- ▶  $\frac{\partial z}{\partial w_j} = x_j$

- ▶ For the perceptron loss:  $\frac{\partial L}{\partial z} = \hat{y} - y$

# Gradients in Perceptrons

- ▶  $\frac{\partial L}{\partial w_j} = (\hat{y} - y)x_j$ 
  - ▶  $\frac{\partial z}{\partial w_j} = x_j$
  - ▶ For the perceptron loss:  $\frac{\partial L}{\partial z} = \hat{y} - y$
- ▶  $\frac{\partial L}{\partial b} = \hat{y} - y$ 
  - ▶  $\frac{\partial z}{\partial b} = 1$
  - ▶ ...