

3. First bud production

Contents

1	Environment preparation	2
2	Data formatting	2
3	Survival object	3
4	Proportional hazards assumption check : Schoenfeld residuals	3
5	AIC comparison of different distributions	4
6	Model selection	4
6.1	Random effects selection	4
6.2	Fixed effects selection	5
6.3	Goodness-of-fit evaluation	6
6.4	Selected model summary	7
7	Median times	14
7.1	By sleep condition	14
7.2	By lineage	14
7.3	By sleep condition and lineage	15
7.4	By tumoral state	16
8	Cumulative incidences	16
8.1	By week and sleep condition	16
8.2	By week and lineage	17
8.3	By week, sleep condition, and lineage	18
8.4	By week and tumoral state	20
9	Cumulative proportions	21
9.1	By week and sleep condition	21
9.2	By week and lineage	22
9.3	By week, sleep condition, and lineage	23
9.4	By week and tumoral state	25

10 Plots of cumulative incidence of first bud production	26
10.1 Common themes	26
10.2 Panel A: interaction between sleep condition and lineage	27
10.2.1 HO_SPT	27
10.2.2 HO_VLN	27
10.2.3 HC_MT	28
10.3 Panel B: by tumoral state	28
10.4 Panel C: by lineage	29
10.4.1 HO_MT	29
10.4.2 HO_SPC	29
10.4.3 HO_SPT	29
10.4.4 HO_VLN	30
10.4.5 HC_MT	30
10.4.6 HV_GAL	30
10.5 Assembly	31

1 Environment preparation

Run scripts with the needed packages, functions, and the raw data frame.

```
source("0a_packages.R")
source("0b_script_data.R")
source("0c_functions.R")
```

2 Data formatting

Create a new data frame adapted for survival analysis.

```
FB_df <-

  #' Load the raw data frame.
  df %>%

  #' Define the follow-up time for each individual:
  #' if the hydra produced a first bud -> time of first bud production (event),
  #' if the hydra died -> time of death (censor),
  #' if the hydra did not produce a first bud and stayed alive
  #' -> 42 days (no event, censor at the end of the experiment).
  mutate(age_followup_days = case_when(
    FB_state == 1 ~ age_FB,
    FB_state == 0 & death_state == 1 ~ age_death,
    TRUE ~ 42),
```

```

age_followup = age_followup_days / 7,
FB_event = FB_state) %>%

#' Build a competing-risk variable for the cumulative incidence functions:
#' if the hydra did not produce a first bud and stayed alive -> 0 (no event),
#' if the hydra produced a first bud -> 1 (event),
#' if the hydra died -> 2 (competing risk of death).
mutate(status_competing = factor(case_when(FB_event == 1 ~ 1,
                                           death_state == 1 ~ 2,
                                           TRUE ~ 0))) %>%

#' Build the variable tum_state_early: an individual is considered
#' tumoral only if the tumor appeared before or at the same time as the
#' supernumerary tentacle event.
mutate(age_tum_weeks = age_tum / 7,
       tum_state_early = case_when(tum_state == "Tumoral" &
                                   !is.na(age_tum_weeks) &
                                   age_tum_weeks <= age_followup ~ "Tumoral",
                                   TRUE ~ "Healthy"),
       tum_state_early = factor(tum_state_early, levels = c("Healthy", "Tumoral"))) %>%

#' Remove missing data.
filter(!is.na(age_followup)) %>%

#' Keep only the variables needed for the analysis.
select(age_followup, FB_event, status_competing, sleep_condition, lineage,
       tum_state_early, parent_unique, replicate_unique)

```

Set reference levels for the categorical variables.

```

FB_df$lineage <- relevel(factor(FB_df$lineage), ref = "HQ_MT")
FB_df$sleep_condition <- relevel(factor(FB_df$sleep_condition), ref = "Control")

```

3 Survival object

Create a survival object with follow-up time and event indicator.

```

FB_surv_object <- Surv(FB_df$age_followup, FB_df$FB_event)

```

4 Proportional hazards assumption check : Schoenfeld residuals

Fit a preliminary full Cox model (fixed effects only).

```

FB_cox <- coxph(Surv(age_followup, FB_event) ~ sleep_condition * lineage *
               tum_state_early, data = FB_df)

```

Test the proportional hazards assumption.

```
print(cox.zph(FB_cox))
```

```
##                chisq df      p
## sleep_condition    5.55  1  0.019
## lineage           131.26  5 < 2e-16
## tum_state_early    4.39  1  0.036
## sleep_condition:lineage  55.59  5 9.9e-11
## sleep_condition:tum_state_early  6.19  1  0.013
## lineage:tum_state_early  5.44  2  0.066
## sleep_condition:lineage:tum_state_early  2.50  1  0.114
## GLOBAL            176.70 16 < 2e-16
```

The proportional hazards assumption is violated for at least one term, an accelerated failure time mixed-effects model will be used.

5 AIC comparison of different distributions

Compare intercept-only accelerated failure time models fitted with different parametric distributions to identify the best-fitting one.

```
FB_dist_weib <- SurvregME(Surv(age_followup, FB_event) ~ 1, data = FB_df,
                          dist = "weibull")

FB_dist_lnorm <- SurvregME(Surv(age_followup, FB_event) ~ 1, data = FB_df,
                           dist = "lognormal")

FB_dist_llog <- SurvregME(Surv(age_followup, FB_event) ~ 1, data = FB_df,
                           dist = "loglogistic")

FB_dist_exp <- SurvregME(Surv(age_followup, FB_event) ~ 1, data = FB_df,
                          dist = "exponential")

AIC(FB_dist_weib, FB_dist_lnorm, FB_dist_llog, FB_dist_exp) %>% arrange(AIC)
```

```
##          df      AIC
## FB_dist_lnorm  2 2521.574
## FB_dist_llog   2 2530.357
## FB_dist_weib   2 2634.431
## FB_dist_exp    1 2648.340
```

The model with a log-normal distribution has the lowest AIC, we'll use it for the models.

6 Model selection

6.1 Random effects selection

With the full fixed-effect structure held constant, compare different random-effect structures to find the optimal one.

```

FB_R_0 <- SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage * tum_state_early,
  data = FB_df, dist = "lognormal")

FB_R_1 <- SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage * tum_state_early +
  (1 | parent_unique),
  data = FB_df, dist = "lognormal")

FB_R_2 <- SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage * tum_state_early +
  (1 | replicate_unique),
  data = FB_df, dist = "lognormal")

FB_R_3 <- SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage * tum_state_early +
  (1 | parent_unique) + (1 | replicate_unique),
  data = FB_df, dist = "lognormal")

AIC(FB_R_0, FB_R_1, FB_R_2, FB_R_3) %>% arrange(AIC)

```

```

##      df      AIC
## FB_R_2 26 2041.720
## FB_R_3 27 2043.720
## FB_R_0 25 2197.253
## FB_R_1 26 2199.253

```

The model with the replicate id as a random effect (FB_R_2) has the lowest AIC, this random effect structure is therefore selected.

6.2 Fixed effects selection

With the selected random-effect structure fixed, compare different fixed-effect structures to find the optimal one.

```

FB_F_0 <- SurvregME(Surv(age_followup, FB_event) ~ 1 + (1 | replicate_unique), data =
FB_df, dist = "lognormal")
FB_F_1 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_2 <- SurvregME(Surv(age_followup, FB_event) ~ lineage + (1 | replicate_unique), data
= FB_df, dist = "lognormal")
FB_F_3 <- SurvregME(Surv(age_followup, FB_event) ~ tum_state_early + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_4 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition + lineage + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_5 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition + tum_state_early + (1
| replicate_unique), data = FB_df, dist = "lognormal")
FB_F_6 <- SurvregME(Surv(age_followup, FB_event) ~ lineage + tum_state_early + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_7 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition * lineage + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_8 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition * tum_state_early + (1
| replicate_unique), data = FB_df, dist = "lognormal")

```

```

FB_F_9 <- SurvregME(Surv(age_followup, FB_event) ~ lineage * tum_state_early + (1 |
replicate_unique), data = FB_df, dist = "lognormal")
FB_F_10 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition + lineage +
tum_state_early + (1 | replicate_unique), data = FB_df, dist = "lognormal")
FB_F_11 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition * lineage +
tum_state_early + (1 | replicate_unique), data = FB_df, dist = "lognormal")
FB_F_12 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition * tum_state_early +
lineage + (1 | replicate_unique), data = FB_df, dist = "lognormal")
FB_F_13 <- SurvregME(Surv(age_followup, FB_event) ~ lineage * tum_state_early +
sleep_condition + (1 | replicate_unique), data = FB_df, dist = "lognormal")
FB_F_14 <- SurvregME(Surv(age_followup, FB_event) ~ sleep_condition * lineage *
tum_state_early + (1 | replicate_unique), data = FB_df, dist = "lognormal")

AIC(FB_F_0, FB_F_1, FB_F_2, FB_F_3, FB_F_4, FB_F_5, FB_F_6, FB_F_7,
    FB_F_8, FB_F_9, FB_F_10, FB_F_11, FB_F_12, FB_F_13, FB_F_14) %>% arrange(AIC)

```

```

##      df      AIC
## FB_F_11 15 2022.674
## FB_F_12 11 2039.850
## FB_F_10 10 2040.381
## FB_F_14 26 2041.720
## FB_F_6   9 2044.891
## FB_F_13 15 2047.613
## FB_F_9   14 2051.542
## FB_F_7   14 2114.692
## FB_F_4    9 2123.404
## FB_F_2    8 2132.611
## FB_F_8    6 2207.268
## FB_F_5    5 2209.009
## FB_F_3    4 2214.100
## FB_F_1    4 2266.732
## FB_F_0    3 2275.753

```

The model with the interaction between the sleep condition and the lineage, and with the additive effect of tumoral state has the lowest AIC and is therefore selected (FB_F_11).


6.3 Goodness-of-fit evaluation

Assess goodness-of-fit of the selected model using Cox-Snell residuals.

```

plot_cox_snell_survregme(model = FB_F_11, data = FB_df, time_col = "age_followup",
event_col = "FB_event")

```



The curve follows a 45 degree exponential, we validate the fit of the selected model.

6.4 Selected model summary

#' Releveling of "lineage" to obtain all pairwise comparisons from the same model.

```
FB_df$lineage <- relevel(FB_df$lineage, ref = "HO_MT")
summary(SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
  data = FB_df, dist = "lognormal"))
```

```
##
## Mixed-Effects Log-Normal Linear Regression Model
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
## (1 | replicate_unique)
##
## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##
##
```

	Estimate	Std. Error
## sleep_conditionMechanical sleep disruption	0.048981	0.170051
## lineageHC_MT	2.211147	0.298500
## lineageHO_SPC	0.185987	0.257831
## lineageHO_SPT	1.474533	0.266992

```

## lineageHO_VLN                2.069535    0.252228
## lineageHV_GAL                1.971254    0.267067
## tum_state_earlyTumoral       2.234789    0.234979
## sleep_conditionMechanical sleep disruption:lineageHC_MT  0.578781    0.269384
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -0.036154    0.238204
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -0.516544    0.242203
## sleep_conditionMechanical sleep disruption:lineageHO_VLN  0.547600    0.225972
## sleep_conditionMechanical sleep disruption:lineageHV_GAL  0.212865    0.239548
##                               z value  Pr(>|z|)
## sleep_conditionMechanical sleep disruption            0.2880    0.77332
## lineageHC_MT                7.4075 1.287e-13 ***
## lineageHO_SPC                0.7214    0.47069
## lineageHO_SPT                5.5228 3.337e-08 ***
## lineageHO_VLN                8.2050 2.306e-16 ***
## lineageHV_GAL                7.3811 1.570e-13 ***
## tum_state_earlyTumoral       9.5106 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHC_MT  2.1485    0.03167 *
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -0.1518    0.87936
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -2.1327    0.03295 *
## sleep_conditionMechanical sleep disruption:lineageHO_VLN  2.4233    0.01538 *
## sleep_conditionMechanical sleep disruption:lineageHV_GAL  0.8886    0.37421
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)

```

```

FB_df$lineage <- relevel(FB_df$lineage, ref = "HO_SPC")
summary(SurvregME(Surv(age_followup, FB_event) ~
                  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
                  data = FB_df, dist = "lognormal"))

```

```

##
## Mixed-Effects Log-Normal Linear Regression Model
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
##   (1 | replicate_unique)
##
## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##

```

```

##                                     Estimate Std. Error
## sleep_conditionMechanical sleep disruption    0.012831    0.167869
## lineageHO_MT                                -0.185979    0.257831
## lineageHC_MT                                2.025164    0.296385
## lineageHO_SPT                               1.288545    0.265838
## lineageHO_VLN                               1.883548    0.249845
## lineageHV_GAL                               1.785271    0.264853
## tum_state_earlyTumoral                      2.234794    0.234979
## sleep_conditionMechanical sleep disruption:lineageHO_MT 0.036143    0.238204
## sleep_conditionMechanical sleep disruption:lineageHC_MT 0.614928    0.267964
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -0.480394    0.240519
## sleep_conditionMechanical sleep disruption:lineageHO_VLN 0.583752    0.224262
## sleep_conditionMechanical sleep disruption:lineageHV_GAL 0.249012    0.237980
##                                     z value  Pr(>|z|)
## sleep_conditionMechanical sleep disruption    0.0764    0.939071
## lineageHO_MT                                -0.7213    0.470712
## lineageHC_MT                                6.8329 8.322e-12 ***
## lineageHO_SPT                               4.8471 1.253e-06 ***
## lineageHO_VLN                               7.5389 4.741e-14 ***
## lineageHV_GAL                               6.7406 1.577e-11 ***
## tum_state_earlyTumoral                      9.5106 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHO_MT 0.1517    0.879400
## sleep_conditionMechanical sleep disruption:lineageHC_MT 2.2948    0.021744 *
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -1.9973    0.045790 *
## sleep_conditionMechanical sleep disruption:lineageHO_VLN 2.6030    0.009242 **
## sleep_conditionMechanical sleep disruption:lineageHV_GAL 1.0464    0.295396
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)

```

```

FB_df$lineage <- relevel(FB_df$lineage, ref = "HO_SPT")
summary(SurvregME(Surv(age_followup, FB_event) ~
                  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
                  data = FB_df, dist = "lognormal"))

```

```

##
## Mixed-Effects Log-Normal Linear Regression Model
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
##      (1 | replicate_unique)
##

```

```

## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##
##                                     Estimate Std. Error
## sleep_conditionMechanical sleep disruption -0.46757    0.17350
## lineageHO_SPC -1.28855    0.26584
## lineageHO_MT -1.47453    0.26699
## lineageHC_MT 0.73661    0.29238
## lineageHO_VLN 0.59499    0.24624
## lineageHV_GAL 0.49671    0.26189
## tum_state_earlyTumoral 2.23480    0.23498
## sleep_conditionMechanical sleep disruption:lineageHO_SPC 0.48040    0.24052
## sleep_conditionMechanical sleep disruption:lineageHO_MT 0.51654    0.24220
## sleep_conditionMechanical sleep disruption:lineageHC_MT 1.09533    0.27264
## sleep_conditionMechanical sleep disruption:lineageHO_VLN 1.06416    0.22974
## sleep_conditionMechanical sleep disruption:lineageHV_GAL 0.72942    0.24241
##                                     z value Pr(>|z|)
## sleep_conditionMechanical sleep disruption -2.6949 0.007041 **
## lineageHO_SPC -4.8471 1.253e-06 ***
## lineageHO_MT -5.5228 3.337e-08 ***
## lineageHC_MT 2.5194 0.011756 *
## lineageHO_VLN 2.4163 0.015678 *
## lineageHV_GAL 1.8966 0.057877 .
## tum_state_earlyTumoral 9.5106 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHO_SPC 1.9974 0.045787 *
## sleep_conditionMechanical sleep disruption:lineageHO_MT 2.1327 0.032950 *
## sleep_conditionMechanical sleep disruption:lineageHC_MT 4.0176 5.880e-05 ***
## sleep_conditionMechanical sleep disruption:lineageHO_VLN 4.6319 3.623e-06 ***
## sleep_conditionMechanical sleep disruption:lineageHV_GAL 3.0090 0.002621 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)

FB_df$lineage <- relevel(FB_df$lineage, ref = "HO_VLN")
summary(SurvregME(Surv(age_followup, FB_event) ~
                  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
                  data = FB_df, dist = "lognormal"))

##
## Mixed-Effects Log-Normal Linear Regression Model

```

```
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
##   (1 | replicate_unique)
##
## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##
##                                     Estimate Std. Error
## sleep_conditionMechanical sleep disruption    0.596595    0.149017
## lineageHO_SPT                                -0.594993    0.246240
## lineageHO_SPC                                -1.883540    0.249845
## lineageHO_MT                                 -2.069524    0.252228
## lineageHC_MT                                 0.141629    0.272716
## lineageHV_GAL                                -0.098264    0.240411
## tum_state_earlyTumoral                       2.234804    0.234979
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -1.064162    0.229745
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -0.583767    0.224262
## sleep_conditionMechanical sleep disruption:lineageHO_MT  -0.547622    0.225972
## sleep_conditionMechanical sleep disruption:lineageHC_MT   0.031167    0.255372
## sleep_conditionMechanical sleep disruption:lineageHV_GAL -0.334754    0.224607
##                                     z value  Pr(>|z|)
## sleep_conditionMechanical sleep disruption    4.0035 6.240e-05 ***
## lineageHO_SPT                                -2.4163  0.01568 *
## lineageHO_SPC                                -7.5388 4.742e-14 ***
## lineageHO_MT                                 -8.2050 2.307e-16 ***
## lineageHC_MT                                 0.5193  0.60353
## lineageHV_GAL                                -0.4087  0.68273
## tum_state_earlyTumoral                       9.5107 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -4.6319 3.623e-06 ***
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -2.6031  0.00924 **
## sleep_conditionMechanical sleep disruption:lineageHO_MT  -2.4234  0.01538 *
## sleep_conditionMechanical sleep disruption:lineageHC_MT   0.1220  0.90286
## sleep_conditionMechanical sleep disruption:lineageHV_GAL -1.4904  0.13612
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)
```

```
FB_df$lineage <- relevel(FB_df$lineage, ref = "HC_MT")
summary(SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
```

```
data = FB_df, dist = "lognormal"))
```

```
##
## Mixed-Effects Log-Normal Linear Regression Model
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
##   (1 | replicate_unique)
##
## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##
##                                     Estimate Std. Error
## sleep_conditionMechanical sleep disruption    0.627755    0.209080
## lineageHO_VLN                                -0.141603    0.272716
## lineageHO_SPT                                -0.736598    0.292376
## lineageHO_SPC                                -2.025149    0.296384
## lineageHO_MT                                 -2.211131    0.298500
## lineageHV_GAL                                -0.239892    0.287121
## tum_state_earlyTumoral                        2.234790    0.234979
## sleep_conditionMechanical sleep disruption:lineageHO_VLN -0.031172    0.255372
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -1.095320    0.272637
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -0.614927    0.267964
## sleep_conditionMechanical sleep disruption:lineageHO_MT  -0.578778    0.269384
## sleep_conditionMechanical sleep disruption:lineageHV_GAL -0.365902    0.268223
##                                     z value Pr(>|z|)
## sleep_conditionMechanical sleep disruption      3.0025  0.002678 **
## lineageHO_VLN                                -0.5192  0.603600
## lineageHO_SPT                                -2.5194  0.011757 *
## lineageHO_SPC                                -6.8329 8.324e-12 ***
## lineageHO_MT                                 -7.4075 1.287e-13 ***
## lineageHV_GAL                                -0.8355  0.403433
## tum_state_earlyTumoral                        9.5106 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHO_VLN -0.1221  0.902847
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -4.0175 5.882e-05 ***
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -2.2948  0.021744 *
## sleep_conditionMechanical sleep disruption:lineageHO_MT  -2.1485  0.031672 *
## sleep_conditionMechanical sleep disruption:lineageHV_GAL -1.3642  0.172514
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)
```

```

FB_df$lineage <- relevel(FB_df$lineage, ref = "HV_GAL")
summary(SurvregME(Surv(age_followup, FB_event) ~
  sleep_condition * lineage + tum_state_early + (1 | replicate_unique),
  data = FB_df, dist = "lognormal"))

```

```

##
## Mixed-Effects Log-Normal Linear Regression Model
##
## Formula:
## Surv(age_followup, FB_event) ~ sleep_condition * lineage + tum_state_early +
##   (1 | replicate_unique)
##
## Fitted to dataset FB_df
##
## Fixed effects parameters:
## =====
##
##                                     Estimate Std. Error
## sleep_conditionMechanical sleep disruption    0.261860   0.168784
## lineageHC_MT                                0.239910   0.287122
## lineageHO_VLN                               0.098294   0.240411
## lineageHO_SPT                              -0.496709   0.261894
## lineageHO_SPC                              -1.785257   0.264853
## lineageHO_MT                               -1.971238   0.267067
## tum_state_earlyTumoral                      2.234796   0.234979
## sleep_conditionMechanical sleep disruption:lineageHC_MT  0.365898   0.268223
## sleep_conditionMechanical sleep disruption:lineageHO_VLN  0.334720   0.224607
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -0.729426   0.242415
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -0.249033   0.237980
## sleep_conditionMechanical sleep disruption:lineageHO_MT -0.212888   0.239548
##                                     z value  Pr(>|z|)
## sleep_conditionMechanical sleep disruption    1.5515  0.120793
## lineageHC_MT                                0.8356  0.403398
## lineageHO_VLN                               0.4089  0.682642
## lineageHO_SPT                              -1.8966  0.057880 .
## lineageHO_SPC                              -6.7406 1.578e-11 ***
## lineageHO_MT                               -7.3811 1.570e-13 ***
## tum_state_earlyTumoral                      9.5106 < 2.2e-16 ***
## sleep_conditionMechanical sleep disruption:lineageHC_MT  1.3642  0.172519
## sleep_conditionMechanical sleep disruption:lineageHO_VLN  1.4902  0.136159
## sleep_conditionMechanical sleep disruption:lineageHO_SPT -3.0090  0.002621 **
## sleep_conditionMechanical sleep disruption:lineageHO_SPC -1.0464  0.295356
## sleep_conditionMechanical sleep disruption:lineageHO_MT -0.8887  0.374160
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Random effects:
## =====
##
## Grouping factor: replicate_unique (432 levels)
## Standard deviation:
## (Intercept)

```

```
##      1.1703
##
##
## Log-likelihood: -996.3371 (npar = 15)
```

7 Median times

7.1 By sleep condition

Estimate cumulative incidence functions for the first bud event, accounting for the competing risk of death, stratified by sleep condition.

```
FB_cuminc_sleep <- cuminc(Surv(age_followup, status_competing) ~ sleep_condition, data =
FB_df)
```

For each stratum, extract the median time to event and its 95CI.

```
FB_medians_sleep <- tidy(FB_cuminc_sleep) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::group_by(strata) %>%
  dplyr::arrange(time) %>%
  dplyr::summarise(
    median_time = { idx <- which(estimate >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_low = { idx <- which(conf.high >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_high = { idx <- which(conf.low >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    .groups = "drop")

print(FB_medians_sleep)
```

```
## # A tibble: 2 x 4
##   strata                median_time ci_low ci_high
##   <fct>                <dbl>   <dbl>   <dbl>
## 1 Control                1.14     1.14     1.14
## 2 Mechanical sleep disruption 1.14     1.14     1.29
```

7.2 By lineage

Same method as above, but stratified by lineage.

```
FB_cuminc_lineage <- cuminc(Surv(age_followup, status_competing) ~ lineage, data = FB_df)

FB_medians_lineage <- tidy(FB_cuminc_lineage) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::group_by(strata) %>%
  dplyr::arrange(time) %>%
  dplyr::summarise(
    median_time = { idx <- which(estimate >= 0.5)[1]
```

```

    if (is.na(idx)) NA_real_ else time[idx] },
    ci_low = { idx <- which(conf.high >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_high = { idx <- which(conf.low >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    .groups = "drop")

print(FB_medians_lineage)

```

```

## # A tibble: 6 x 4
##   strata median_time ci_low ci_high
##   <fct>      <dbl> <dbl> <dbl>
## 1 HV_GAL      1.71  1.29    2
## 2 HC_MT        3    2.43    3
## 3 HO_VLN       1     1    2.43
## 4 HO_SPT      1.29  1.14    1.29
## 5 HO_SPC      0.286 0.286    1.14
## 6 HO_MT       0.286 0.286    1

```

7.3 By sleep condition and lineage

Same method as above, but stratified by sleep condition and lineage.

```

FB_cuminc_interaction <- cuminc(Surv(age_followup, status_competing) ~ sleep_condition +
lineage, data = FB_df)

FB_medians_interaction <- tidy(FB_cuminc_interaction) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::group_by(strata) %>%
  dplyr::arrange(time) %>%
  dplyr::summarise(
    median_time = { idx <- which(estimate >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_low = { idx <- which(conf.high >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_high = { idx <- which(conf.low >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    .groups = "drop")

print(FB_medians_interaction)

```

```

## # A tibble: 12 x 4
##   strata                                median_time ci_low ci_high
##   <fct>                                <dbl> <dbl> <dbl>
## 1 Control, HV_GAL                      1.71  1.71    2
## 2 Mechanical sleep disruption, HV_GAL  1.71  1    2.43
## 3 Control, HC_MT                      1.43  1.14    3
## 4 Mechanical sleep disruption, HC_MT    3     3     3
## 5 Control, HO_VLN                      1     1    1.14
## 6 Mechanical sleep disruption, HO_VLN  1.14  1    3.14
## 7 Control, HO_SPT                     1.29  1.14    1.57
## 8 Mechanical sleep disruption, HO_SPT  1.14  1.14    1.29

```

```
## 9 Control, HO_SPC 0.286 0.286 1.14
## 10 Mechanical sleep disruption, HO_SPC 0.286 0.286 1.14
## 11 Control, HO_MT 0.286 0.286 1
## 12 Mechanical sleep disruption, HO_MT 0.286 0.286 1
```

7.4 By tumoral state

Same method as above, but stratified by tumoral state.

```
FB_cuminc_tum <- cuminc(Surv(age_followup, status_competing) ~ tum_state_early, data =
FB_df)

FB_medians_tum <- tidy(FB_cuminc_tum) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::group_by(strata) %>%
  dplyr::arrange(time) %>%
  dplyr::summarise(
    median_time = { idx <- which(estimate >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_low = { idx <- which(conf.high >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    ci_high = { idx <- which(conf.low >= 0.5)[1]
    if (is.na(idx)) NA_real_ else time[idx] },
    .groups = "drop")

print(FB_medians_tum)
```

```
## # A tibble: 2 x 4
##   strata median_time ci_low ci_high
##   <fct>      <dbl> <dbl> <dbl>
## 1 Healthy      1.14   1.14   1.14
## 2 Tumoral       3     2.29   NA
```

8 Cumulative incidences

8.1 By week and sleep condition

Convert the continuous cumulative incidence function into a weekly summary table: for each stratum of sleep condition and each week, show the estimate with its 95CI.

```
FB_incidence_sleep <- tidy(FB_cuminc_sleep) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::mutate(week = ceiling(time)) %>%
  dplyr::group_by(strata, week) %>%
  dplyr::filter(time == max(time)) %>%
  dplyr::slice(1) %>%
  dplyr::ungroup() %>%
  dplyr::select(strata, week, estimate_cuminc = estimate, conf.low, conf.high) %>%
  dplyr::arrange(strata, week)

print(FB_incidence_sleep)
```

```
## # A tibble: 14 x 5
##   strata                week estimate_cuminc conf.low conf.high
##   <fct>                <dbl>         <dbl>    <dbl>    <dbl>
## 1 Control                0             0      NA      NA
## 2 Control                1          0.433    0.386    0.479
## 3 Control                2          0.736    0.692    0.775
## 4 Control                3          0.894    0.860    0.919
## 5 Control                4          0.907    0.876    0.931
## 6 Control                5          0.931    0.902    0.951
## 7 Control                6          0.947    0.921    0.964
## 8 Mechanical sleep disruption 0             0      NA      NA
## 9 Mechanical sleep disruption 1          0.421    0.374    0.467
## 10 Mechanical sleep disruption 2          0.650    0.603    0.693
## 11 Mechanical sleep disruption 3          0.789    0.748    0.825
## 12 Mechanical sleep disruption 4          0.822    0.782    0.855
## 13 Mechanical sleep disruption 5          0.880    0.845    0.907
## 14 Mechanical sleep disruption 6          0.882    0.848    0.909
```

8.2 By week and lineage

Same method as above, but stratified by lineage.

```
FB_incidence_lineage <- tidy(FB_cuminc_lineage) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::mutate(week = ceiling(time)) %>%
  dplyr::group_by(strata, week) %>%
  dplyr::filter(time == max(time)) %>%
  dplyr::slice(1) %>%
  dplyr::ungroup() %>%
  dplyr::select(strata, week, estimate_cuminc = estimate, conf.low, conf.high) %>%
  dplyr::arrange(strata, week)

print(FB_incidence_lineage)
```

```
## # A tibble: 42 x 5
##   strata week estimate_cuminc conf.low conf.high
##   <fct> <dbl>         <dbl>    <dbl>    <dbl>
## 1 HV_GAL 0             0      NA      NA
## 2 HV_GAL 1          0.319    0.245    0.396
## 3 HV_GAL 2          0.597    0.512    0.672
## 4 HV_GAL 3          0.868    0.800    0.914
## 5 HV_GAL 4          0.875    0.808    0.920
## 6 HV_GAL 5          0.931    0.874    0.962
## 7 HV_GAL 6          0.938    0.882    0.967
## 8 HC_MT 0             0      NA      NA
## 9 HC_MT 1          0.0729    0.0320    0.137
## 10 HC_MT 2          0.375    0.279    0.471
## 11 HC_MT 3          0.844    0.754    0.903
## 12 HC_MT 4          0.917    0.838    0.958
## 13 HC_MT 5          0.938    0.862    0.972
## 14 HC_MT 6          0.938    0.862    0.972
## 15 HO_VLN 0             0      NA      NA
## 16 HO_VLN 1          0.536    0.463    0.604
```

```
## 17 HO_VLN      2      0.573    0.500    0.640
## 18 HO_VLN      3      0.672    0.600    0.734
## 19 HO_VLN      4      0.703    0.633    0.763
## 20 HO_VLN      5      0.813    0.749    0.861
## 21 HO_VLN      6      0.849    0.789    0.893
## 22 HO_SPT      0      0        NA      NA
## 23 HO_SPT      1      0.215    0.152    0.286
## 24 HO_SPT      2      0.819    0.746    0.874
## 25 HO_SPT      3      0.868    0.800    0.914
## 26 HO_SPT      4      0.882    0.816    0.925
## 27 HO_SPT      5      0.882    0.816    0.925
## 28 HO_SPT      6      0.882    0.816    0.925
## 29 HO_SPC      0      0        NA      NA
## 30 HO_SPC      1      0.549    0.463    0.626
## 31 HO_SPC      2      0.882    0.816    0.925
## 32 HO_SPC      3      0.924    0.865    0.957
## 33 HO_SPC      4      0.951    0.899    0.977
## 34 HO_SPC      5      0.979    0.931    0.994
## 35 HO_SPC      6      0.979    0.931    0.994
## 36 HO_MT       0      0        NA      NA
## 37 HO_MT       1      0.715    0.634    0.781
## 38 HO_MT       2      0.847    0.778    0.896
## 39 HO_MT       3      0.931    0.874    0.962
## 40 HO_MT       4      0.931    0.874    0.962
## 41 HO_MT       5      0.931    0.874    0.962
## 42 HO_MT       6      0.931    0.874    0.962
```

8.3 By week, sleep condition, and lineage

Same method as above, but stratified by sleep condition and lineage.

```
FB_incidence_interaction <- tidy(FB_cuminc_interaction) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::mutate(week = ceiling(time)) %>%
  dplyr::group_by(strata, week) %>%
  dplyr::filter(time == max(time)) %>%
  dplyr::slice(1) %>%
  dplyr::ungroup() %>%
  dplyr::select(strata, week, estimate_cuminc = estimate, conf.low, conf.high) %>%
  dplyr::arrange(strata, week)

print(FB_incidence_interaction)
```

```
## # A tibble: 84 x 5
##   strata                week estimate_cuminc conf.low conf.high
##   <fct>                <dbl>         <dbl>    <dbl>    <dbl>
## 1 Control, HV_GAL      0           0        NA        NA
## 2 Control, HV_GAL      1         0.222    0.134    0.324
## 3 Control, HV_GAL      2         0.639    0.515    0.739
## 4 Control, HV_GAL      3         0.986    0.898    0.998
## 5 Control, HV_GAL      4          NA        NA        NA
## 6 Control, HV_GAL      5          NA        NA        NA
## 7 Control, HV_GAL      6          NA        NA        NA
```

## 8	Mechanical sleep disruption, HV_GAL	0	0	NA	NA
## 9	Mechanical sleep disruption, HV_GAL	1	0.417	0.302	0.528
## 10	Mechanical sleep disruption, HV_GAL	2	0.556	0.432	0.662
## 11	Mechanical sleep disruption, HV_GAL	3	0.75	0.631	0.835
## 12	Mechanical sleep disruption, HV_GAL	4	0.75	0.631	0.835
## 13	Mechanical sleep disruption, HV_GAL	5	0.861	0.755	0.924
## 14	Mechanical sleep disruption, HV_GAL	6	0.875	0.771	0.934
## 15	Control, HC_MT	0	0	NA	NA
## 16	Control, HC_MT	1	0.0833	0.0263	0.183
## 17	Control, HC_MT	2	0.521	0.370	0.652
## 18	Control, HC_MT	3	0.917	0.788	0.969
## 19	Control, HC_MT	4	0.979	0.777	0.998
## 20	Control, HC_MT	5	0.979	0.777	0.998
## 21	Control, HC_MT	6	0.979	0.777	0.998
## 22	Mechanical sleep disruption, HC_MT	0	0	NA	NA
## 23	Mechanical sleep disruption, HC_MT	1	0.0625	0.0160	0.156
## 24	Mechanical sleep disruption, HC_MT	2	0.229	0.122	0.356
## 25	Mechanical sleep disruption, HC_MT	3	0.771	0.623	0.867
## 26	Mechanical sleep disruption, HC_MT	4	0.854	0.713	0.929
## 27	Mechanical sleep disruption, HC_MT	5	0.896	0.759	0.957
## 28	Mechanical sleep disruption, HC_MT	6	0.896	0.759	0.957
## 29	Control, HO_VLN	0	0	NA	NA
## 30	Control, HO_VLN	1	0.604	0.499	0.694
## 31	Control, HO_VLN	2	0.635	0.530	0.723
## 32	Control, HO_VLN	3	0.75	0.650	0.825
## 33	Control, HO_VLN	4	0.771	0.672	0.843
## 34	Control, HO_VLN	5	0.865	0.776	0.920
## 35	Control, HO_VLN	6	0.938	0.863	0.972
## 36	Mechanical sleep disruption, HO_VLN	0	0	NA	NA
## 37	Mechanical sleep disruption, HO_VLN	1	0.469	0.366	0.565
## 38	Mechanical sleep disruption, HO_VLN	2	0.510	0.406	0.606
## 39	Mechanical sleep disruption, HO_VLN	3	0.594	0.488	0.685
## 40	Mechanical sleep disruption, HO_VLN	4	0.635	0.530	0.723
## 41	Mechanical sleep disruption, HO_VLN	5	0.760	0.661	0.835
## 42	Mechanical sleep disruption, HO_VLN	6	0.760	0.661	0.835
## 43	Control, HO_SPT	0	0	NA	NA
## 44	Control, HO_SPT	1	0.194	0.112	0.293
## 45	Control, HO_SPT	2	0.819	0.707	0.892
## 46	Control, HO_SPT	3	0.847	0.738	0.914
## 47	Control, HO_SPT	4	0.847	0.738	0.914
## 48	Control, HO_SPT	5	0.847	0.738	0.914
## 49	Control, HO_SPT	6	0.847	0.738	0.914
## 50	Mechanical sleep disruption, HO_SPT	0	0	NA	NA
## 51	Mechanical sleep disruption, HO_SPT	1	0.236	0.145	0.340
## 52	Mechanical sleep disruption, HO_SPT	2	0.819	0.707	0.892
## 53	Mechanical sleep disruption, HO_SPT	3	0.889	0.786	0.944
## 54	Mechanical sleep disruption, HO_SPT	4	0.917	0.818	0.963
## 55	Mechanical sleep disruption, HO_SPT	5	0.917	0.818	0.963
## 56	Mechanical sleep disruption, HO_SPT	6	0.917	0.818	0.963
## 57	Control, HO_SPC	0	0	NA	NA
## 58	Control, HO_SPC	1	0.542	0.419	0.649
## 59	Control, HO_SPC	2	0.889	0.787	0.944
## 60	Control, HO_SPC	3	0.972	0.870	0.994
## 61	Control, HO_SPC	4	0.972	0.870	0.994

## 62 Control, HO_SPC	5	NA	NA	NA
## 63 Control, HO_SPC	6	NA	NA	NA
## 64 Mechanical sleep disruption, HO_SPC	0	0	NA	NA
## 65 Mechanical sleep disruption, HO_SPC	1	0.556	0.433	0.662
## 66 Mechanical sleep disruption, HO_SPC	2	0.875	0.771	0.934
## 67 Mechanical sleep disruption, HO_SPC	3	0.875	0.771	0.934
## 68 Mechanical sleep disruption, HO_SPC	4	0.931	0.836	0.972
## 69 Mechanical sleep disruption, HO_SPC	5	0.972	0.874	0.994
## 70 Mechanical sleep disruption, HO_SPC	6	0.972	0.874	0.994
## 71 Control, HO_MT	0	0	NA	NA
## 72 Control, HO_MT	1	0.778	0.663	0.858
## 73 Control, HO_MT	2	0.875	0.771	0.934
## 74 Control, HO_MT	3	0.944	0.852	0.980
## 75 Control, HO_MT	4	0.944	0.852	0.980
## 76 Control, HO_MT	5	0.944	0.852	0.980
## 77 Control, HO_MT	6	0.944	0.852	0.980
## 78 Mechanical sleep disruption, HO_MT	0	0	NA	NA
## 79 Mechanical sleep disruption, HO_MT	1	0.653	0.530	0.751
## 80 Mechanical sleep disruption, HO_MT	2	0.819	0.709	0.891
## 81 Mechanical sleep disruption, HO_MT	3	0.917	0.821	0.962
## 82 Mechanical sleep disruption, HO_MT	4	0.917	0.821	0.962
## 83 Mechanical sleep disruption, HO_MT	5	0.917	0.821	0.962
## 84 Mechanical sleep disruption, HO_MT	6	0.917	0.821	0.962

8.4 By week and tumoral state

Same method as above, but stratified by tumoral state.

```
FB_incidence_tum <- tidy(FB_cuminc_tum) %>%
  dplyr::filter(outcome == "1") %>%
  dplyr::mutate(week = ceiling(time)) %>%
  dplyr::group_by(strata, week) %>%
  dplyr::filter(time == max(time)) %>%
  dplyr::slice(1) %>%
  dplyr::ungroup() %>%
  dplyr::select(strata, week, estimate_cuminc = estimate, conf.low, conf.high) %>%
  dplyr::arrange(strata, week)

print(FB_incidence_tum)
```

## # A tibble: 14 x 5					
## strata week estimate_cuminc conf.low conf.high					
## <fct> <dbl> <dbl> <dbl> <dbl>					
## 1 Healthy 0 0 NA NA					
## 2 Healthy 1 0.447 0.413 0.481					
## 3 Healthy 2 0.712 0.680 0.742					
## 4 Healthy 3 0.858 0.832 0.880					
## 5 Healthy 4 0.880 0.856 0.900					
## 6 Healthy 5 0.919 0.898 0.935					
## 7 Healthy 6 0.928 0.908 0.944					
## 8 Tumoral 0 0 NA NA					
## 9 Tumoral 1 0.0244 0.00185 0.112					
## 10 Tumoral 2 0.317 0.181 0.462					

```
## 11 Tumoral      3      0.512  0.349      0.654
## 12 Tumoral      4      0.561  0.394      0.699
## 13 Tumoral      5      0.634  0.464      0.763
## 14 Tumoral      6      0.634  0.464      0.763
```

9 Cumulative proportions

9.1 By week and sleep condition

Compute raw cumulative proportions of first bud production per week, stratified by sleep condition.

```
FB_proportions_sleep <- FB_df %>%
  dplyr::mutate(week_FB_event = ceiling(age_followup)) %>%
  dplyr::group_by(sleep_condition) %>%
  dplyr::summarise(total_N = n(),
    n1 = sum(week_FB_event <= 1 & FB_event == 1),
    n2 = sum(week_FB_event <= 2 & FB_event == 1),
    n3 = sum(week_FB_event <= 3 & FB_event == 1),
    n4 = sum(week_FB_event <= 4 & FB_event == 1),
    n5 = sum(week_FB_event <= 5 & FB_event == 1),
    n6 = sum(week_FB_event <= 6 & FB_event == 1),
    p1 = n1 / total_N,
    p2 = n2 / total_N,
    p3 = n3 / total_N,
    p4 = n4 / total_N,
    p5 = n5 / total_N,
    p6 = n6 / total_N,
    .groups = "drop") %>%
  tidyr::pivot_longer(cols = matches("[np][1-6]$"),
    names_to = c(".value", "week"),
    names_pattern = "(n|p)([1-6])") %>%
  dplyr::mutate(week = as.numeric(week)) %>%
  dplyr::select(sleep_condition, week, n, total_N, raw_proportion = p)

print(FB_proportions_sleep)
```

```
## # A tibble: 12 x 5
##   sleep_condition      week      n total_N raw_proportion
##   <fct>          <dbl> <int>   <int>         <dbl>
## 1 Control            1    187    432         0.433
## 2 Control            2    318    432         0.736
## 3 Control            3    386    432         0.894
## 4 Control            4    392    432         0.907
## 5 Control            5    402    432         0.931
## 6 Control            6    409    432         0.947
## 7 Mechanical sleep disruption 1    182    432         0.421
## 8 Mechanical sleep disruption 2    281    432         0.650
## 9 Mechanical sleep disruption 3    341    432         0.789
## 10 Mechanical sleep disruption 4    355    432         0.822
## 11 Mechanical sleep disruption 5    380    432         0.880
## 12 Mechanical sleep disruption 6    381    432         0.882
```

9.2 By week and lineage

Same method as above, but stratified by lineage.

```
FB_proportions_lineage <- FB_df %>%
  dplyr::mutate(week_FB_event = ceiling(age_followup)) %>%
  dplyr::group_by(lineage) %>%
  dplyr::summarise(total_N = n(),
    n1 = sum(week_FB_event <= 1 & FB_event == 1),
    n2 = sum(week_FB_event <= 2 & FB_event == 1),
    n3 = sum(week_FB_event <= 3 & FB_event == 1),
    n4 = sum(week_FB_event <= 4 & FB_event == 1),
    n5 = sum(week_FB_event <= 5 & FB_event == 1),
    n6 = sum(week_FB_event <= 6 & FB_event == 1),
    p1 = n1 / total_N,
    p2 = n2 / total_N,
    p3 = n3 / total_N,
    p4 = n4 / total_N,
    p5 = n5 / total_N,
    p6 = n6 / total_N,
    .groups = "drop") %>%
  tidyr::pivot_longer(cols = matches("^np[1-6]$",),
    names_to = c(".value", "week"),
    names_pattern = "(n|p)([1-6])") %>%
  dplyr::mutate(week = as.numeric(week)) %>%
  dplyr::select(lineage, week, n, total_N, raw_proportion = p)

print(FB_proportions_lineage)
```

```
## # A tibble: 36 x 5
##   lineage week   n total_N raw_proportion
##   <fct>   <dbl> <int>   <int>         <dbl>
## 1 HV_GAL     1    46    144         0.319
## 2 HV_GAL     2    86    144         0.597
## 3 HV_GAL     3   125    144         0.868
## 4 HV_GAL     4   126    144         0.875
## 5 HV_GAL     5   134    144         0.931
## 6 HV_GAL     6   135    144         0.938
## 7 HC_MT      1     7     96         0.0729
## 8 HC_MT      2    36     96         0.375
## 9 HC_MT      3    81     96         0.844
## 10 HC_MT     4    88     96         0.917
## 11 HC_MT     5    90     96         0.938
## 12 HC_MT     6    90     96         0.938
## 13 HO_VLN     1   103    192         0.536
## 14 HO_VLN     2   110    192         0.573
## 15 HO_VLN     3   129    192         0.672
## 16 HO_VLN     4   135    192         0.703
## 17 HO_VLN     5   156    192         0.812
## 18 HO_VLN     6   163    192         0.849
## 19 HO_SPT     1    31    144         0.215
## 20 HO_SPT     2   118    144         0.819
## 21 HO_SPT     3   125    144         0.868
## 22 HO_SPT     4   127    144         0.882
```

```
## 23 HO_SPT      5   127   144      0.882
## 24 HO_SPT      6   127   144      0.882
## 25 HO_SPC      1    79   144      0.549
## 26 HO_SPC      2   127   144      0.882
## 27 HO_SPC      3   133   144      0.924
## 28 HO_SPC      4   137   144      0.951
## 29 HO_SPC      5   141   144      0.979
## 30 HO_SPC      6   141   144      0.979
## 31 HO_MT       1   103   144      0.715
## 32 HO_MT       2   122   144      0.847
## 33 HO_MT       3   134   144      0.931
## 34 HO_MT       4   134   144      0.931
## 35 HO_MT       5   134   144      0.931
## 36 HO_MT       6   134   144      0.931
```

9.3 By week, sleep condition, and lineage

Same method as above, but stratified by sleep condition and lineage.

```
FB_proportions_interaction <- FB_df %>%
  dplyr::mutate(week_FB_event = ceiling(age_followup)) %>%
  dplyr::group_by(sleep_condition, lineage) %>%
  dplyr::summarise(total_N = n(),
    n1 = sum(week_FB_event <= 1 & FB_event == 1),
    n2 = sum(week_FB_event <= 2 & FB_event == 1),
    n3 = sum(week_FB_event <= 3 & FB_event == 1),
    n4 = sum(week_FB_event <= 4 & FB_event == 1),
    n5 = sum(week_FB_event <= 5 & FB_event == 1),
    n6 = sum(week_FB_event <= 6 & FB_event == 1),
    p1 = n1 / total_N,
    p2 = n2 / total_N,
    p3 = n3 / total_N,
    p4 = n4 / total_N,
    p5 = n5 / total_N,
    p6 = n6 / total_N,
    .groups = "drop") %>%
  tidyr::pivot_longer(cols = matches("^np[1-6]$",
    names_to = c(".value", "week"),
    names_pattern = "(n|p)([1-6])") %>%
  dplyr::mutate(week = as.numeric(week)) %>%
  dplyr::select(sleep_condition, lineage, week, n, total_N, raw_proportion = p)

print(FB_proportions_interaction)
```

```
## # A tibble: 72 x 6
##   sleep_condition lineage week    n total_N raw_proportion
##   <fct>          <fct> <dbl> <int>   <int>         <dbl>
## 1 Control      HV_GAL     1    16     72         0.222
## 2 Control      HV_GAL     2    46     72         0.639
## 3 Control      HV_GAL     3    71     72         0.986
## 4 Control      HV_GAL     4    72     72          1
## 5 Control      HV_GAL     5    72     72          1
## 6 Control      HV_GAL     6    72     72          1
```

## 7 Control	HC_MT	1	4	48	0.0833
## 8 Control	HC_MT	2	25	48	0.521
## 9 Control	HC_MT	3	44	48	0.917
## 10 Control	HC_MT	4	47	48	0.979
## 11 Control	HC_MT	5	47	48	0.979
## 12 Control	HC_MT	6	47	48	0.979
## 13 Control	HO_VLN	1	58	96	0.604
## 14 Control	HO_VLN	2	61	96	0.635
## 15 Control	HO_VLN	3	72	96	0.75
## 16 Control	HO_VLN	4	74	96	0.771
## 17 Control	HO_VLN	5	83	96	0.865
## 18 Control	HO_VLN	6	90	96	0.938
## 19 Control	HO_SPT	1	14	72	0.194
## 20 Control	HO_SPT	2	59	72	0.819
## 21 Control	HO_SPT	3	61	72	0.847
## 22 Control	HO_SPT	4	61	72	0.847
## 23 Control	HO_SPT	5	61	72	0.847
## 24 Control	HO_SPT	6	61	72	0.847
## 25 Control	HO_SPC	1	39	72	0.542
## 26 Control	HO_SPC	2	64	72	0.889
## 27 Control	HO_SPC	3	70	72	0.972
## 28 Control	HO_SPC	4	70	72	0.972
## 29 Control	HO_SPC	5	71	72	0.986
## 30 Control	HO_SPC	6	71	72	0.986
## 31 Control	HO_MT	1	56	72	0.778
## 32 Control	HO_MT	2	63	72	0.875
## 33 Control	HO_MT	3	68	72	0.944
## 34 Control	HO_MT	4	68	72	0.944
## 35 Control	HO_MT	5	68	72	0.944
## 36 Control	HO_MT	6	68	72	0.944
## 37 Mechanical sleep disruption	HV_GAL	1	30	72	0.417
## 38 Mechanical sleep disruption	HV_GAL	2	40	72	0.556
## 39 Mechanical sleep disruption	HV_GAL	3	54	72	0.75
## 40 Mechanical sleep disruption	HV_GAL	4	54	72	0.75
## 41 Mechanical sleep disruption	HV_GAL	5	62	72	0.861
## 42 Mechanical sleep disruption	HV_GAL	6	63	72	0.875
## 43 Mechanical sleep disruption	HC_MT	1	3	48	0.0625
## 44 Mechanical sleep disruption	HC_MT	2	11	48	0.229
## 45 Mechanical sleep disruption	HC_MT	3	37	48	0.771
## 46 Mechanical sleep disruption	HC_MT	4	41	48	0.854
## 47 Mechanical sleep disruption	HC_MT	5	43	48	0.896
## 48 Mechanical sleep disruption	HC_MT	6	43	48	0.896
## 49 Mechanical sleep disruption	HO_VLN	1	45	96	0.469
## 50 Mechanical sleep disruption	HO_VLN	2	49	96	0.510
## 51 Mechanical sleep disruption	HO_VLN	3	57	96	0.594
## 52 Mechanical sleep disruption	HO_VLN	4	61	96	0.635
## 53 Mechanical sleep disruption	HO_VLN	5	73	96	0.760
## 54 Mechanical sleep disruption	HO_VLN	6	73	96	0.760
## 55 Mechanical sleep disruption	HO_SPT	1	17	72	0.236
## 56 Mechanical sleep disruption	HO_SPT	2	59	72	0.819
## 57 Mechanical sleep disruption	HO_SPT	3	64	72	0.889
## 58 Mechanical sleep disruption	HO_SPT	4	66	72	0.917
## 59 Mechanical sleep disruption	HO_SPT	5	66	72	0.917
## 60 Mechanical sleep disruption	HO_SPT	6	66	72	0.917

```
## 61 Mechanical sleep disruption HO_SPC      1    40    72    0.556
## 62 Mechanical sleep disruption HO_SPC      2    63    72    0.875
## 63 Mechanical sleep disruption HO_SPC      3    63    72    0.875
## 64 Mechanical sleep disruption HO_SPC      4    67    72    0.931
## 65 Mechanical sleep disruption HO_SPC      5    70    72    0.972
## 66 Mechanical sleep disruption HO_SPC      6    70    72    0.972
## 67 Mechanical sleep disruption HO_MT       1    47    72    0.653
## 68 Mechanical sleep disruption HO_MT       2    59    72    0.819
## 69 Mechanical sleep disruption HO_MT       3    66    72    0.917
## 70 Mechanical sleep disruption HO_MT       4    66    72    0.917
## 71 Mechanical sleep disruption HO_MT       5    66    72    0.917
## 72 Mechanical sleep disruption HO_MT       6    66    72    0.917
```

9.4 By week and tumoral state

Same method as above, but stratified by tumoral state.

```
FB_proportions_tum <- FB_df %>%
  dplyr::mutate(week_FB_event = ceiling(age_followup)) %>%
  dplyr::group_by(tum_state_early) %>%
  dplyr::summarise(total_N = n(),
    n1 = sum(week_FB_event <= 1 & FB_event == 1),
    n2 = sum(week_FB_event <= 2 & FB_event == 1),
    n3 = sum(week_FB_event <= 3 & FB_event == 1),
    n4 = sum(week_FB_event <= 4 & FB_event == 1),
    n5 = sum(week_FB_event <= 5 & FB_event == 1),
    n6 = sum(week_FB_event <= 6 & FB_event == 1),
    p1 = n1 / total_N,
    p2 = n2 / total_N,
    p3 = n3 / total_N,
    p4 = n4 / total_N,
    p5 = n5 / total_N,
    p6 = n6 / total_N,
    .groups = "drop") %>%
  tidyr::pivot_longer(cols = matches("^np[1-6]$",
    names_to = c(".value", "week"),
    names_pattern = "(n|p)([1-6])") %>%
  dplyr::mutate(week = as.numeric(week)) %>%
  dplyr::select(tum_state_early, week, n, total_N, raw_proportion = p)

print(FB_proportions_tum)
```

```
## # A tibble: 12 x 5
##   tum_state_early week      n total_N raw_proportion
##   <fct>          <dbl> <int>   <int>         <dbl>
## 1 Healthy        1    368    823         0.447
## 2 Healthy        2    586    823         0.712
## 3 Healthy        3    706    823         0.858
## 4 Healthy        4    724    823         0.880
## 5 Healthy        5    756    823         0.919
## 6 Healthy        6    764    823         0.928
## 7 Tumoral        1      1     41         0.0244
## 8 Tumoral        2     13     41         0.317
```

## 9 Tumoral	3	21	41	0.512
## 10 Tumoral	4	23	41	0.561
## 11 Tumoral	5	26	41	0.634
## 12 Tumoral	6	26	41	0.634

10 Plots of cumulative incidence of first bud production

10.1 Common themes

Theme of y-axis, and its presence or absence on the plots

```
scale_y <- scale_y_continuous(limits = c(0, 1), breaks = seq(0, 1, by = 0.25),
                              labels = percent, position = "right")
```

Fix to hide y-axis numbers, ticks, and titles on the right and left

```
hide_y <- theme(axis.text.y.right = element_blank(),
                axis.ticks.y.right = element_blank(),
                axis.title.y.right = element_blank(),
                axis.text.y = element_blank(),
                axis.ticks.y = element_blank(),
                axis.title.y = element_blank())
```

Fix to show y-axis numbers and ticks on the right

```
show_y <- theme(axis.text.y.right = element_text(color = "black", size = 14),
                axis.ticks.y.right = element_line(),
                axis.text.y = element_blank(),
                axis.ticks.y = element_blank(),
                axis.title.y = element_blank())
```

Theme of x-axis, and its presence or absence on the plots.

```
scale_x <- scale_x_continuous(limits = c(0, 6), breaks = seq(0, 6, by = 2))

hide_x <- theme(axis.text.x = element_blank(),
                axis.ticks.x = element_blank(),
                axis.title.x = element_blank())

show_x <- theme(axis.text.x = element_text(color = "black", size = 14),
                axis.ticks.x = element_line())
```

Color scale for tumoral state

```
tum_scale <- list(
  scale_color_manual(values = tum_colors),
  scale_fill_manual(values = tum_colors))
```

Color scales for sleep condition

```
sleep_scale <- list(
  scale_color_manual(values = sleep_condition_colors, breaks = c("Control", "Mechanical
sleep disruption")),
  scale_fill_manual(values = sleep_condition_colors, breaks = c("Control", "Mechanical
sleep disruption")))
```

Theme shared by panels A, B, and C (angle = 90 for bottom-to-top reading)

```
theme_common <- theme(
  axis.title.x = element_text(color = "black", size = 14, margin = margin(t = 10)),
  axis.title.y.right = element_text(color = "black", size = 14, angle = 90, margin =
margin(l = 15)),
  plot.margin = margin(l = 5, r = 5, t = 5, b = 5))
```

Specific label configurations for precise placement

```
labs_base <- labs(x = NULL, y = NULL, title = NULL)
labs_x_only <- labs(x = "Time (weeks)", y = NULL, title = NULL)
labs_y_only <- labs(x = NULL, y = "Cumulative incidence of\nfirst bud production", title
= NULL)
labs_xy <- labs(x = "Time (weeks)", y = "Cumulative incidence of\nfirst bud
production", title = NULL)
```

10.2 Panel A: interaction between sleep condition and lineage

10.2.1 HO_SPT

```
A1 <- cuminc(Surv(age_followup, factor(status_competing)) ~ sleep_condition,
  data = FB_df %>% filter(lineage == "HO_SPT") %>%
  mutate(sleep_condition = factor(sleep_condition, levels = c("Control",
"Mechanical sleep disruption")))) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_base + scale_y + scale_x + sleep_scale +
theme_bw() + theme_common + hide_y +
theme(axis.text.x = element_text(color = "black", size = 14),
  legend.position = "none") +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") *
bold("_SPT")),
  color = "black", size = 5, hjust = 1, vjust = 0)
```

10.2.2 HO_VLN

```
A2 <- cuminc(Surv(age_followup, factor(status_competing)) ~ sleep_condition,
  data = FB_df %>% filter(lineage == "HO_VLN") %>%
  mutate(sleep_condition = factor(sleep_condition, levels = c("Control",
"Mechanical sleep disruption")))) %>%
```

```
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_x_only + scale_y + scale_x + sleep_scale +
theme_bw() + theme_common + hide_y +
theme(axis.text.x = element_text(color = "black", size = 14),
       legend.position = "none") +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") *
bold("_VLN")),
        color = "black", size = 5, hjust = 1, vjust = 0)
```

10.2.3 HC_MT

```
A3 <- cuminc(Surv(age_followup, factor(status_competing)) ~ sleep_condition,
             data = FB_df %>% filter(lineage == "HC_MT") %>%
             mutate(sleep_condition = factor(sleep_condition, levels = c("Control",
             "Mechanical sleep disruption")))) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_y_only + scale_y + scale_x + sleep_scale +
theme_bw() + theme_common + show_y +
theme(axis.text.x = element_text(color = "black", size = 14),
       legend.position = "none") +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Hc") * bold("_MT")),
        color = "black", size = 5, hjust = 1, vjust = 0)
```

Extract legend from the plot A3

```
shared_treatment_legend <- cowplot::get_legend(
  A3 + theme(legend.position = "bottom",
             legend.direction = "horizontal",
             legend.text = element_text(size = 13),
             legend.title = element_text(size = 14, face = "bold")))
```

Combine A1, A2, and A3 into panel “A”

```
A <- (A1 + A2 + A3 + plot_layout(nrow = 1)) +
plot_annotation(title = "Interaction between sleep condition and lineage",
               theme = theme(plot.title = element_text(hjust = 0.5, face = "bold",
               size = 16, margin = margin(b = 10))))
```

10.3 Panel B: by tumoral state

Cumulative incidence of first bud production by tumoral state.

```
B <- cuminc(Surv(age_followup, factor(status_competing)) ~ tum_state_early, data = FB_df)
%>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_xy + scale_y + tum_scale + scale_x +
```

```

theme_bw() + theme_common + show_y +
theme(axis.text.x = element_text(color = "black", size = 14),
      axis.text.y.right = element_text(color = "black", size = 14),
      axis.ticks.y.right = element_line(),
      legend.position = "none") +
plot_annotation(title = "Tumoral state effect",
                theme = theme(plot.title = element_text(hjust = 0.5, face = "bold",
                size = 16, margin = margin(b = 10))))

```

Extract legend from this panel

```

shared_tumor_legend <- cowplot::get_legend(
  B + theme(legend.position = "bottom",
            legend.direction = "horizontal",
            legend.text = element_text(size = 13),
            legend.title = element_text(size = 14, face = "bold")))

```

10.4 Panel C: by lineage

10.4.1 HO_MT

```

C1 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
             data = FB_df %>% filter(lineage == "HO_MT")) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_base + scale_y + scale_x +
theme_bw() + theme_common + hide_y + hide_x +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") * bold("_MT")),
           color = "black", size = 5, hjust = 1, vjust = 0)

```

10.4.2 HO_SPC

```

C2 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
             data = FB_df %>% filter(lineage == "HO_SPC")) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_base + scale_y + scale_x +
theme_bw() + theme_common + hide_y + hide_x +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") *
bold("_SPC")),
           color = "black", size = 5, hjust = 1, vjust = 0)

```

10.4.3 HO_SPT

```

C3 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
             data = FB_df %>% filter(lineage == "HO_SPT")) %>%

```

```
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_base + scale_y + scale_x +
theme_bw() + theme_common + hide_y + hide_x +
theme(axis.text.y.right = element_text(color = "black", size = 14),
axis.ticks.y.right = element_line()) +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") *
bold("_SPT")),
color = "black", size = 5, hjust = 1, vjust = 0)
```

10.4.4 HO_VLN

```
C4 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
data = FB_df %>% filter(lineage == "HO_VLN")) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_base + scale_y + scale_x +
theme_bw() + theme_common + hide_y + show_x +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Ho") *
bold("_VLN")),
color = "black", size = 5, hjust = 1, vjust = 0)
```

10.4.5 HC_MT

```
C5 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
data = FB_df %>% filter(lineage == "HC_MT")) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_x_only + scale_y + scale_x +
theme_bw() + theme_common + hide_y + show_x +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Hc") * bold("_MT")),
color = "black", size = 5, hjust = 1, vjust = 0)
```

10.4.6 HV_GAL

```
C6 <- cuminc(Surv(age_followup, factor(status_competing)) ~ 1,
data = FB_df %>% filter(lineage == "HV_GAL")) %>%
ggcuminc(outcome = "1") +
add_confidence_interval() +
labs_y_only + scale_y + scale_x +
theme_bw() + theme_common + show_y + show_x +
annotate("text", x = 5.8, y = 0.05, label = expression(bolditalic("Hv") *
bold("_GAL")),
color = "black", size = 5, hjust = 1, vjust = 0)
```

Create panel C combining plots C1 to C6.

```
C <- (C1 + C2 + C3) / (C4 + C5 + C6) +
  plot_layout(heights = c(1, 1)) +
  plot_annotation(title = "Lineage effect",
                  theme = theme(plot.title = element_text(
                    hjust = 0.5, face = "bold", size = 16, margin = margin(b = 10))))
```

10.5 Assembly

Add a letter tag to the first sub-plot of each panel.

```
A[[1]] <- A[[1]] + labs(tag = "A") + theme(plot.tag = element_text(size = 20, face =
"bold"))
B <- B + labs(tag = "B") + theme(plot.tag = element_text(size = 20, face = "bold"))
C[[1]][[1]] <- C[[1]][[1]] + labs(tag = "C") + theme(plot.tag = element_text(size = 20,
face = "bold"))
```

Convert each panel into a graphical object.

```
A <- patchwork::patchworkGrob(A)
```

Wrap B in patchwork before conversion to force perfect axis alignment with A and C


```
B <- patchwork::patchworkGrob(B + patchwork::plot_layout(ncol = 1))
C <- patchwork::patchworkGrob(C)
```

Combine panels and shared legends vertically into a single column, aligning axes.

```
final_FB_panel <- cowplot::plot_grid(A,
                                     shared_treatment_legend,
                                     B,
                                     shared_tumor_legend,
                                     C,
                                     ncol = 1,
                                     align = "v",
                                     axis = "lr",
                                     rel_heights = c(1, 0.1, 1, 0.1, 1.8)) +
  theme(plot.margin = margin(l = 5, r = 5, t = 5, b = 5))
```

Show the final figure.

```
print(final_FB_panel)
```



```
ggsave("figure_4.png", plot = final_FB_panel, width = 21, height = 29.7, units = "cm",
dpi = 300)
```