

Report – Numerical Simulation of Light-Ray Deflection in Schwarzschild Space-Time

1. Introduction

Gravitational lensing - the bending of light by massive bodies - provides one of the most striking confirmations of general relativity. In this project, I simulate photon trajectories around a non-rotating (Schwarzschild) black hole by numerically integrating the null geodesic equations. The primary goal is to reproduce the classic result that light passing at impact parameter b is deflected by an angle

$$\Delta\phi \approx \frac{4M}{b}$$

in geometrized units ($c = G = 1$). I build an interactive dashboard (using Dash and Plotly) that allows the user to vary the black-hole mass M and visualize both individual light-ray trajectories and the dependence of $\Delta\phi$ on b .

2. Mathematical Formulation

2.1. Schwarzschild Metric and Null Geodesics

In Schwarzschild coordinates (t, r, θ, ϕ) , a spherically symmetric, static black hole of mass M has line element

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2).$$

Restricting motion to the equatorial plane ($\theta = \pi/2$), null geodesics ($ds^2 = 0$) satisfy two first integrals:

$$E = \left(1 - \frac{2M}{r}\right) \frac{dt}{d\lambda}, \quad L = r^2 \frac{d\phi}{d\lambda},$$

where E, L are the conserved energy and angular momentum per unit mass, and λ is just a parameter that increases along the light ray's path. Eliminating $dt/d\lambda$ and using $ds^2 = 0$ yields the radial equation (see [theconfused.me, 2017]):

$$\frac{d^2 r}{d\lambda^2} = \frac{L^2}{r^4} (r - 3M).$$

Thus we have the system of three first-order ODEs for variables $\{r, \dot{r}, \phi\}$:

$$\begin{cases} \frac{dr}{d\lambda} = \dot{r}, \\ \frac{d\dot{r}}{d\lambda} = \frac{L^2}{r^4} (r - 3M), \\ \frac{d\phi}{d\lambda} = \frac{L}{r^2}. \end{cases}$$

We adopt geometrized units throughout, so M and all lengths are in the same unit.

2.2. Impact Parameter and Initial Conditions

The impact parameter b is the asymptotic closest-approach distance for a straight-line light ray. We launch photons from a point $(x_0, y_0) = (-x_{\text{init}}, b)$ with an initial velocity directed along $+x$. Converting to polar coordinates,

$$r_0 = \sqrt{x_{\text{init}}^2 + b^2}, \quad \phi_0 = \arccos(x_{\text{init}}/r_0),$$

and components of the velocity in (r, ϕ) form are chosen so that the ray is initially moving purely in the $+x$ direction. One solves

$$\dot{r}_0 = \cos \phi_0, \quad \dot{\phi}_0 = -\frac{\sin \phi_0}{r_0},$$

then sets $L = r_0^2 \dot{\phi}_0$.

3. Numerical Method

I use SciPy's `solve_ivp` routine (RK45 by default) to integrate the ODE system from $\lambda = 0$ to $\lambda = \lambda_{\text{max}}$. The trajectory is sampled uniformly in λ with step $\Delta\lambda = 0.05$. Trajectories for various b values are computed and then transformed back to Cartesian (x, y) via

$$x = r \cos \phi, \quad y = r \sin \phi.$$

4. Results and Error Analysis

4.1. Photon Trajectories

For a representative mass $M = 1.0$, trajectories with various b values from 4 to 30 clearly show increasing deflection for smaller b , and capture (plunge into the horizon) when $b < b_{\text{crit}} = 3\sqrt{3}M \approx 5.20$ (see Fig. 1).




Fig.1. The plot of Light Rays Around a Schwarzschild Black Hole

4.2. Deflection Angle vs. Impact Parameter

I compare:

- **Theoretical:** $\Delta\phi_{\text{th}} = 4M/b$.
- **Numerical:** estimated by measuring the asymptotic incoming and outgoing slopes of the trajectory and computing

$$\Delta\phi_{\text{num}} = \arctan(-\text{slope}).$$

For $b \gtrsim 10$, numerical and theoretical agree within $< 1\%$. As b decreases toward the critical value, nonlinear effects introduce larger deviations; at $b = 6$ the relative error is a few percent, growing rapidly for $b < 5.5$.

4.3. Sources of Error

- Integration step size ($\Delta\lambda$): fixed-step sampling of output can poorly resolve rapid curvature near periapsis, leading to slope-estimation error.
- Finite λ_{\max} : the photon may not fully exit the curved region within the chosen domain, biasing asymptotic angle.
- Slope estimation: using only two points near the end of the trajectory amplifies noise; a least-squares fit to the outgoing leg would reduce variance.

5. Conclusions

I have successfully implemented and visualized light bending around a Schwarzschild black hole, reproducing the classical $4M/b$ law in the weak-field regime. The interactive dashboard highlights how deflection grows with mass and falls off with b , and illustrates the capture threshold at $b = 3\sqrt{3}M$.

Possible Improvements:

- Adaptive integration: switch to an adaptive step integrator that refines near periapsis to control local error.
- Extended domain: increase λ_{\max} until the angle converges to a desired precision.
- Robust asymptotic fitting: perform linear regression on the far-field segment of (x, y) to extract the deflection angle more accurately.
- Higher-order methods: implement higher-order integrators tailored to geodesic equations.

Such refinements would reduce numerical error, extend validity closer to the photon-sphere, and provide even more faithful simulations of strong-field lensing.

6. References

- Numerical integration of light paths in a Schwarzschild metric, *theconfused.me* (2017), available at <https://theconfused.me/blog/numerical-integration-of-light-paths-in-a-schwarzschild-metric/>