


Certified homotopy tracking using the Krawczyk method

Joint work with Tim Duff



Kisun Lee (Clemson University) - kisunl@clemson.edu


The International Symposium on Symbolic and Algebraic Computation (ISSAC) 2024

Homotopy continuation


Finding solutions by tracking homotopy

$$H(t, x) = (1 - t)\gamma G(x) + tF(x), \quad t \in [0, 1]$$

Solve F (**target system**) by constructing a homotopy with G (**start system**) whose solutions are known



Homotopy continuation




Homotopy continuation

■ Predictor step

Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Predictor step


Solving ODE

$$x'(t_0) = (\partial_x F_t)^{-1} \partial_t F_t \Big|_{x=x_0, t=t_0}$$

to get the prediction.

- Corrector step

Refine the approximation using
Newton's method



Homotopy continuation

- Homotopy continuation produces **numerical solutions** (approximations) to the system (**Certification** needed)

Homotopy continuation

- Homotopy continuation produces **numerical solutions** (approximations) to the system (**Certification** needed)
- For a system F and a numerical solution $x \in \mathbb{C}^n$ to F , we say that x is **certified** if x is (possibly) contained in a small region that also contains a **unique exact solution** x^\star to F

Homotopy continuation

- Homotopy continuation produces **numerical solutions** (approximations) to the system (**Certification** needed)
- For a system F and a numerical solution $x \in \mathbb{C}^n$ to F , we say that x is **certified** if x is (possibly) contained in a small region that also contains a **unique** exact solution x^\star to F (So that x can be refined to be arbitrarily close to x^\star using a finite procedure)

Homotopy continuation

- Homotopy continuation produces **numerical solutions** (approximations) to the system (**Certification** needed)
- For a system F and a numerical solution $x \in \mathbb{C}^n$ to F , we say that x is **certified** if x is (possibly) contained in a small region that also contains a unique exact solution x^\star to F (So that x can be refined to be arbitrarily close to x^\star using a finite procedure)
(a posteriori certification)


Certified homotopy tracking

- Post-processing certification does not verify the correctness of path tracking

Certified homotopy tracking


- Post-processing certification does not verify the correctness of path tracking
- In some algorithms or computations (e.g. monodromy or Galois group computation), the order and correspondence of solutions are important

Why certified homotopy tracking?




Why certified homotopy tracking?

- Scenario 1) Path jumping



Why certified homotopy tracking?


- Scenario 1) Path jumping



Why certified homotopy tracking?

- Scenario 1) Path jumping

Path jumping may result in **false multiple solutions** and a **lost solution**.




Why certified homotopy tracking?

- Scenario 1) Path jumping

Path jumping may result in **false multiple solutions** and a **lost solution**.

- Scenario 2) Path crossing




Why certified homotopy tracking?

- Scenario 1) Path jumping

Path jumping may result in **false multiple solutions** and a **lost solution**.

- Scenario 2) Path crossing




Why certified homotopy tracking?

- Scenario 1) Path jumping

Path jumping may result in **false multiple solutions** and a **lost solution**.

- Scenario 2) Path crossing

Path crossing **loses correspondence** between the start solution and the target solution.



Why certified homotopy tracking?


- Scenario 1) Path jumping

Path jumping may result in **false multiple solutions** and a **lost solution**.

- Scenario 2) Path crossing

Path crossing **loses correspondence** between the start solution and the target solution.


Crucial for some algorithms (e.g. monodromy), and hard to detect.



Why certified homotopy tracking?

- Certified homotopy guarantees tracking the same path from start to finish **without jumping or crossing**


**(Beltran-Leykin 2012 2013,
Hauenstein-Haywood-Liddell 2014,
van der Hoeven 2015,
Xu-Burr-Yap 2018,**



Why certified homotopy tracking?


- Certified homotopy guarantees tracking the same path from start to finish **without jumping or crossing**

(Beltran-Leykin 2012 2013,
Hauenstein-Haywood-Liddell 2014,
van der Hoeven 2015,
Xu-Burr-Yap 2018,
Guillemot-Lairez 2024)



Interval arithmetic

- interval arithmetic




Interval arithmetic

- interval arithmetic

for an arithmetic operator \odot , define

$$[a, b] \odot [c, d] = \{x \odot y \mid x \in [a, b], y \in [c, d]\}$$

for conservative and guaranteed computation



Interval arithmetic

- interval arithmetic

for an arithmetic operator \odot , define


$$[a, b] \odot [c, d] = \{x \odot y \mid x \in [a, b], y \in [c, d]\}$$

for conservative and guaranteed computation

ex) $f(x) = x^3 - 2x$ on $x \in [-2, 2]$.

Then, $f(x) \in [-2, 2]^3 - 2 \cdot [-2, 2] = [-12, 12]$

(much larger than the actual image $[-4, 4]$)



Interval arithmetic

■ interval arithmetic

for an arithmetic operator \odot , define

$$[a, b] \odot [c, d] = \{x \odot y \mid x \in [a, b], y \in [c, d]\}$$

for conservative and guaranteed computation


ex) $f(x) = x^3 - 2x$ on $x \in [-2, 2]$.

$$\text{Then, } f(x) \in [-2, 2]^3 - 2 \cdot [-2, 2] = [-12, 12]$$

(much larger than the actual image $[-4, 4]$)

If a certain property is true for $[-12, 12]$,

then it is true for any point in the actual image of $f(x)$ over $[-2, 2]$



Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

Theorem (Krawczyk 1969). For the Krawczyk operator the following hold :

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

Theorem (Krawczyk 1969). For the Krawczyk operator the following hold :

1. If $K_{x,Y}(I) \subset I$, then there is a solution of F in I (**existence** of a solution)

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

Theorem (Krawczyk 1969). For the Krawczyk operator the following hold :

1. If $K_{x,Y}(I) \subset I$, then there is a solution of F in I (**existence** of a solution)
2. If I has a solution and $\sqrt{2} \|Id - Y \square \frac{dF}{dx}(I)\| < 1$, then there is a solution of F in I , and it is unique (**uniqueness** of a solution)

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix}$$

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix} \text{ with } x = \begin{bmatrix} 1.4 \\ 1 \end{bmatrix} \text{ and } Y = JF(x)^{-1}$$

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix} \text{ with } x = \begin{bmatrix} 1.4 \\ 1 \end{bmatrix} \text{ and } Y = JF(x)^{-1}$$

$$\text{Then, } K_{x,Y}(I) = \begin{bmatrix} [1.3476\dots, 1.4810\dots] \\ [.89333\dots, 1.1067\dots] \end{bmatrix}$$

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix} \text{ with } x = \begin{bmatrix} 1.4 \\ 1 \end{bmatrix} \text{ and } Y = JF(x)^{-1}$$

$$\text{Then, } K_{x,Y}(I) = \begin{bmatrix} [1.3476\dots, 1.4810\dots] \\ [.89333\dots, 1.1067\dots] \end{bmatrix} \subset I.$$

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix} \text{ with } x = \begin{bmatrix} 1.4 \\ 1 \end{bmatrix} \text{ and } Y = JF(x)^{-1}$$

$$\text{Then, } K_{x,Y}(I) = \begin{bmatrix} [1.3476\dots, 1.4810\dots] \\ [.89333\dots, 1.1067\dots] \end{bmatrix} \subset I.$$

Therefore, I contains a root of F (which is $(\sqrt{2}, 1)$),

Krawczyk method

Combine interval arithmetic and Newton's method. Define the **Krawczyk operator**

$$K_{x,Y}(I) = x - YF(x) + \left(Id - Y \square \frac{dF}{dx}(I) \right)(I - x)$$

F : a square differentiable system of polynomial equations

I : an interval

$\square F(I) \supset \{F(x) \mid x \in I\}$: an interval extension of F over I

x : a point in I

Y : an invertible matrix

ex) $F = \begin{bmatrix} x^2 - y - 1 \\ -\frac{1}{2}x^2 + y^2 \end{bmatrix}$ has a root $(\sqrt{2}, 1)$.

(But let us pretend not to know!)

$$I = \begin{bmatrix} [1.2, 1.6] \\ [0.8, 1.2] \end{bmatrix} \text{ with } x = \begin{bmatrix} 1.4 \\ 1 \end{bmatrix} \text{ and } Y = JF(x)^{-1}$$

$$\text{Then, } K_{x,Y}(I) = \begin{bmatrix} [1.3476\dots, 1.4810\dots] \\ [.89333\dots, 1.1067\dots] \end{bmatrix} \subset I.$$

Therefore, I contains a root of F (which is $(\sqrt{2}, 1)$), uniquely.

Q. Can we use the Krawczyk method for certified homotopy tracking?

Q. Can we use the Krawczyk method for certified homotopy tracking?

A. Consider the Krawczyk method “for a system with parameters**”**

Krawczyk method

Combine interval arithmetic and Newton's method.

Define the **Krawczyk operator**

$$K_{x,Y}(I, \textcolor{red}{T}) = x - Y \square H(x, \textcolor{red}{T}) + \left(Id - Y \square \frac{\partial H(I, \textcolor{red}{T})}{\partial x} \right) (I - x).$$

$H(x, t)$: a homotopy with a parameter t from

$\mathbb{C}^n \times [0,1]$ to \mathbb{C}^n

(e.g. $H(x, t) = t\gamma G(x) + (1 - t)F(x)$)

I : an interval

T : an interval contained in $[0,1]$

Krawczyk method

Combine interval arithmetic and Newton's method.
Define the **Krawczyk operator**

$$K_{x,Y}(I, \textcolor{red}{T}) = x - Y \square H(x, \textcolor{red}{T}) + \left(Id - Y \square \frac{\partial H(I, \textcolor{red}{T})}{\partial x} \right) (I - x).$$

$H(x, t)$: a homotopy with a parameter t from
 $\mathbb{C}^n \times [0,1]$ to \mathbb{C}^n

(e.g. $H(x, t) = t\gamma G(x) + (1 - t)F(x)$)

I : an interval

T : an interval contained in $[0,1]$

Theorem (Duff-L.) For the Krawczyk operator,
the following hold :

1. If $K_{x,Y}(I, \textcolor{red}{T}) \subset I$, then there is a solution of
 $H(x, t)$ in I for all $t \in T$
(**existence** of a solution path)

2. If I has a solution and

$$\sqrt{2} \left\| Id - Y \square \frac{\partial H(I, \textcolor{red}{T})}{\partial x} \right\| < 1, \text{ then}$$


for all $t \in T$, there is a solution of F in I ,
and it is unique
(**uniqueness** of a solution path)


$F(x)$


$G(x)$


$t = 1$

$t = 0$

$F(x)$ $G(x)$ $t = 1$ $t = 0$ 

$F(x)$ $G(x)$ $t = 1$ $t = 0$ 

$F(x)$ $G(x)$ $t = 1$ $t = 0$  $I \times T$




$F(x)$

$G(x)$

$t = 1$

$t = 0$




$F(x)$

$G(x)$

$t = 1$

$t = 0$





$F(x)$


$G(x)$

$t = 1$

$t = 0$









Preconditioning

Constructing a tilted interval box




Preconditioning

Constructing a tilted interval box




Preconditioning

Constructing a tilted interval box




Preconditioning

Constructing a tilted interval box




Preconditioning

Constructing a tilted interval box




Preconditioning

Constructing a tilted interval box



Preconditioning


Constructing a tilted interval box



Construct a tilted interval box
centered at the line segment

Preconditioning


Constructing a tilted interval box



Construct a tilted interval box
centered at the line segment



Preconditioning

Constructing a tilted interval box



Preconditioning

Constructing a tilted interval box




After the construction, apply the **shearing map** to make the rectangular interval box to avoid **overestimation** from interval arithmetic

Algorithm

Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $F(x)$
 $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant



Algorithm

Input :

- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test passed **then**

Set $r = \lambda r$ and $dt = \lambda dt$

Set $t_0 = t_1$ and $t_1 = t_0 + dt$



Algorithm

Input :


- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test passed **then**

Set $r = \lambda r$ and $dt = \lambda dt$

Set $t_0 = t_1$ and $t_1 = t_0 + dt$




Algorithm

Input :

- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test failed **then**



Algorithm


Input :

- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test failed **then**

Set $r = \frac{1}{\lambda}r$ and $dt = \frac{1}{\lambda}dt$



Algorithm


Input :

- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test failed **then**

Set $r = \frac{1}{\lambda}r$ and $dt = \frac{1}{\lambda}dt$



Algorithm


Input :

- $H(x, t) = F(x; p(t))$ a parameter homotopy from $t = 0$ to $t = 1$
- $x(t)$: a nonsingular solution path in \mathbb{C}^n
- I_r : a square interval box with a radius r whose midpoint $x_0 = m(I_r)$ approximates $x(0)$
- $dt \in (0, 1)$: the initial step size
- $T_{t_0, dt} = [t_0, t_1]$ where $t_0 = 0$ and $t_1 = t_0 + dt$
- $\lambda > 1$: a scaling constant

While $t_0 < 1$ **do**

If Krawczyk test failed **then**

Set $r = \frac{1}{\lambda}r$ and $dt = \frac{1}{\lambda}dt$



Algorithm

Algorithm 3 Krawczyk homotopy (tilted)

Input:

- A parameter homotopy $H(x, t) = F(x; p(t)) : \mathbb{C}^n \times [0, 1] \rightarrow \mathbb{C}^n$ analytic in x and linear in p ,
- a point x_0 approximating $x(0)$, for some nonsingular solution path $x(t) : [0, 1] \rightarrow \mathbb{C}^n$ such that $H(x(t), t) = 0$,
- a positive number $r > 0$ for the initial radius,
- a time-step size $dt \in (0, 1)$, and
- a scaling constant $\lambda > 1$.

Output: A certified approximation of $x(1)$.

- 1: Set $t_0 = 0$ and $t_1 = dt$.
- 2: Run **Preconditioning**($H(x, t), r, t_0, t_1, x_0$) to compute $x_1, \hat{H}(x, t), I_r$ and $T_{t_0, dt}$.
- 3: Compute $Y := \partial_x \hat{H}(0, t_0)^{-1}$.
- 4: **while** $t_0 < 1$ **do**
- 5: Run Krawczyk test with $K_{0,Y}(I_r, T_{t_0, dt})$.
- 6: **if** Krawczyk test passed **then**
- 7: Set $r = \lambda r$ and $dt = \lambda dt$.
- 8: Refine x_1
- 9: Set $x_0 = x_1, t_0 = t_1$, and $t_1 = t_0 + dt$.
- 10: Run **Preconditioning**($H(x, t), r, t_0, t_1, x_0$) to compute $x_1, \hat{H}(x, t), I_r$ and $T_{t_0, dt}$.
- 11: Compute $Y := \partial_x \hat{H}(0, t_0)^{-1}$.
- 12: **else**
- 13: Set $r = \frac{1}{\lambda}r$ and $dt = \frac{1}{\lambda}dt$.
- 14: Run **Preconditioning**($H(x, t), r, t_0, t_1, x_0$) to compute $x_1, \hat{H}(x, t), I_r$ and $T_{t_0, dt}$.
- 15: **end if**
- 16: **end while**
- 17: Find a point x_1 by refining $s(1)$ with the system $H(x, 1)$.
- 18: Return x_1 .

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$.

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$. Consider a fixed constant $L \geq \parallel \square \partial_x^2 H(I_1, [0,1]) \parallel$.

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$. Consider a fixed constant $L \geq \parallel \square \partial_x^2 H(I_1, [0,1]) \parallel$. Assuming that $x(t)$ is nonsingular at any $t \in [0,1]$,

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$. Consider a fixed constant $L \geq \parallel \square \partial_x^2 H(I_1, [0,1]) \parallel$.

Assuming that $x(t)$ is nonsingular at any $t \in [0,1]$, if dt and r satisfy

$$1 - \|Y\| \cdot \frac{dt}{r} \cdot \|G(x^\star) - F(x^\star)\| > 0 \quad \text{and} \quad \sqrt{nr^2 + dt^2} < \frac{1}{\sqrt{2} \cdot \|Y\| \cdot L}$$

where $Y = \partial_x H(x^\star, t_0)^{-1}$, then the Krawczyk method succeeds.

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$. Consider a fixed constant $L \geq \parallel \square \partial_x^2 H(I_1, [0,1]) \parallel$.

Assuming that $x(t)$ is nonsingular at any $t \in [0,1]$, if dt and r satisfy

$$1 - \|Y\| \cdot \frac{dt}{r} \cdot \|G(x^\star) - F(x^\star)\| > 0 \quad \text{and} \quad \sqrt{n r^2 + dt^2} < \frac{1}{\sqrt{2} \cdot \|Y\| \cdot L}$$

where $Y = \partial_x H(x^\star, t_0)^{-1}$, then the Krawczyk method succeeds.

Termination

Theorem (Duff-L.). Let $H(x, t) = t\gamma G(x) + (1 - t)F(x)$ be twice differentiable with an exact solution x^\star to $H(x, t_0)$. Consider a fixed constant $L \geq \|\square \partial_x^2 H(I_1, [0,1])\|$.

Assuming that $x(t)$ is nonsingular at any $t \in [0,1]$, if dt and r satisfy

$$1 - \|Y\| \cdot \frac{dt}{r} \cdot \|G(x^\star) - F(x^\star)\| > 0 \quad \text{and} \quad \sqrt{nr^2 + dt^2} < \frac{1}{\sqrt{2} \cdot \|Y\| \cdot L}$$

where $Y = \partial_x H(x^\star, t_0)^{-1}$, then the Krawczyk method succeeds.

Corollary (Duff-L.). Assuming that $x(t)$ is nonsingular at any $t \in [0,1]$, the tracking terminates in finitely many steps for the homotopy $H(x, t)$.

Implementation detail

Implementation in Macaulay2

Not the optimal version yet due to incomplete implementation of interval arithmetic in M2 (a proof-of-concept implementation)

A scaling constant $\lambda = 3$ is used

Experiments

- Comparison with the known method

The certified homotopy (**Beltran-Leykin 2012 2013**) is based on alpha theory. Compare the number of steps for each solution path for benchmark problems. The initial radius and step-size, $r = 0.4$ and $dt = 0.1$ were used

Experiments

- Comparison with the known method

The certified homotopy (**Beltran-Leykin 2012 2013**) is based on alpha theory. Compare the number of steps for each solution path for benchmark problems. The initial radius and step-size, $r = 0.4$ and $dt = 0.1$ were used

System	# roots	Average	Max	Min	Average(BL 2012)
Random (2,2,2)	8	317.75	509	203	198.5
Random (2,2,2,2)	16	563.25	1257	211	813.812
Random (2,2,2,2,2)	32	675.3125	5119	209	1542.5
Random (2,2,2,2,2,2)	64	1166.421875	5415	267	2211.58

- In many examples, Krawczyk homotopy requires less number of steps in average than BL 2012

Experiments

- Performance according to changes in step size and radius

Solved the random (2,2,2) problem by changing the step size dt and the radius r , and measured the iterations.

Experiments

■ Performance according to changes in step size and radius

Solved the random (2,2,2) problem by changing the step size dt and the radius r , and measured the iterations.

	Average	Max	Min		Average	Max	Min
$(dt, r) = (.2,.4)$	134.75	181	83	$(.02,.04)$	135.625	189	85
$(.4,.4)$	96.5	170	44	$(.04,.04)$	99.5	175	48
$(.6,.4)$	98.625	166	32	$(.06,.04)$	98.875	171	34
$(.8,.4)$	96.75	194	24	$(.08,.04)$	99.5	195	28
$(.9,.4)$	100.5	216	23	$(.09,.04)$	101.375	215	24

Experiments

■ Performance according to changes in step size and radius

Solved the random (2,2,2) problem by changing the step size dt and the radius r , and measured the iterations.

	Average	Max	Min		Average	Max	Min
$(dt, r) = (.2,.4)$	134.75	181	83	$(.02,.04)$	135.625	189	85
$(.4,.4)$	96.5	170	44	$(.04,.04)$	99.5	175	48
$(.6,.4)$	98.625	166	32	$(.06,.04)$	98.875	171	34
$(.8,.4)$	96.75	194	24	$(.08,.04)$	99.5	195	28
$(.9,.4)$	100.5	216	23	$(.09,.04)$	101.375	215	24

Experiments

- Performance according to changes in step size and radius

Solved the random (2,2,2) problem by changing the step size dt and the radius r , and measured the iterations.

	Average	Max	Min		Average	Max	Min
$(dt, r) = (.2,.4)$	134.75	181	83	$(.02,.04)$	135.625	189	85
$(.4,.4)$	96.5	170	44	$(.04,.04)$	99.5	175	48
$(.6,.4)$	98.625	166	32	$(.06,.04)$	98.875	171	34
$(.8,.4)$	96.75	194	24	$(.08,.04)$	99.5	195	28
$(.9,.4)$	100.5	216	23	$(.09,.04)$	101.375	215	24

- The performance of the algorithm depends on the ratio between dt and r (rather than their specific values)

Summary

- Certified homotopy tracking guarantees the correctness of homotopy tracking without path jumping or crossing
- Certified tracking assisted by interval arithmetic may be more competitive to alpha theory-based known methods
- Robust implementation to deal with larger size examples is expected

Thanks For Your Attention!

(<https://arxiv.org/abs/2402.07053>)