



# Current concepts in breast cancer genomics: An evidence based review by the CGC breast cancer working group

Katherine B. Geiersbach\*, Hui Chen, Rajyasree Emmadi, Gloria T. Haskell, Xinyan Lu, Yajuan J. Liu, Karen Swisshelm

Mayo Clinic, 200 First Street SW, Rochester, MN 55905, United States

## ARTICLE INFO

### Article history:

Received 22 September 2019

Revised 18 January 2020

Accepted 5 February 2020

## ABSTRACT

**Background:** Genomic abnormalities in breast cancer have been described according to diverse conceptual frameworks, including histologic subtypes, clinical molecular subtypes, intrinsic DNA, RNA, and epigenetic profiles, and activated molecular pathways.

**Methods:** The Cancer Genomics Consortium (CGC) Breast Cancer Workgroup performed an evidence based literature review to summarize current knowledge of clinically significant genomic alterations in breast cancer using CGC levels of evidence. Targetable or disease-defining alterations were prioritized.

**Results:** We summarized genomic alterations in breast cancer within a framework of existing clinical tools for diagnosis, risk stratification, and therapeutic management. Using CGC levels of evidence, we catalog copy number profiles, gene expression profiles, and mutations in clinically significant genes. We also describe emerging molecular markers such as methylation profiling and immunotherapy biomarkers.

**Conclusion:** A summary of currently available information on breast cancer genomics will enhance precision medicine by serving as an interpretive resource for clinical laboratory geneticists, providing a foundation for future practice guidelines, and identifying knowledge gaps to address in future research.

© 2020 Elsevier Inc. All rights reserved.

## Introduction and Background

Current diagnosis and management of breast cancer relies primarily upon histology, clinical and pathologic staging, and a limited set of biomarkers for hormone therapy and anti-HER2 therapy. The principal biomarkers are estrogen receptor alpha (ER- $\alpha$ ) (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER-2/neu, *ERBB2*). Pathologic work up of invasive tumors includes immunohistochemistry to evaluate for ER and PR expression, and immunohistochemistry and/or in situ hybridization (ISH) for determination of HER2 status. Histopathology, clinical data, and biomarker data are integrated to determine a patient's prognostic stage group. The clinical prognostic stage group includes all available clinical and imaging data as well as histopathology and biomarker status on the diagnostic breast biopsy and/or distant metastases. The pathologic prognostic stage group incorporates histopathologic findings from surgical resection, including the final tumor size, the status of surgical margins and any excised lymph nodes, and distant metastatic disease, if applicable. In the

neoadjuvant setting, therapy is administered after the diagnostic biopsy but before tumor resection, and clinical prognostic staging is employed; the pathologic stage post neoadjuvant treatment is used to assess response to treatment and to guide post-surgery adjuvant treatment. Additional testing for risk stratification and therapeutic management typically includes a proliferation marker (mitotic index and/or Ki67 expression), and for a subset of patients with early stage ER positive, HER2 negative tumors, clinically approved gene expression profiling ("genomic") assays are performed on a tumor sample to determine the risk of tumor recurrence. Surgery is the mainstay of breast cancer treatment, and aside from *HER2* (*ERBB2*) amplification, specific genomic abnormalities are most important in the setting of inoperable or metastatic disease. Genomic profiling is becoming more relevant as additional actionable genetic abnormalities are discovered. This review summarizes recent advances and current concepts in breast cancer genomics.

## Methods

We performed a systematic review to identify original articles on human breast cancer genetics or genomics published in priority journals in the English language since 2010. The full search

\* Corresponding author.

E-mail address: [geiersbach.katherine@mayo.edu](mailto:geiersbach.katherine@mayo.edu) (K.B. Geiersbach).

**Table 1**  
Cancer Genomics Consortium Levels of Evidence.

Tier	Data Source(s)	Interpretation
1	FDA approved therapies, professional guidelines, multiple large clinical studies	Strong evidence supporting clinical utility of variant(s) for diagnosis, selection of therapies, or predicting disease outcome
2	One large study or multiple case reports	Emerging evidence supporting clinical utility of variant(s)
3	Case reports or expert opinion	Unknown clinical significance
4	Published evidence indicating lack of pathogenicity of variant(s)	Benign or likely benign

strategy performed in the Ovid MEDLINE database is shown in the Data Supplement. Additionally, we manually searched major journals, textbooks, and clinical practice guidelines to capture relevant articles regardless of publication date. We excluded non-malignant neoplasms, non-epithelial neoplasms, and metastases to the breast originating in other organs. We prioritized articles according to size of the study, the inclusion of clinical data, and the clinical significance of the aberrations described. We then triaged articles according to the type of aberration described (DNA copy number, RNA expression profiling, somatic sequence variation, or other mutations).

Somatic variants were categorized at the gene level according to the Cancer Genomics Consortium levels of evidence shown in Table 1. We excluded Tier 3 and Tier 4 variants from this review. We included genes associated with an inherited risk for breast cancer according to current literature and guidelines of the National Comprehensive Cancer Network (NCCN) because inherited mutations can also be detected in cancers, and hence mutations in these genes should be regarded as either somatic or germline in origin.

### Disease-Defining molecular alterations

While rare, a few recurrent gene rearrangements define rare subtypes of invasive breast carcinoma. The *ETV6-NTRK3* fusion is associated with secretory breast carcinoma [1,2], and the *MYB-NFIB* fusion is associated with adenoid cystic carcinoma [3]; each accounts for less than 1% of invasive carcinomas. Loss of expression of the cell adhesion molecule cadherin 1 (*CDH1*) is observed in almost all invasive lobular carcinomas as well as some carcinomas with mixed ductal and lobular histologic features [4]. *CDH1* mutations are usually truncating and most frequently co-occur with heterozygous loss of 16q (either by deletion or copy neutral LOH) to induce complete loss of *CDH1* expression. While the mechanisms for loss of *CDH1* expression are heterogeneous, comprehensive genomic profiling of lobular carcinomas has shown inactivation of *CDH1* in virtually all cases [5]. Loss of *CDH1* function is clinically relevant not only for diagnosis but also for patient management, since germline *CDH1* mutations are associated with a hereditary predisposition to breast and gastric cancers with a characteristic infiltrating growth pattern and characteristic “signet ring” cells [6].

Other than the two recurrent gene fusions and *CDH1* mutations associated with loss of protein expression, genetic abnormalities in breast cancer are not specific for a particular histopathologic diagnosis. The spectrum of mutations does vary in frequency across histologic subtypes, however. *TP53* mutations are especially common in medullary carcinomas and triple negative carcinomas, and *GATA3* mutations are more common in mucinous carcinomas [7]. In addition to *CDH1* mutations, lobular carcinomas are enriched for *PTEN* mutations, which are mutually exclusive with *PIK3CA* mutations; additionally, *FOXA1* mutations are more frequent and *GATA3* mutations are less frequent in lobular carcinomas, suggesting differences in estrogen pathway modulation between lobular carcinomas and ductal carcinomas [5]. Frequent expression of *FOXA1* and *GATA3* in breast cancer also makes these markers helpful in the diagnostic workup of breast cancer [5]. Lobular carcinomas also more frequently harbor *ERBB2* and *ERBB3* sequence variants [8]. Aside from a close association with loss of *CDH1* expression, however, there are no specific genomic aberrations that are diagnostic of lobular carcinoma.

### Clinical molecular subtypes

Three biomarkers are an essential part of the diagnostic workup and clinical management of breast cancer: estrogen receptor alpha (*ER-α*), progesterone receptor (PR), and human epidermal growth factor receptor 2 (*HER-2/neu*, commonly referred to as *HER2*). Expression of ER, PR, and *HER2* is measured by immunohistochemistry (IHC). In situ hybridization (ISH) is also used to determine *HER2* status by the presence or absence of *ERBB2* (*HER2*) gene amplification, since *ERBB2* amplification is very closely correlated with *HER2* overexpression [9]. In the United States, evaluation of biomarkers is performed either according to FDA approved criteria or according to current guidelines published by the American Society of Clinical Oncology and the College of American Pathologists (Table 2) [10,11]. Diagnostic evaluation of breast cancer specimens includes the determination of histologic subtypes, tumor grade, and stage, as well as an evaluation of hormone receptor status and *HER2* status; a mitotic index or evaluation of Ki-67 is typically also performed as a marker of proliferation. The majority of breast cancers express estrogen and progesterone receptors,

**Table 2**  
Clinical Molecular Subtypes of Breast Cancer.

Clinical molecular subtype	Estrogen receptor IHC [10]	Progesterone receptor IHC [10]	HER2 IHC [11]	HER2 ISH [11]
Hormone receptor positive, HER2 negative	≥1% positive nuclear staining (1–10% nuclear staining is considered “low positive”) [10]	Any staining pattern	Negative (0 or 1+) or equivocal (2+) staining with negative ISH	HER2/centromere ratio <2.0 and HER2/cell <4.0 (Group 5) or equivocal (2+) IHC staining with negative IHC-guided ISH score in Groups 2, 4, or 5
HER2 positive	Any staining pattern	Any staining pattern	Positive (3+) or equivocal (2+) staining with positive ISH	HER2/centromere ratio ≥2.0 and HER2/cell ≥4.0 (Group 1) or equivocal (2+) reflex IHC staining with ISH score in Groups 1 or 3
Triple negative	<1% positive nuclear staining	<1% positive nuclear staining	Negative (0 or 1+) or equivocal (2+) staining with negative ISH	HER2/centromere ratio <2.0 and HER2/cell <4.0 (Group 5) or equivocal (2+) IHC staining with negative IHC-guided ISH score in Groups 2, 4, or 5

and 15–20% of breast cancers overexpress HER2 on the cell membrane, while a minority (15–20%) lack expression of estrogen receptors, progesterone receptors, or HER2 receptors and are called “triple negative.” Triple negative breast cancer is currently a very active area for research and drug development. Breast cancers that lack expression of ER, PR, or HER2 (and/or: lack *ERBB2* amplification) are quite heterogeneous in terms of their expression profile and their genomic signatures [12]. Triple negative breast cancers have the worst overall prognosis, but the genomic profiles of this subgroup are quite heterogeneous at the DNA level and represent a mix of tumors with good and poor prognoses.

### Intrinsic molecular subtypes

In 2000, Perou and colleagues published their seminal study on the hierarchical clustering of differential expression levels for over 1700 genes in breast cancer; this study established five intrinsic subtypes of breast cancer (luminal A, luminal B, HER2 enriched, basal like, and normal like) [13]. Perou's five intrinsic subtypes were based on expression levels of genes associated with cellular proliferation, basal vs. luminal epithelium, estrogen responsiveness, and HER2 activation. The “normal” pattern described by Perou has since been challenged as a potential artifact of inadequate tumor enrichment, and most researchers now accept four intrinsic subtypes of breast cancer: Luminal A, Luminal B, HER2-enriched, and Basal-like [14]. More recently, Perou and others have described a “claudin low” subtype with low expression of proliferation markers and high expression of genes involved in the immune system response and multiple other gene families [15–17]. Perou's intrinsic subtypes have been reproduced in numerous subsequent genomic studies on breast cancer and remain a core conceptual framework for the genomic classification of breast cancers [18,19].

Hormone receptor expression most likely precedes and promotes carcinogenesis in luminal cells of the breast [20]. Expression of hormone receptors and responsiveness to hormone therapy in breast cancers evolve with treatment, and secondary resistance mutations in *ESR1* are common [21–23]. Recent large genomic studies have confirmed that hormone receptor positive tumors correlate well (but not perfectly) with the luminal intrinsic subtypes [18,19]. Rare breast cancers with *ESR1* (*ERα*) expression can show a basal rather than a luminal molecular subtype due to aberrant expression of an alternatively spliced form referred to as EΔ7. This form of *ERα* is missing exon 7 and has a dominant negative function in repressing *ERα*-responsive genes [24].

Like hormone receptor expression, HER2 expression may precede and promote carcinogenesis. Most but not all breast cancers with HER2 overexpression show evidence of *ERBB2* (*HER2*) gene amplification. Rare activating mutations also occur and can be targeted with small molecule inhibitors (lapatinib, afatinib, and others) [25]. Tumors with a HER2 positive clinical molecular subtype are a mix of HER2 enriched and luminal intrinsic subtypes; the latter are typically “Luminal B” and characterized by a higher proliferation index.

Basal like breast cancers are defined by expression of basal keratins and low or absent expression of luminal keratins. This subgroup has been found to be heterogeneous at the molecular level and includes the aforementioned “claudin low” subtype. All are characterized by minimal or absent expression of genes associated with estrogen responsiveness, and minimal or absent expression of HER2; the associated clinical molecular subtype is most often triple negative.

### Genomic profiling of RNA expression in the clinical setting

A number of RNA-based genomic assays are clinically approved for disease stratification in patients with ER-positive, HER2-

negative disease. These assays were designed to generate a score reflecting the risk for recurrent disease based on the overall expression profile observed in a breast cancer sample, resulting in better prognostic stratification for these patients, and aiding therapeutic decisions. Commonly referred to as “genomic assays,” these RNA expression assays determine the relative expression of luminal and basal keratins, proliferation markers, HER2 and associated genes, genes involved in hormone signaling, and genes associated with the immune response, together presenting an overall picture of the activated gene pathways and the proliferation rate of a cancer. Oncotype DX® is the most frequent assay performed in the United States (Genomic Health, Redwood City, California), and it has been endorsed by several professional organizations including the American Joint Committee on Cancer (AJCC) and the National Comprehensive Cancer Network (NCCN). The TAILORx trial included long term follow up of patients whose tumors were tested by Oncotype DX, and confirmed a low risk of recurrence for most patients with an intermediate or low risk score, particularly for older patients without additional clinical risk factors [26,27]. The MammaPrint assay was developed before Oncotype DX and is widely used in European countries; testing is offered by Agendia (Amsterdam, The Netherlands). The MINDACT trial demonstrated the prognostic utility of the MammaPrint assay [28,29], and this assay is endorsed by multiple professional organizations in Europe and the United States. The MammaPrint assay was clinically validated for patients with either node negative or node positive disease. The Oncotype DX assay was originally validated for breast cancer patients with node negative disease as part of the National Surgical Adjuvant Breast and Bowel Project (NSABP B-14 and B-20) [27,30,31], but Oncotype DX can also stratify risk in patients with node positive disease, as demonstrated by the Southwest Oncology Group (SWOG) [32,33]. A third gene expression assay is Prosigna, previously known as the PAM50 assay; this is licensed to independent CLIA-certified laboratories by Nanostring (Seattle, Washington). The Prosigna assay generates a risk of recurrence score, but it is also useful for intrinsic molecular subtyping [14]. The Blueprint assay from Agendia works similarly to generate an intrinsic subtype (Luminal-type, HER2-type, or Basal-type) using gene expression profiling. Additional RNA expression profiling assays include the EndoPredict assay [34,35] and the Breast Cancer Index [36–39], both of which are used to generate a risk of recurrence similar to the aforementioned assays. Immunohistochemical markers are also useful for predicting prognosis in breast cancer. Beyond the clinical molecular subtypes, quantification of immunohistochemical staining and other protein expression assays can generate prognostic information; the most common of these are the IHC4 assay [40] and the Magee Equation recurrence score using combined quantitative information from ER, PR, HER2, and Ki-67 immunohistochemistry [41,42]. Another expression assay that is performed on the protein level is measurement of levels of urokinase-type plasminogen activator (uPA)/plasminogen activator inhibitor-1 (PAI-1) by using an enzyme-linked immunosorbent assay (ELISA) [43]. The uPA/PAI-1 method is fairly inexpensive and simple to perform, and it has been endorsed and employed mainly in European countries to identify breast cancer patients at an elevated risk for recurrence, similar to RNA expression assays. While extended clinical use of gene profiling assays is advocated for by some, genomic expression profiling does not always provide helpful or accurate information beyond the intended use of the assay [44], and the clinical value of genomic assays appears to be quite limited for those breast cancers with low grade histopathology and unequivocal biomarker status [41]. A list of clinically approved genomic RNA expression profiling assays is shown in Table 3.

**Table 3**

Assays Used to Predict Risk of Recurrence in Hormone Receptor Positive, HER2 Negative Breast Cancer.

Assay	Manufacturer	AJCC Evidence Level	Eligible Tumor Type(s)	Description
Oncotype DX®	Genomic Health	1	Node negative	21-gene assay categorizes cancers as low, intermediate, or high risk
MammaPrint	Agendia	2	Node negative or positive < 5 cm	70-gene assay, low or high risk
Prosigna®	Nanostring	2	Node negative, tamoxifen treated	50-gene assay, low or high risk
PAM50				
EndoPredict®	Myriad Genetics	2	Node negative or positive (1–3 nodes), pre- or postmenopausal	8-gene assay combined with tumor size and nodal status to assign risk score; low or high risk
Breast Cancer Index	Biotheranostics	2	Node negative, untreated or tamoxifen treated	2-gene assay generates H/I index associated with low or high risk

Evidence levels for different RNA based gene expression assays assigned by the American Joint Committee on Cancer. Part XI: Chapter 48, Breast. In: M.B. Amin et al. (eds.), AJCC Cancer Staging Manual, Eighth Edition, American College of Surgeons, Chicago, IL, USA.

### Genomic profiling of copy number alterations (CNAs) and loss of heterozygosity (LOH)

Early genetic studies on breast cancer employed karyotyping, FISH, flow cytometry, and comparative genomic hybridization (CGH) testing to describe recurrent chromosomal abnormalities. Such studies established many recurrent genomic abnormalities in breast cancer, including frequent hyperdiploidy [45–47], polyclonality [48–53], and gene amplification including *ERBB2* [9,54–56] and/or *MYC* [56–58]. The latter abnormality, amplification, could be observed in homogeneously staining regions of chromosomes in karyotype preparations of G-banded metaphase cell preparations [59–61]. Perhaps due to differences in short term and long term cytogenetic culture preparations [50], and perhaps also due to a limited number of samples studied, early studies yielded different and sometimes conflicting information on recurrent copy number alterations (CNAs) in breast cancer [53,62,63]. Array CGH technology allowed for the study of breast cancer genomics without the requirement for cell culture [62–68]. Clear patterns emerged regarding the prognostic implications of CNAs in breast cancer. Early array CGH studies also established an important link between gene amplification and overexpression [69].

Two large genomic studies published in 2012 yielded a wealth of information on the landscape of copy number alterations and allelic imbalances in breast cancer [18,19]. These studies demonstrated many of the recurrent copy number abnormalities that had been previously described, including gains of 1q, 8q, and 17q and losses of 8p, 16q, and 17p. Both studies included long term clinical follow up and in depth molecular characterization of primary breast cancers; but the two studies grouped copy number alterations differently based on the combined evidence from multiple genomic platforms. The Molecular Taxonomy of Breast Cancer International Consortium (METABRIC) study described ten integrative clusters weighted more toward recurrent DNA copy number profiles [19]; and The Cancer Genome Atlas Network (TCGA) described five integrated subtypes weighted more toward RNA expression profiles [18]. In both studies, the genomic copy number groups shared similar common alterations among subtypes, suggesting limited specificity and perhaps limited clinical utility of genome wide copy number / LOH profiles. However, a follow up study from the METABRIC dataset has confirmed several important prognostic biomarkers in the copy number landscape: amplicons on 8p12, 8q24, 11q13, and 17q23 [70]. All are associated with an increased risk of late relapse (recurrence of disease up to decades after the initial diagnosis) in women diagnosed with ER positive, HER2 negative breast cancers [70]. Most individual copy number alterations of clinical significance in breast cancer are adverse prognostic indicators; these include amplification of 8p11.2 [71], 8q24 [57,58], 17q23 [72–74], and 20q [75,76], and deletion or loss of heterozygosity (LOH) of 10q23 (including *PTEN*) and 17p (including *TP53*). Recently, amplification of a genomic region on

9p24.1 including *JAK2*, *CD274* (PD-L1 gene), and *PDCD1LG2* (PD-L2 gene) has been identified in 1–2% of breast cancers [77]. 9p24.1 amplification is more common in triple negative breast cancers and has been associated with responsiveness to immunotherapy and JAK2 inhibition in this tumor type [77]. In addition to providing prognostic and predictive information, genomic copy number profiling can also identify recurrent patterns associated with histologic subtypes of breast cancer. Simple or nearly normal copy number profiles are characteristic of tubular carcinoma, adenoid cystic carcinoma, mucinous carcinoma, and carcinoma with neuroendocrine features [78]. Very complex copy number profiles with “firestorm” or “sawtooth” patterns are typical of medullary carcinoma, metaplastic carcinoma, and invasive micropapillary carcinoma [78]. Table 4 summarizes the recurrent clinically relevant copy number changes and regions of LOH seen in breast cancer.

Comprehensive genomic testing modalities for somatic variant detection can identify copy number changes in genes associated with increased hereditary breast cancer risk [79]. Therefore, it is important to evaluate for the possibility of a germline alteration when a deletion or other potentially inactivating structural rearrangement is detected in one of these genes (Table 5).

### Sequence alterations

Pathogenic sequence alterations in DNA samples from primary breast cancers occur most commonly in *PIK3CA*, *GATA3*, and *TP53*; other genes are individually mutated in less than 10% of breast cancers overall [18]. Genes commonly affected by copy number alterations or LOH in breast cancer can alternatively harbor point mutations or other forms of sequence variation (including small insertion-deletion mutations). Certain genes are correlated with histologic or molecular subtypes of breast cancer (for example, *CDH1* mutations in lobular carcinomas [4,6,80,81]). Inactivating mutations in genes associated with hereditary breast and ovarian cancer syndromes should prompt consideration of a germline mutation, and genetic counseling can be very helpful in this context for guiding follow up constitutional genetic testing for the patient and potentially affected family members [82,83]. Germline mutations associated with a hereditary risk for breast cancer are generally autosomal dominant with reduced penetrance; thus, not all individuals carrying a pathogenic germline mutation will develop cancer. The potential for a hereditary risk varies based on the specific alteration. *TP53* mutations are among the most common sequence alterations in breast cancer and are usually somatic rather than inherited [84]. Mutations in *BRCA1* or *BRCA2*, however, are fairly uncommon, and so these should generally raise clinical suspicion of a hereditary cancer predisposition (Table 5). Often, the initial mutational event in a tumor suppressor gene (either germline or somatic) is a sequence alteration, and the “second hit” causing loss of tumor suppressor function is a large deletion or LOH event [79].



**Table 4**  
Clinically significant copy number alterations in breast cancer.

Alteration	Relevant Gene(s)	CGC Evidence Level	Subgroup Association(s)
1q gain	unknown	2	Most common copy number alteration, often with 16q loss; all subtypes
8p11.2 amplification	<i>FGFR1, ZNF703</i>	2	METABRIC IntClust6, ER positive
8q24 amplification	<i>MYC</i>	2	METABRIC IntClust9, ER positive
9p24 amplification	<i>JAK2, CD274, PDCD1LG2</i>	2	Enriched in TNBC
10q23.3 loss or LOH	<i>PTEN</i>	2	Enriched in TNBC and in lobular carcinoma
11q13-q14 gain / amplification	<i>CCND1, EMS1, and others</i>	2	METABRIC IntClust2
16q loss / LOH	<i>CDH1</i>	2	METABRIC IntClust2, ER positive
17p loss / LOH	<i>TP53</i>	2	TNBC, basal-like intrinsic subtype
17q12 amplification	<i>ERBB2 (HER2)</i>	1	METABRIC IntClust5, HER2-enriched
17q21 amplification	<i>TOP2A</i>	2	METABRIC IntClust5, HER2-enriched
17q23 amplification (“17q distal amplicon”)	<i>RPS6KB, others</i>	2	METABRIC IntClust1
19q12	<i>CCNE1</i>	2	METABRIC IntClust5; HER2-enriched
20q gain; 20q13 amp	<i>AURKA, GNAS, ZNF217</i>	2	METABRIC IntClust1, ER Positive

**Table 5**  
Genes with known hereditary risk associations in breast cancer.

Gene	Associated Syndrome; Breast Cancer Subtype
<i>ATM</i>	Ataxia telangiectasia syndrome
<i>BARD1</i>	TNBC
<i>BRCA1</i>	BRCA-Related Breast/ Ovarian Cancer Syndrome; TNBC
<i>BRCA2</i>	BRCA-Related Breast/ Ovarian Cancer Syndrome; TNBC
<i>CDH1</i>	Hereditary Diffuse Gastric Cancer and Lobular Breast Cancer
<i>CHEK2</i>	Inherited breast cancer
<i>NBN</i>	Nijmegen Breakage Syndrome
<i>NF1</i>	Neurofibromatosis type 1
<i>PALB2</i>	Fanconi anemia
<i>PTEN</i>	Cowden syndrome
<i>RAD51C</i>	TNBC
<i>RAD51D</i>	TNBC
<i>STK11</i>	Peutz-Jeghers syndrome
<i>TP53</i>	Li-Fraumeni syndrome

Somatic genomic testing based on DNA sequencing, such as large NGS cancer panels, is not usually performed for surgically resectable breast cancer. In the setting of unresectable, metastatic, or recurrent disease, however, genomic sequencing tests can be very useful for identifying targetable abnormalities. A recent study demonstrated that *PIK3CA* mutations respond to targeted therapy in advanced hormone receptor positive, HER2 negative breast cancers [85]. Large sequencing panels can identify gene mutations that are individually rare but potentially targetable, including *ERBB2* and other ERBB family members (*EGFR*, *ERBB3*, and *ERBB4*) [25,86,87], and other kinase genes.

Sequence alterations associated with therapy resistance commonly emerge in the setting of residual or recurrent disease. *ESR1* mutations are frequent in breast cancer patients treated with hormone therapy [22,88] and can be detected in circulating cell free DNA, also called “liquid biopsy” [89–92]. Liquid biopsy is increasingly used in the clinical setting to detect subclonal resistance mutations in *ESR1*. Likewise, resistance mutations in ERBB gene family members (*EGFR*, *ERBB2/HER2*, *ERBB3*, *ERBB4*) can occur after exposure to HER2-targeted therapy [93]; *HER2* kinase domain mutations are a recurrent abnormality in this setting [94]. A list of genes associated with clinically significant sequence alterations in breast cancer is shown in Table 6.

### Gene fusions and oncogene activation by structural rearrangements

Recurrent gene fusions in cancer include the *ETV6-NTRK3* rearrangement associated with secretory breast cancer [1,2] and *MYB-NFIB* rearrangement associated with adenoid cystic carcinoma [3].

These along with another rare fusion, *MECT1-MAML2* [95], are associated with origin in adnexal structures of the breast that phenotypically resemble salivary gland epithelium.

Rare rearrangements involving kinase genes are highly targetable. The *MAGI3-AKT3* fusion is a recurrent event that is targetable by AKT small molecule inhibitors [86]. Fusions involving *RET*, *FGFR1*, and *FGFR2* have also been identified in breast cancer and are highly targetable [96–98]. *NTRK* inhibitors are FDA approved for advanced or metastatic tumors harboring *NTRK* gene fusions regardless of tumor type. Routine genomic profiling including fusion detection in breast cancer is likely to identify more targetable rearrangements in the future.

While most clinically important rearrangements are activating, rearrangements that impair or inactivate genes can also be clinically significant. Rearrangements involving the ligand binding domain of *ESR1* have been identified in hormone resistant breast cancers, some of which also harbored *ESR1* resistance mutations [21]. Functionally inactivating rearrangements of tumor suppressor genes are a recurrent abnormality in all cancers, and these can occasionally represent a germline alteration.

Other rearrangements appear frequently in breast cancer but have no definitive clinical significance. Rearrangements of neuregulin 1 (*NRG1*) are a recurrent finding in breast cancers [99,100] and are frequently found in cases with 8p11-p12 amplification [71]. *NRG1* fusions have been associated with an oncogenic transcriptional isoform called gamma-hereregulin [101], but the clinical significance of *NRG1* rearrangement remains for the most part unclear aside from its possible use as a marker of poor prognosis [102].

### Genomic signatures of defective DNA repair

Genomic signatures are hallmark patterns of mutations that are not specific to a particular genomic region but rather are observed anywhere in the genome. Several different genomic signatures have been identified in breast and other cancers. Chromothripsis, or “chromosome shattering,” can occur during cancer development, leaving a signature pattern of alternating gains and losses (typically alternating between two or maybe three different copy states) in the affected chromosome regions [103]. Chromothripsis is theorized to result from damage occurring when one or more chromosomes or chromosome regions are retained within micronuclei [103]. Another pattern of DNA damage is kataegis, or a “thunderstorm” composed of clustered mutations in a region of the genome. Kataegis has been attributed to abnormal activity of the apolipoprotein B mRNA editing enzyme catalytic polypeptide-like (APOBEC) family of cytidine deaminases and is characterized by C>T or C>G substitutions, typically within TpCpN trinucleotides

**Table 6**  
Major clinically significant genes associated with somatic sequence alterations in breast cancer.

Gene(s)	CGC Evidence Level	Clinical Significance and Subgroup Association(s)	Therapy Implication(s)
<i>AKT1</i>	2	Metastatic BC	AKT inhibitors
<i>ATM</i>	1	Possible hereditary risk; TNBC	PARP inhibitors (germline)
<i>BRCA1, BRCA2</i>	1	Often hereditary risk; TNBC	Platinum based therapy; PARP inhibitors (germline)
<i>CBFB</i>	2	ER-positive, Metastatic BC	
<i>CCND1, CCNE1*</i>	2	HER2-enriched	CDK4/6 inhibitors
<i>CDK4, CDK6*</i>	2	ER-positive, Metastatic BC	CDK4/6 inhibitors
<i>CDH1</i>	1	Lobular histology; Possible hereditary risk	
<i>CDKN2A</i>	2	Metastatic BC	
<i>CHEK2</i>	1	Often hereditary risk	PARP inhibitors (germline)
<i>ERBB2*</i>	1	Rare activating sequence alterations; kinase domain resistance mutations; HER2-enriched	HER2-targeted therapy
<i>ESR1</i>	1	Metastatic ER-positive	Hormone therapy resistance
<i>FGFR1-4</i>	2	ER-positive	FGFR inhibitors
<i>FOXA1</i>	2	ER-positive, Luminal subtype, lobular histology	
<i>GATA3</i>	2	ER-positive, Luminal subtype	
<i>JAK2*</i>	2	TNBC	JAK2 inhibitors, immunotherapy
<i>MAP2K4</i>	2	Metastatic BC	
<i>MAP3K1</i>	2	ER-positive, Metastatic BC	
<i>MYC*</i>	2		
<i>NBN</i>	1	Possible hereditary risk	PARP inhibitors (germline)
<i>NF1</i>	1	Possible hereditary risk	mTOR/PI3K/AKT inhibitors (germline)
<i>NTRK1-3</i>	1		NTRK inhibitors
<i>PALB2</i>	1	Often hereditary risk	PARP inhibitors (germline)
<i>PIK3CA</i>	1	ER-Positive, Luminal subtype	PI3K inhibitors for selected hotspot mutations; acquired hormone resistance
<i>PTEN</i>	2	Loss in lobular BC; Possible hereditary risk	mTOR/PI3K/AKT inhibitors; radiation contraindicated
<i>RB1</i>	2	Metastatic BC	Acquired hormone resistance
<i>STK11</i>	1	Possible hereditary risk	
<i>TBX3</i>	2	Lobular BC	
<i>TOP2A*</i>	2		Anthracycline inhibitors
<i>TP53</i>	1	TNBC, HER2-enriched, Metastatic BC Possible hereditary risk	Radiation contraindicated

\* Indicates genes more commonly activated by amplification than by sequence variation BC, breast cancer. TNBC, triple negative breast cancer.

[104]. Several recurrent patterns of genomic mutation, termed “Signature 2” and “Signature 13,” are characterized by kataegis related to APOBEC dysfunction in breast cancer [104]. Another genomic signature has been termed a “firestorm,” and like chromothripsis, this signature affects a localized area of the genome. Unlike chromothripsis, a firestorm shows very high level amplification (up to 30 copies or more) of a very narrow genomic region that is copied many times over in a particular region of the genome [105]. A firestorm is thought to result from cyclic breakage-fusion-bridge DNA repair events occurring early in cancer development [105,106]. The firestorm pattern of DNA amplification is seen most commonly in Luminal B and HER2 enriched breast cancers and has been associated with a poor prognosis [105,107]. The genes *ERBB2* (*HER2*) and *CCND1* are frequently amplified in these so called “firestorms” on single nucleotide polymorphism (SNP) array testing.

Some genomic signatures are currently used as biomarkers to identify patients for certain targeted therapies. *BRCA1/2*-deficient tumors can show a characteristic pattern of DNA damage called “Signature 3” [104,108]. On SNP array, *BRCA1/2*-deficient tumors frequently show a pattern of numerous large scale transitions in copy number and allelic imbalance or loss of heterozygosity (LOH), the latter including telomeric allelic imbalance, throughout the

genome [109–111]. These *BRCA1/2*-deficient patterns result from deficient homologous recombination at double strand DNA breaks, resulting in abnormal non-homologous end joining. Termed homologous recombination deficiency or HRD, this genomic signature (also referred to as a “sawtooth pattern” [105] or “genomic scar” [110,111]) is a useful predictor of responsiveness to PARP inhibitors and platinum based chemotherapy agents (cisplatin, carboplatin) [110–112]. The same signature, when observed in triple negative breast cancers lacking a detectable pathogenic germline alteration in *BRCA1* or *BRCA2*, likewise appears to be a useful predictor of responsiveness to PARP inhibitors and platinum based chemotherapy [112–115]. Another related pattern predicting responsiveness to platinum based chemotherapy is the tandem duplicator phenotype [116]. This pattern of duplications is found in triple negative breast cancers with loss of both *BRCA1* and *TP53* function and is thought to be a mechanism by which such tumors acquire pathogenic alterations in tumor suppressor genes and oncogenes [116]. Sequencing shows microhomologies at the breakpoints of tandem duplications attributable to microhomology-mediated breakage induced repair. There is a bimodal size distribution of these tandem duplications with peaks at ~10 kb and ~150 kb [116]. These tandem duplications are challenging to detect with currently available SNP array

platforms that are typically designed to detect copy number alterations on a scale of 200 kb or larger [117].

Some patterns of mutation are infrequent in breast cancer but are highly significant if identified, due to the availability of targeted therapy. Mismatch repair deficiency (MMRd) or microsatellite instability (MSI), is associated with “Signature 6” [104] and a high tumor mutation burden (TMB). Breast cancers with MMRd/MSI-high status are fairly rare and typically involve somatic rather than germline mutations in MMR genes, although germline mutations have also been observed in breast cancer [118]. Due to tumor-agnostic approval by the FDA for this class of molecular alteration, these breast cancers are targetable with the immune checkpoint inhibitor pembrolizumab (Keytruda). New markers predicting responsiveness to immunotherapy are emerging, including an amplicon on 9p24.1 containing *JAK2* and the genes encoding PD-L1 and PD-L2; preliminary evidence suggests that this is an important biomarker for responsiveness to immunotherapy in a small fraction (1–2%) of breast cancers, especially triple negative breast cancers [77]. Outside of genomic profiling for advanced or metastatic breast cancers, measurement of tumor infiltrating lymphocytes (TILs) and immunohistochemical evaluation of PD-L1 expression are clinically used to identify candidates for immunotherapy.

### Emerging genomic biomarkers: miRNAs, methylation, and beyond

Large genomic studies have shown differential expression of miRNAs that are recurrent in breast cancer [18]. Certain miRNA expression patterns have been associated with *TP53* mutation [18] or *PTEN* mutation [119] and are associated with poor prognosis. Likewise, global patterns of hypermethylation or hypomethylation have been identified in breast cancer [18]. The hypermethylated pattern affects CpG islands in gene promoter regions and has been associated with the Luminal B expression subtype; the hypomethylated pattern is seen primarily in the basal-like expression subtype [120]. Overall patterns observed on methylation profiling have also provided insights on breast cancer development [121]. In addition to global methylation patterns, hypermethylation is an important mechanism for inactivation of tumor suppressor genes, including *BRCA1* [114].

### Conclusions

A major reason that genomic profiling has not been widely adopted in the diagnostic workup of breast cancer is that these assays have not been completely standardized to permit platform-agnostic correlation with patient outcomes. Furthermore, genomic assays are more challenging to interpret in clinical specimens containing not only invasive tumor but also non-tumor cells and non-invasive (in situ) cancer precursors; in contrast, clinically available IHC and ISH assays allow cell by cell interpretation of abnormalities in selected cell populations. Furthermore, both polyploidy and polyclonality can confound the interpretation of genomic abnormalities, including the degree of copy number and allelic imbalances. Despite these substantial challenges, many genomic RNA expression assays are routinely employed to aid clinical decision making, and DNA-based genomic assays are routinely used in patients with recurrent, progressive, non-resectable, or metastatic disease. Adoption of routine genomic testing for all invasive breast cancers will require standardization across platforms and professional guidelines for application and interpretation of genomic sequence and copy number data. Given the increasing availability of therapies based on genomic signatures and rare but highly targetable genetic abnormalities, the clinical utility of genomic profiling for breast cancer is evident. Routine use of genomic testing

may facilitate patient management and contain costs by personalizing breast cancer treatment. The immediate task for clinical laboratories is the development of technical standards to guide broader use of genomic profiling in breast cancer.

### Declaration of Competing Interest

None.

### Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.cancergen.2020.02.002](https://doi.org/10.1016/j.cancergen.2020.02.002).

### CRediT authorship contribution statement

**Katherine B. Geiersbach:** Conceptualization, Methodology, Investigation, Resources, Writing - original draft, Writing - review & editing. **Hui Chen:** Methodology, Investigation, Writing - review & editing. **Rajyasree Emmadi:** Methodology, Investigation, Writing - review & editing. **Gloria T. Haskell:** Methodology, Investigation, Writing - original draft, Writing - review & editing. **Xinyan Lu:** Methodology, Investigation, Writing - review & editing. **Yajuan J. Liu:** Methodology, Investigation, Writing - review & editing. **Karen Swisshelm:** Methodology, Investigation, Writing - review & editing.

### References

- [1] Makretsov N, He M, Hayes M, Chia S, Horsman DE, Sorensen PH, Huntsman DG. A fluorescence in situ hybridization study of ETV6-NTRK3 fusion gene in secretory breast carcinoma. *Genes Chromosomes Cancer* 2004;40:152–7.
- [2] Tognon C, Knezevich SR, Huntsman D, Roskelley CD, Melnyk N, Mathers JA, et al. Expression of the ETV6-NTRK3 gene fusion as a primary event in human secretory breast carcinoma. *Cancer Cell* 2002;2:367–76.
- [3] Persson M, Andren Y, Mark J, Horlings HM, Persson F, Stenman G. Recurrent fusion of MYB and NFIB transcription factor genes in carcinomas of the breast and head and neck. *Proc Natl Acad Sci U S A* 2009;106:18740–4.
- [4] Berx G, Cleton-Janssen AM, Strumane K, de Leeuw WJ, Nollet F, van Roy F, et al. E-cadherin is inactivated in a majority of invasive human lobular breast cancers by truncation mutations throughout its extracellular domain. *Oncogene* 1996;13:1919–25.
- [5] Ciriello G, Gatza ML, Beck AH, Wilkerson MD, Rhie SK, Pastore A, et al. Comprehensive molecular portraits of invasive lobular breast cancer. *Cell* 2015;163:506–19.
- [6] Keller G, Vogelsang H, Becker I, Hutter J, Ott K, Candidus S, Grundei T, Becker KF, Mueller J, Siewert JR, Hofler H. Diffuse type gastric and lobular breast carcinoma in a familial gastric cancer patient with an E-cadherin germline mutation. *Am J Pathol* 1999;155:337–42.
- [7] Pereira B, Chin SF, Rueda OM, Volland HK, Provenzano E, Bardwell HA, et al. The somatic mutation profiles of 2,433 breast cancers refine their genomic and transcriptomic landscapes. *Nat Commun* 2016;7:11479.
- [8] Desmedt C, Zoppoli G, Gundem G, Pruneri G, Larsimont D, Fornili M, Fumagalli D, Brown D, Rothe F, Vincent D, Kheddoumi N, Rouas G, Majaj S, Brohee S, Van Loo P, Maisonneuve P, Salgado R, Van Brussel T, Lambrechts D, Bose R, Metzger O, Galant C, Bertucci F, Piccart-Gebhart M, Viale G, Biganzoli E, Campbell PJ, Sotiriou C. Genomic characterization of primary invasive lobular breast cancer. *J Clin Oncol* 2016;34:1872–81.
- [9] Kallioniemi OP, Kallioniemi A, Kurisu W, Thor A, Chen LC, Smith HS, Waldman FM, Pinkel D, Gray JW. ERBB2 amplification in breast cancer analyzed by fluorescence in situ hybridization. *Proc Natl Acad Sci U S A* 1992;89:5321–5325.
- [10] Allison KH, Hammond MEH, Dowsett M, McKernin SE, Carey LA, Fitzgibbons PL, Hayes DF, Lakhani SR, Chavez-MacGregor M, Perlmutter J, Perou CM, Regan MM, Rimm DL, Symmans WF, Torlakovic EE, Varella L, Viale G, Weisberg TF, McShane LM, Wolff AC. Estrogen and progesterone receptor testing in breast cancer: asco/cap guideline update. *J Clin Oncol* 2020 JCO1902309.
- [11] Wolff AC, Hammond MEH, Allison KH, Harvey BE, Mangu PB, Bartlett JMS, et al. Human epidermal growth factor receptor 2 testing in breast cancer: american society of clinical oncology/college of american pathologists clinical practice guideline focused update. *J Clin Oncol* 2018;36:2105–2122.
- [12] Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, Pietenpol JA. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 2011;121:2750–67.

- [13] Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature* 2000;406:747–52.
- [14] Wallden B, Storhoff J, Nielsen T, Dowidar N, Schaper C, Ferree S, Liu S, Leung S, Geiss G, Snider J, Vickery T, Davies SR, Mardis ER, Gnant M, Sestak I, Ellis MJ, Perou CM, Bernard PS, Parker JS. Development and verification of the PAM50-based prosigna breast cancer gene signature assay. *BMC Med Genomics* 2015;8:54.
- [15] Herschkowitz JI, Simin K, Weigman VJ, Mikaelian I, Usary J, Hu Z, Rasmussen KE, Jones LP, Assefnia S, Chandrasekharan S, Backlund MG, Yin Y, Khramtsov AI, Bastein R, Quackenbush J, Glazer RI, Brown PH, Green JE, Kopelovich L, Furth PA, Palazzo JP, Olopade OI, Bernard PS, Churchill GA, Van Dyke T, Perou CM. Identification of conserved gene expression features between murine mammary carcinoma models and human breast tumors. *Genome Biol* 2007;8:R76.
- [16] Prat A, Pineda E, Adamo B, Galvan P, Fernandez A, Gaba L, Diez M, Viladot M, Arance A, Munoz M. Clinical implications of the intrinsic molecular subtypes of breast cancer. *Breast* 2015;24(Suppl 2):S26–35.
- [17] Dias K, Dvorkin-Gheva A, Hallett RM, Wu Y, Hassell J, Pond GR, et al. Claudin-Low breast cancer; clinical & pathological characteristics. *PLoS ONE* 2017;12:e0168669.
- [18] Cancer Genome Atlas N. Comprehensive molecular portraits of human breast tumours. *Nature* 2012;490:61–70.
- [19] Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature* 2012;486:346–52.
- [20] Collaborative G, Davies C, Godwin J, Gray R, Clarke M, Cutter D, Darby S, McGale P, Pan HC, Taylor C, Wang YC, Dowsett M, Ingle J, Peto REarly Breast Cancer Trialists'. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. *Lancet* 2011;378:771–84.
- [21] Hartmaier RJ, Trabucco SE, Friedigkeit N, Chung JH, Parachoniak CA, Vanden Borre P, Morley S, Rosenzweig M, Gay LM, Goldberg ME, Suh J, Ali SM, Ross J, Leyland-Jones B, Young B, Williams C, Park B, Tsai M, Haley B, Peguero J, Callahan RD, Sachelarie I, Cho J, Atkinson JM, Bahreini A, Nagle AM, Puhalla SL, Watters RJ, Erdogan-Yildirim Z, Cao L, Oesterreich S, Mathew A, Lucas PC, Davidson NE, Brufsky AM, Frampton GM, Stephens PJ, Chmielecki J, Lee AV. Recurrent hyperactive ESR1 fusion proteins in endocrine therapy-resistant breast cancer. *Ann Oncol* 2018;29:872–80.
- [22] Robinson DR, Wu YM, Vats P, Su F, Lonigro RJ, Cao X, Kalyana-Sundaram S, Wang R, Ning Y, Hodges L, Gursky A, Siddiqui J, Tomlins SA, Roychowdhury S, Pienta KJ, Kim SY, Roberts JS, Rae JM, Van Poznak SH, Hayes DF, Chugh R, Kunju LP, Talpaz M, Schott AF, Chinnaiyan AM. Activating ESR1 mutations in hormone-resistant metastatic breast cancer. *Nat Genet* 2013;45:1446–51.
- [23] Jeselsohn R, Buchwalter G, De Angelis C, Brown M, Schiff R. ESR1 mutations—a mechanism for acquired endocrine resistance in breast cancer. *Nat Rev Clin Oncol* 2015;12:573–83.
- [24] Groenendijk FH, Zwart W, Floore A, Akbari S, Bernards R. Estrogen receptor splice variants as a potential source of false-positive estrogen receptor status in breast cancer diagnostics. *Breast Cancer Res Treat* 2013;140:475–84.
- [25] Bose R, Kavuri SM, Searleman AC, Shen W, Shen D, Koboldt DC, Monsey J, Goel N, Aronson AB, Li S, Ma CX, Ding L, Mardis ER, Ellis MJ. Activating HER2 mutations in HER2 gene amplification negative breast cancer. *Cancer Discov* 2013;3:224–37.
- [26] Sparano JA, Gray RJ, Makower DF, Pritchard KI, Albain KS, Hayes DF, Geyer CE Jr, Dees EC, Goetz MP, Olson JA Jr, Lively T, Badve SS, Saphner TJ, Wagner LJ, Whelan TJ, Ellis MJ, Paik S, Wood WC, Ravdin PM, Keane MM, Gomez Moreno HL, Reddy PS, Goggins TF, Mayer JA, Brufsky AM, Toppmeyer DL, Kaklamani VG, Berenberg JL, Abrams J, Sledge GW Jr. Adjuvant chemotherapy guided by a 21-Gene expression assay in breast Cancer. *N Engl J Med* 2018;379:111–21.
- [27] Sparano JA, Paik S. Development of the 21-gene assay and its application in clinical practice and clinical trials. *J Clin Oncol* 2008;26:721–8.
- [28] Cardoso F, Van't Veer L, Rutgers E, Loi S, Mook S, Piccart-Gebhart MJ. Clinical application of the 70-gene profile: the mindact trial. *J Clin Oncol* 2008;26:729–35.
- [29] Cardoso F, van't Veer LJ, Bogaerts J, Slaets L, Viale G, Delalogue S, Pierga JY, Brain E, Causeret S, DeLorenzi M, Glas AM, Goulinopoulos V, Goulioti T, Knox S, Matos E, Meulemans B, Neijenhuis PA, Nitz U, Passalacqua R, Ravdin P, Rubio IT, Saghatian M, Smilde TJ, Sotiriou C, Stork L, Straehle C, Thomas G, Thompson AM, van der Hoeven JM, Vuylsteke P, Bernards R, Tryfonidis K, Rutgers E, Piccart M. 70-Gene Signature as an Aid to treatment decisions in early-stage breast cancer. *N Engl J Med* 2016;375:717–29.
- [30] Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, Baehner FL, Walker MG, Watson D, Park T, Hiller W, Fisher ER, Wickerham DL, Bryant J, Wolmark N. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 2004;351:2817–26.
- [31] Paik S, Tang G, Shak S, Kim C, Baker J, Kim W, Cronin M, Baehner FL, Watson D, Bryant J, Costantino JP, Geyer CE Jr, Wickerham DL, Wolmark N. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol* 2006;24:3726–34.
- [32] Woodward WA, Barlow WE, Jagsi R, Buchholz TA, Shak S, Baehner F, Whelan TJ, Davidson NE, Ingle JN, King TA, Ravdin PM, Osborne CK, Tripathy D, Livingston RB, Gralow JR, Hortobagyi GN, Hayes DF, Albain KS. Association between 21-Gene assay recurrence score and locoregional recurrence rates in patients with node-positive breast cancer. *JAMA Oncol* 2020.
- [33] Albain KS, Barlow WE, Shak S, Hortobagyi GN, Livingston RB, Yeh IT, Ravdin P, Bugarini R, Baehner FL, Davidson NE, Sledge GW, Winer EP, Hudis C, Ingle JN, Perez EA, Pritchard KI, Shepherd L, Gralow JR, Yoshizawa C, Allred DC, Osborne CK, Hayes DF. Breast cancer intergroup of north A. prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol* 2010;11:55–65.
- [34] Buus R, Sestak I, Kronenwett R, Denkert C, Dubsky P, Krappmann K, Scheer M, Petry C, Cuzick J, Dowsett M. Comparison of endopredict and EPclin with oncotype dx recurrence score for prediction of risk of distant recurrence after endocrine therapy. *J Natl Cancer Inst* 2016;108.
- [35] Filipits M, Rudas M, Jakesz R, Dubsky P, Fitzal F, Singer CF, Dietze O, Greil R, Jelen A, Sevela P, Freibauer C, Muller V, Janicek F, Schmidt M, Kolbl H, Rody A, Kaufmann M, Schroth W, Brauch H, Schwab M, Fritz P, Weber KE, Feder IS, Hennig G, Kronenwett R, Gehrmann M, Gnant M, Investigators EP. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin Cancer Res* 2011;17:6012–20.
- [36] Sgroi DC, Sestak I, Cuzick J, Zhang Y, Schnabel CA, Schroeder B, Erlander MG, Dunbier A, Sidhu K, Lopez-Knowles E, Goss PE, Dowsett M. Prediction of late distant recurrence in patients with oestrogen-receptor-positive breast cancer: a prospective comparison of the breast-cancer index (BCI) assay, 21-gene recurrence score, and IHC4 in the transatac study population. *Lancet Oncol* 2013;14:1067–76.
- [37] Sgroi DC, Carney E, Zarrella E, Steffel L, Binns SN, Finkelstein DM, Szymonifka J, Bhan AK, Shepherd LE, Zhang Y, Schnabel CA, Erlander MG, Ingle JN, Porter P, Muss HB, Pritchard KI, Tu D, Rimm DL, Goss PE. Prediction of late disease recurrence and extended adjuvant letrozole benefit by the hoxb13/il17br biomarker. *J Natl Cancer Inst* 2013;105:1036–42.
- [38] Zhang Y, Schnabel CA, Schroeder BE, Jerevall PL, Jankowitz RC, Fornander T, Stal O, Brufsky AM, Sgroi D, Erlander MG. Breast cancer index identifies early-stage estrogen receptor-positive breast cancer patients at risk for early- and late-distant recurrence. *Clin Cancer Res* 2013;19:4196–205.
- [39] Sanft T, Aktas B, Schroeder B, Bossuyt V, DiGiovanna M, Abu-Khalaf M, Chung G, Silber A, Hofstadter E, Mougalian S, Epstein L, Hatzis C, Schnabel C, Pusztai L. Prospective assessment of the decision-making impact of the breast cancer index in recommending extended adjuvant endocrine therapy for patients with early-stage ER-positive breast cancer. *Breast Cancer Res Treat* 2015;154:533–41.
- [40] Bartlett JM, Christiansen J, Gustavson M, Rimm DL, Piper T, van de Velde CJ, Hasenburger A, Kieback DG, Putter H, Markopoulos CJ, Dirix LY, Seynaeve C, Rea DW. Validation of the IHC4 breast cancer prognostic algorithm using multiple approaches on the multinational team clinical trial. *Arch Pathol Lab Med* 2016;140:66–74.
- [41] Hou Y, Zynger DL, Li X, Li Z. Comparison of oncotype dx with modified magee equation recurrence scores in low-grade invasive carcinoma of breast. *Am J Clin Pathol* 2017;148:167–72.
- [42] Bhargava R, Clark BZ, Dabbs DJ. Breast cancers with magee equation score of less than 18, or 18–25 and mitosis score of 1, do not require oncotype dx testing: a value study. *Am J Clin Pathol* 2019;151:316–23.
- [43] Harbeck N, Schmitt M, Meisner C, Friedel C, Untch M, Schmidt M, Sweep CG, Lisboa BW, Lux MP, Beck T, Hasmuller S, Kiechle M, Janicek F, Thomssen C, Chemo NSG. Ten-year analysis of the prospective multicentre chemo-n0 trial validates american society of clinical oncology (ASCO)-recommended biomarkers uPA and PAI-1 for therapy decision making in node-negative breast cancer patients. *Eur J Cancer* 2013;49:1825–35.
- [44] Dabbs DJ, Klein ME, Mohsin SK, Tubbs RR, Shuai Y, Bhargava R. High false-negative rate of HER2 quantitative reverse transcription polymerase chain reaction of the oncotype dx test: an independent quality assurance study. *J Clin Oncol* 2011;29:4279–85.
- [45] Dressler LG, Seamer LC, Owens MA, Clark GM, McGuire WL. DNA flow cytometry and prognostic factors in 1331 frozen breast cancer specimens. *Cancer* 1988;61:420–7.
- [46] Wenger CR, Beardslee S, Owens MA, Pounds G, Oldaker T, Vendely P, et al. DNA ploidy, S-phase, and steroid receptors in more than 127,000 breast cancer patients. *Breast Cancer Res. Treat.* 1993;28:9–20.
- [47] Carpenter R, Gibbs N, Matthews J, Cooke T. Importance of cellular dna content in pre-malignant breast disease and pre-invasive carcinoma of the female breast. *Br J Surg* 1987;74:905–6.
- [48] Teixeira MR, Pandis N, Bardi G, Andersen JA, Mandahl N, Mitelman F, et al. Cytogenetic analysis of multifocal breast carcinomas: detection of karyotypically unrelated clones as well as clonal similarities between tumour foci. *Br. J. Cancer* 1994;70:922–7.
- [49] Teixeira MR, Pandis N, Bardi G, Andersen JA, Mitelman F, Heim S. Clonal heterogeneity in breast cancer: karyotypic comparisons of multiple intra- and extra-tumorous samples from 3 patients. *International journal of cancer* 1995;63:63–8.
- [50] Steinarsdottir M, Petursdottir I, Snorraddottir S, Eyfjord JE, Ogmundsdottir HM. Cytogenetic studies of breast carcinomas: different karyotypic profiles detected by direct harvesting and short-term culture. *Genes Chromosomes Cancer* 1995;13:239–48.
- [51] Kuukasjarvi T, Karhu R, Tanner M, Kahkonen M, Schaffer A, Nupponen N, Pennanen S, Kallioniemi A, Kallioniemi OP, Isola J. Genetic heterogeneity and clonal evolution underlying development of asynchronous metastasis in human breast cancer. *Cancer Res* 1997;57:1597–604.



- [52] Going JJ, Abd El-Monem HM, Craft JA. Clonal origins of human breast cancer. *J Pathol* 2001;194:406–12.
- [53] Teixeira MR, Pandis N, Heim S. Cytogenetic clues to breast carcinogenesis. *Genes Chromosomes Cancer* 2002;33:1–16.
- [54] Schechter AL, Stern DF, Vaidyanathan L, Decker SJ, Drebin JA, Greene MI, et al. The neu oncogene: an erb-B-related gene encoding a 185,000-Mr tumour antigen. *Nature* 1984;312:513–16.
- [55] Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 1987;235:177–82.
- [56] Seshadri R, Matthews C, Dobrovic A, Horsfall DJ. The significance of oncogene amplification in primary breast cancer. *Int J Cancer* 1989;43:270–2.
- [57] Escot C, Theillet C, Lidereau R, Spyrtos F, Champeme MH, Gest J, Callahan R. Genetic alteration of the c-myc protooncogene (MYC) in human primary breast carcinomas. *Proc Natl Acad Sci U S A* 1986;83:4834–8.
- [58] Borg A, Baldetorp B, Fernö M, Olsson H, Sigurdsson H. c-myc amplification is an independent prognostic factor in postmenopausal breast cancer. *Int J Cancer* 1992;51:687–91.
- [59] Zafrani B, Gerbault-Seureau M, Mosseri V, Dutrillaux B. Cytogenetic study of breast cancer: clinicopathological significance of homogeneously staining regions in 84 patients. *Hum Pathol* 1992;23:542–7.
- [60] Muleris M, Almeida A, Gerbault-Seureau M, Malfoy B, Dutrillaux B. Detection of DNA amplification in 17 primary breast carcinomas with homogeneously staining regions by a modified comparative genomic hybridization technique. *Genes Chromosomes Cancer* 1994;10:160–70.
- [61] Muleris M, Almeida A, Gerbault-Seureau M, Malfoy B, Dutrillaux B. Identification of amplified DNA sequences in breast cancer and their organization within homogeneously staining regions. *Genes Chromosomes Cancer* 1995;14:155–63.
- [62] Isola JJ, Kallioniemi OP, Chu LW, Fuqua SA, Hilsenbeck SG, Osborne CK, Waldman FM. Genetic aberrations detected by comparative genomic hybridization predict outcome in node-negative breast cancer. *Am J Pathol* 1995;147:905–11.
- [63] Hislop RG, Pratt N, Stocks SC, Steel CM, Sales M, Goudie D, Robertson A, Thompson AM. Karyotypic aberrations of chromosomes 16 and 17 are related to survival in patients with breast cancer. *Br J Surg* 2002;89:1581–6.
- [64] Zudaire I, Otero MD, Caballero C, Valenti C, Martinez-Penuela JM, Isola J, et al. Genomic imbalances detected by comparative genomic hybridization are prognostic markers in invasive ductal breast carcinomas. *Histopathology* 2002;40:547–55.
- [65] Aubele M, Auer G, Braselmann H, Nahrig J, Zitzelsberger H, Quintanilla-Martinez L, Smida J, Walch A, Hofler H, Werner M. Chromosomal imbalances are associated with metastasis-free survival in breast cancer patients. *Anal Cell Pathol* 2002;24:77–87.
- [66] Deltas A, Torhorst J, Schultheiss E, Mihatsch MJ, Moch H. DNA sequence losses on chromosomes 11p and 18q are associated with clinical outcome in lymph node-negative ductal breast cancer. *Clin Cancer Res* 2002;8:1210–1216.
- [67] Rennstam K, Ahlstedt-Soini M, Baldetorp B, Bendahl PO, Borg A, Karhu R, Tanner M, Tirkkonen M, Isola J. Patterns of chromosomal imbalances defines subgroups of breast cancer with distinct clinical features and prognosis. a study of 305 tumors by comparative genomic hybridization. *Cancer Res* 2003;63:8861–8.
- [68] Naylor TL, Greshock J, Wang Y, Colligon T, Yu QC, Clemmer V, Zaks TZ, Weber BL. High resolution genomic analysis of sporadic breast cancer using array-based comparative genomic hybridization. *Breast Cancer Res* 2005;7:R1186–98.
- [69] Pollack JR, Sorlie T, Perou CM, Rees CA, Jeffrey SS, Lønning PE, Tibshirani R, Botstein D, Borresen-Dale AL, Brown PO. Microarray analysis reveals a major direct role of DNA copy number alteration in the transcriptional program of human breast tumors. *Proc Natl Acad Sci U S A* 2002;99:12963–8.
- [70] Rueda OM, Sammut SJ, Seoane JA, Chin SF, Caswell-Jin JL, Callari M, et al. Dynamics of breast-cancer relapse reveal late-recurring ER-positive genomic subgroups. *Nature* 2019;567:399–404.
- [71] Gelsi-Boyer V, Orsetti B, Cervera N, Finetti P, Sircoulomb F, Rouge C, La-sorsa L, Letessier A, Ginestier C, Monville F, Esteyries S, Adelaide J, Esterni B, Henry C, Ethier SP, Bibeau F, Mozziconacci MJ, Charafe-Jauffret E, Jacquemier J, Bertucci F, Birnbaum D, Theillet C, Chaffanet M. Comprehensive profiling of 8p11–12 amplification in breast cancer. *Mol Cancer Res* 2005;3:655–67.
- [72] Monni O, Barlund M, Mousses S, Kononen J, Sauter G, Heiskanen M, Paavola P, Avela K, Chen Y, Bittner ML, Kallioniemi A. Comprehensive copy number and gene expression profiling of the 17q23 amplicon in human breast cancer. *Proc Natl Acad Sci U S A* 2001;98:5711–16.
- [73] Barlund M, Monni O, Kononen J, Cornelison R, Torhorst J, Sauter G, Kallioniemi O-P, Kallioniemi A. Multiple genes at 17q23 undergo amplification and overexpression in breast cancer. *Cancer Res* 2000;60:5340–4.
- [74] Parssinen J, Kuukasjarvi T, Karhu R, Kallioniemi A. High-level amplification at 17q23 leads to coordinated overexpression of multiple adjacent genes in breast cancer. *Br. J. Cancer* 2007;96:1258–64.
- [75] Collins C, Rommens JM, Kowbel D, Godfrey T, Tanner M, Hwang SI, Polikoff D, Nonet G, Cochran J, Myambo K, Jay KE, Froula J, Cloutier T, Kuo WL, Yaswen P, Dairkee S, Giovanola J, Hutchinson GB, Isola J, Kallioniemi OP, Palazzolo M, Martin C, Ericsson C, Pinkel D, Albertson D, Li WB, Gray JW. Positional cloning of ZNF217 and NABC1: genes amplified at 20q13.2 and overexpressed in breast carcinoma. *Proc Natl Acad Sci U S A* 1998;95:8703–8.
- [76] Nonet GH, Stampfer MR, Chin K, Gray JW, Collins CC, Yaswen P. The ZNF217 gene amplified in breast cancers promotes immortalization of human mammary epithelial cells. *Cancer Res* 2001;61:1250–1254.
- [77] Gupta S, Vanderbilt CM, Cotzia P, Arias-Stella JA 3rd, Chang JC, Zehir A, et al. Next-Generation sequencing-based assessment of JAK2, PD-L1, and PD-L2 copy number alterations at 9p24.1 in breast cancer: potential implications for clinical management. *J Mol Diagn* 2019;21:307–317.
- [78] Horlings HM, Weigelt B, Anderson EM, Lambros MB, Mackay A, Natrajan R, Ng CK, Geyer FC, van de Vijver MJ, Reis-Filho JS. Genomic profiling of histological special types of breast cancer. *Breast Cancer Res Treat* 2013;142:257–69.
- [79] Riaz N, Blecua P, Lim RS, Shen R, Higginson DS, Weinhold N, Norton L, Weigelt B, Powell SN, Reis-Filho JS. Pan-cancer analysis of bi-allelic alterations in homologous recombination DNA repair genes. *Nat Commun* 2017;8:857.
- [80] Roylance R, Droufakou S, Gorman P, Gillett C, Hart IR, Hanby A, Tomlinson I. The role of E-cadherin in low-grade ductal breast tumourigenesis. *J Pathol* 2003;200:53–8.
- [81] Cleton-Jansen A-M. E-cadherin and loss of heterozygosity at chromosome 16 in breast carcinogenesis: different genetic pathways in ductal and lobular breast cancer? *Breast Cancer Research* 2002;4:5–8.
- [82] Network NCC. NCCN guidelines version 3.2019: genetic/familial high-risk assessment: breast and ovarian. National Comprehensive Cancer Network, Inc.; 2019.
- [83] Shimelis H, LaDuca H, Hu C, Hart SN, Na J, Thomas A, Akinhanmi M, Moore RM, Brauch H, Cox A, Eccles DM, Ewart-Toland A, Fasching PA, Fostira F, Garber J, Godwin AK, Konstantopoulou I, Nevanlinna H, Sharma P, Yan-noukakos D, Yao S, Feng BJ, Tippin Davis B, Lilyquist J, Pesaran T, Goldgar DE, Polley EC, Dolinsky JS, Couch FJ. Triple-Negative breast cancer risk genes identified by multigene hereditary cancer panel testing. *J Natl Cancer Inst* 2018.
- [84] Coles C, Condie A, Chetty U, Steel CM, Evans HJ, Prosser J. p53 mutations in breast cancer. *Cancer Res* 1992;52:5291–8.
- [85] Andre F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS, Iwata H, Conte P, Mayer IA, Kaufman B, Yamashita T, Lu YS, Inoue K, Takahashi M, Pappai Z, Longin AS, Mills D, Wilke C, Hirawat S, Juric D, Group S-S. the S-SG. alpelisib for PIK3CA-Mutated, hormone receptor-positive advanced breast cancer. *N Engl J Med* 2019;380:1929–40.
- [86] Banerji S, Cibulskis K, Rangel-Escareno C, Brown KK, Carter SL, Frederick AM, et al. Sequence analysis of mutations and translocations across breast cancer subtypes. *Nature* 2012;486:405–9.
- [87] Hyman DM, Piha-Paul SA, Won H, Rodon J, Saura C, Shapiro GI, et al. HER kinase inhibition in patients with HER2- and HER3-mutant cancers. *Nature* 2018;554:189–94.
- [88] Lefebvre C, Bachelot T, Filleron T, Pedrero M, Campone M, Soria JC, et al. Mutational profile of metastatic breast cancers: a retrospective analysis. *PLoS Med* 2016;13:e1002201.
- [89] Fribbens C, O'Leary B, Kilburn L, Hrebien S, Garcia-Murillas I, Beaney M, Cristofanilli M, Andre F, Loi S, Loibl S, Jiang J, Bartlett CH, Koehler M, Dowsett M, Bliss JM, Johnston SR, Turner NC. Plasma ESR1 mutations and the treatment of estrogen receptor-positive advanced breast cancer. *J Clin Oncol* 2016;34:2961–8.
- [90] Clatot F, Perdrix A, Augusto L, Beaussire L, Delacour J, Calbrix C, et al. Kinetics, prognostic and predictive values of ESR1 circulating mutations in metastatic breast cancer patients progressing on aromatase inhibitor. *Oncotarget* 2016;7:74448–59.
- [91] Chu D, Paoletti C, Gersch C, VanDenBerg DA, Zabransky DJ, Cochran RL, Wong HY, Toro PV, Cidado J, Croessmann S, Erlanger B, Cravero K, Kyker-Snowman K, Button B, Parsons HA, Dalton WB, Gillani R, Medford A, Aung K, Tokudome N, Chinnaiyan AM, Schott A, Robinson D, Jacks KS, Lauring J, Hurley PJ, Hayes DF, Rae JM, Park BH. ESR1 mutations in circulating plasma tumor dna from metastatic breast cancer patients. *Clin Cancer Res* 2016;22:993–9.
- [92] Chandralapaty S, Chen D, He W, Sung P, Samoil A, You D, et al. Prevalence of ESR1 mutations in cell-free DNA and outcomes in metastatic breast cancer: a secondary analysis of the BOLERO-2 clinical trial. *JAMA Oncol* 2016;2:1310–15.
- [93] Elster N, Toomey S, Fan Y, Cremona M, Morgan C, Weiner Gorzel K, et al. Frequency, impact and a preclinical study of novel ERBB gene family mutations in HER2-positive breast cancer. *Ther Adv Med Oncol* 2018;10.
- [94] Zuo WJ, Jiang YZ, Wang YJ, Xu XE, Hu X, Liu GY, Wu J, Di GH, Yu KD, Shao ZM. Dual characteristics of novel HER2 kinase domain mutations in response to HER2-Targeted therapies in human breast cancer. *Clin Cancer Res* 2016;22:4859–69.
- [95] Kazakov DV, Vanecek T, Belousova IE, Mukensnabl P, Kollertova D, Michal M. Skin-type hidradenoma of the breast parenchyma with t(11;19) translocation: hidradenoma of the breast. *Am J Dermatopathol* 2007;29:457–461.
- [96] Paratala BS, Chung JH, Williams CB, Yilmazel B, Petrosky W, Williams K, Schrock AB, Gay LM, Lee E, Dolfi SC, Pham K, Lin S, Yao M, Kulkarni A, Di-Clemente F, Liu C, Rodriguez-Rodriguez L, Ganesan S, Ross JS, Ali SM, Leyland-Jones B, Hirshfield KM. RET rearrangements are actionable alterations in breast cancer. *Nat Commun* 2018;9:4821.

- [97] Wu YM, Su F, Kalyana-Sundaram S, Khazanov N, Ateeq B, Cao X, Lonigro RJ, Vats P, Wang R, Lin SF, Cheng AJ, Kunju LP, Siddiqui J, Tomlins SA, Wyngaard P, Sadi S, Roychowdhury S, Hussain MH, Feng FY, Zalupski MM, Talpaz M, Pienta KJ, Rhodes DR, Robinson DR, Chinnaiyan AM. Identification of targetable FGFR gene fusions in diverse cancers. *Cancer Discov* 2013;3:636–47.
- [98] Perez-Garcia J, Munoz-Couselo E, Soberino J, Racca F, Cortes J. Targeting FGFR pathway in breast cancer. *Breast* 2018;37:126–33.
- [99] Adelaide J, Huang HE, Murati A, Alsop AE, Orsetti B, Mozziconacci MJ, Popovici C, Ginestier C, Letessier A, Basset C, Courtay-Cahen C, Jacquemier J, Theillet C, Birnbaum D, Edwards PA, Chaffanet M. A recurrent chromosome translocation breakpoint in breast and pancreatic cancer cell lines targets the neuregulin/NGR1 gene. *Genes Chromosomes Cancer* 2003;37:333–45.
- [100] Huang HE, Chin SF, Ginestier C, Bardou VJ, Adelaide J, Iyer NG, Garcia MJ, Pole JC, Callagy GM, Hewitt SM, Gullick WJ, Jacquemier J, Caldas C, Chaffanet M, Birnbaum D, Edwards PA. A recurrent chromosome breakpoint in breast cancer at the NRG1/neuregulin 1/hereregulin gene. *Cancer Res* 2004;64:6840–4.
- [101] Liu X, Baker E, Eyre HJ, Sutherland GR, Zhou M. Gamma-hergulin: a fusion gene of DOC-4 and neuregulin-1 derived from a chromosome translocation. *Oncogene* 1999;18:7110–14.
- [102] Prentice LM, Shadeo A, Lestou VS, Miller MA, deLeeuw RJ, Makretsov N, et al. NRG1 gene rearrangements in clinical breast cancer: identification of an adjacent novel amplicon associated with poor prognosis. *Oncogene* 2005;24:7281–9.
- [103] Stephens PJ, Greenman CD, Fu B, Yang F, Bignell GR, Mudie LJ, et al. Massive genomic rearrangement acquired in a single catastrophic event during cancer development. *Cell* 2011;144:27–40.
- [104] Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature* 2013;500:415–21.
- [105] Hicks J, Krasnitz A, Lakshmi B, Navin NE, Riggs M, Leib E, Esposito D, Alexander J, Troge J, Grubor V, Yoon S, Wigler M, Ye K, Borresen-Dale AL, Naume B, Schlichting E, Norton L, Hagerstrom T, Skoog L, Auer G, Maner S, Lundin P, Zetterberg A. Novel patterns of genome rearrangement and their association with survival in breast cancer. *Genome Res* 2006;16:1465–79.
- [106] Gisselsson D, Pettersson L, Hoglund M, Heidenblad M, Gorunova L, Wiegant J, Mertens F, Dal Cin P, Mitelman F, Mandahl N. Chromosomal breakage-fusion-bridge events cause genetic intratumor heterogeneity. *Proc Natl Acad Sci U S A* 2000;97:5357–62.
- [107] Volland HK, Rueda OM, Chin SF, Curtis C, Turashvili G, Shah S, Lingjaerde OC, Yuan Y, Ng CK, Dunning MJ, Dicks E, Provenzano E, Sammut S, McKinney S, Ellis IO, Pinder S, Purushotham A, Murphy LC, Kristensen VN, Group M, Brenton JD, Pharoah PD, Borresen-Dale AL, Aparicio S, Caldas C. A tumor DNA complex aberration index is an independent predictor of survival in breast and ovarian cancer. *Mol Oncol* 2015;9:115–27.
- [108] Nik-Zainal S, Davies H, Staaf J, Ramakrishna M, Glodzik D, Zou X, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* 2016;534:47–54.
- [109] Timms KM, Abkevich V, Hughes E, Neff C, Reid J, Morris B, Kalva S, Potter J, Tran TV, Chen J, Iliev D, Sangale Z, Tikishvili E, Perry M, Zharkikh A, Gutin A, Lanchbury JS. Association of BRCA1/2 defects with genomic scores predictive of DNA damage repair deficiency among breast cancer subtypes. *Breast Cancer Res* 2014;16:475.
- [110] Watkins JA, Irshad S, Grigoriadis A, Tutt AN. Genomic scars as biomarkers of homologous recombination deficiency and drug response in breast and ovarian cancers. *Breast Cancer Res* 2014;16:211.
- [111] Marquard AM, Eklund AC, Joshi T, Krzystanek M, Favero F, Wang ZC, Richardson AL, Silver DP, Szallasi Z, Birkbak NJ. Pan-cancer analysis of genomic scar signatures associated with homologous recombination deficiency suggests novel indications for existing cancer drugs. *Biomark Res* 2015;3:9.
- [112] Manie E, Popova T, Battistella A, Tarabeux J, Caux-Moncoutier V, Golmard L, Smith NK, Mueller CR, Mariani O, Sigal-Zafrani B, Dubois T, Vincent-Salomon A, Houdayer C, Stoppa-Lyonnet D, Stern MH. Genomic hallmarks of homologous recombination deficiency in invasive breast carcinomas. *Int J Cancer* 2016;138:891–900.
- [113] Telli ML, Jensen KC, Vinayak S, Kurian AW, Lipson JA, Flaherty PJ, et al. Phase II study of gemcitabine, carboplatin, and iniparib as neoadjuvant therapy for triple-negative and BRCA1/2 mutation-associated breast cancer with assessment of a tumor-based measure of genomic instability: prECOG 0105. *J Clin Oncol* 2015;33:1895–901.
- [114] Davies H, Glodzik D, Morganella S, Yates LR, Staaf J, Zou X, Ramakrishna M, Martin S, Boyault S, Sieuwerts AM, Simpson PT, King TA, Raine K, Eyfjord JE, Kong G, Borg A, Birney E, Stunnenberg HG, van de Vijver MJ, Borresen-Dale AL, Martens JW, Span PN, Lakhani SR, Vincent-Salomon A, Sotiriou C, Tutt A, Thompson AM, Van Laere S, Richardson AL, Viari A, Campbell PJ, Stratton MR, Nik-Zainal S. HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on mutational signatures. *Nat Med* 2017;23:517–25.
- [115] Telli ML, Timms KM, Reid J, Hennessy B, Mills GB, Jensen KC, Szallasi Z, Barry WT, Winer EP, Tung NM, Isakoff SJ, Ryan PD, Greene-Colozzi A, Gutin A, Sangale Z, Iliev D, Neff C, Abkevich V, Jones JT, Lanchbury JS, Hartman AR, Garber JE, Ford JM, Silver DP, Richardson AL. Homologous recombination deficiency (HRD) score predicts response to platinum-containing neoadjuvant chemotherapy in patients with triple-negative breast cancer. *Clin Cancer Res* 2016;22:3764–73.
- [116] Menghi F, Inaki K, Woo X, Kumar PA, Grzeda KR, Malhotra A, Yadav V, Kim H, Marquez EJ, Ucar D, Shreckengast PT, Wagner JP, MacIntyre G, Murthy Karuturi KR, Scully R, Keck J, Chuang JH, Liu ET. The tandem duplicator phenotype as a distinct genomic configuration in cancer. *Proc Natl Acad Sci U S A* 2016;113:E2373–82.
- [117] Li MM, Monzon FA, Biegel JA, Jobanputra V, Laffin JJ, Levy B, Leon A, Miron P, Rossi MR, Toruner G, Alvarez K, Doho G, Dougherty MJ, Hu X, Kash S, Streck D, Znoyko I, Hagenkord JM, Wolff DJ. A multicenter, cross-platform clinical validation study of cancer cytogenomic arrays. *Cancer Genet* 2015;208:525–36.
- [118] Davies H, Morganella S, Purdie CA, Jang SJ, Borgen E, Russnes H, et al. Whole-Genome sequencing reveals breast cancers with mismatch repair deficiency. *Cancer Res* 2017;77:4755–62.
- [119] Wang DY, Gendoo DMA, Ben-David Y, Woodgett JR, Zacksenhaus E. A subgroup of microRNAs defines PTEN-deficient, triple-negative breast cancer patients with poorest prognosis and alterations in RB1, MYC, and WNT signaling. *Breast cancer research* 2019;21.
- [120] Stefansson OA, Moran S, Gomez A, Sayols S, Arribas-Jorba C, Sandoval J, Hilmarsson H, Olafsdottir E, Tryggvadottir L, Jonasson JG, Eyfjord J, Esteller M. A DNA methylation-based definition of biologically distinct breast cancer subtypes. *Mol Oncol* 2015;9:555–68.
- [121] Teschendorff AE, Gao Y, Jones A, Ruebner M, Beckmann MW, Wachter DL, Fasching PA, Widschwendter M. DNA methylation outliers in normal breast tissue identify field defects that are enriched in cancer. *Nat Commun* 2016;7:10478.