

Question 1

Let $\{u_1, u_2, u_3\}$ be an orthonormal set of vectors in some vector space with inner product. Let

$$u := u_1 + 2u_2 + 3u_3 \text{ and } v := u_1 - u_3$$

Compute $\langle u, v \rangle$, $\|u\|$, and $\|v\|$.

Solution: Since the basis is orthonormal, the inner product of any two vectors in the basis is 0, and the inner product of a vector in the basis with itself is 1.

$$\begin{aligned}\langle u, v \rangle &= \langle u_1 + 2u_2 + 3u_3, u_1 - u_3 \rangle \\ &= \langle u_1, u_1 - u_3 \rangle + \langle 2u_2, u_1 - u_3 \rangle + \langle 3u_3, u_1 - u_3 \rangle \\ &= \langle u_1, u_1 \rangle - \langle u_1, u_3 \rangle + \langle 2u_2, u_1 \rangle - \langle 2u_2, u_3 \rangle + \langle 3u_3, u_1 \rangle - \langle 3u_3, u_3 \rangle \\ &= 1 - 0 + 0 - 0 + 0 - 3 \\ &= -2\end{aligned}$$

$$\begin{aligned}\|u\|^2 &= \langle u, u \rangle \\ &= \langle u_1 + 2u_2 + 3u_3, u_1 + 2u_2 + 3u_3 \rangle \\ &= \langle u_1, u_1 + 2u_2 + 3u_3 \rangle + \langle 2u_2, u_1 + 2u_2 + 3u_3 \rangle + \langle 3u_3, u_1 + 2u_2 + 3u_3 \rangle \\ &= \langle u_1, u_1 \rangle + \langle 2u_2, 2u_2 \rangle + \langle 3u_3, 3u_3 \rangle \\ &= 1 + 4 + 9 \\ &= 14 \implies \|u\| = \sqrt{14}\end{aligned}$$

$$\begin{aligned}\|v\|^2 &= \langle v, v \rangle \\ &= \langle u_1 - u_3, u_1 - u_3 \rangle \\ &= \langle u_1, u_1 - u_3 \rangle + \langle -u_3, u_1 - u_3 \rangle \\ &= \langle u_1, u_1 \rangle - \langle u_1, u_3 \rangle - \langle u_3, u_1 \rangle + \langle u_3, u_3 \rangle \\ &= 1 - 0 - 0 + 1 \\ &= 2 \implies \|v\| = \sqrt{2}\end{aligned}$$

Question 2

Consider the vector space $C[-1, 1]$ equipped with the inner product:

$$\langle f, g \rangle := \int_{-1}^1 f(x)g(x)dx$$

1. Show that $1, x$ are orthogonal.
2. Compute the norms $\|1\|, \|x\|$.

Solution: Two vectors are orthogonal if their inner product is 0.

$$\langle 1, x \rangle = \int_{-1}^1 1 \cdot x \, dx = \int_{-1}^1 x \, dx = \frac{1}{2}x^2 \Big|_{-1}^1 = \frac{1}{2} - \frac{1}{2} = 0$$

Thus, $1, x$ are orthogonal. The norms of 1 and x are the square root of their inner product with themselves.

$$\begin{aligned} \|1\|^2 &= \langle 1, 1 \rangle = \int_{-1}^1 1 \cdot 1 \, dx = \int_{-1}^1 1 \, dx = x \Big|_{-1}^1 = 1 - (-1) = 2 \\ &\implies \|1\| = \sqrt{2} \\ \|x\|^2 &= \langle x, x \rangle = \int_{-1}^1 x \cdot x \, dx = \int_{-1}^1 x^2 \, dx = \frac{1}{3}x^3 \Big|_{-1}^1 = \frac{1}{3} - \frac{1}{3} = \frac{2}{3} \\ &\implies \|x\| = \sqrt{\frac{2}{3}} \end{aligned}$$

Question 3

Let

$$u_1 = \left(\frac{1}{3\sqrt{2}}, \frac{1}{3\sqrt{2}}, -\frac{4}{3\sqrt{2}} \right)^T, \quad u_2 = \frac{1}{3}(2, 2, 1)^T, \quad u_3 = \frac{1}{\sqrt{2}}(1, -1, 0)^T$$

1. Show that u_1, u_2, u_3 is an orthonormal basis for \mathbb{R}^3 .
2. Let $x = (1, 2, 2)^T$. Find the projection of p of x onto $S := \text{span}\{u_2, u_3\}$.

Solution: A set of vectors form an orthonormal basis if they are orthogonal and their norms are 1. Since we are working in \mathbb{R}^3 , we can use the dot product to check if the vectors are orthogonal.

$$\begin{aligned} \langle u_1, u_2 \rangle &= \frac{1}{3\sqrt{2}} \cdot \frac{2}{3} + \frac{1}{3\sqrt{2}} \cdot \frac{2}{3} + -\frac{4}{3\sqrt{2}} \cdot \frac{1}{3} = \frac{2}{9\sqrt{2}} + \frac{2}{9\sqrt{2}} - \frac{4}{9\sqrt{2}} = 0 \\ \langle u_1, u_3 \rangle &= \frac{1}{3\sqrt{2}} \cdot \frac{1}{\sqrt{2}} + \frac{1}{3\sqrt{2}} \cdot -\frac{1}{\sqrt{2}} + -\frac{4}{3\sqrt{2}} \cdot 0 = \frac{1}{6} - \frac{1}{6} = 0 \\ \langle u_2, u_3 \rangle &= \frac{2}{3} \cdot \frac{1}{\sqrt{2}} + \frac{2}{3} \cdot -\frac{1}{\sqrt{2}} + \frac{1}{3} \cdot 0 = \frac{2}{3\sqrt{2}} - \frac{2}{3\sqrt{2}} = 0 \end{aligned}$$

Thus, u_1, u_2, u_3 are orthogonal. To check if their norms are 1, we can use the formula $\|u\|^2 = \langle u, u \rangle$.

$$\begin{aligned} \|u_1\|^2 &= \langle u_1, u_1 \rangle = \frac{1}{3\sqrt{2}} \cdot \frac{1}{3\sqrt{2}} + \frac{1}{3\sqrt{2}} \cdot \frac{1}{3\sqrt{2}} + -\frac{4}{3\sqrt{2}} \cdot -\frac{4}{3\sqrt{2}} = \frac{1}{18} + \frac{1}{18} + \frac{16}{18} = 1 \\ \|u_2\|^2 &= \langle u_2, u_2 \rangle = \frac{1}{3} \cdot \frac{1}{3} + \frac{1}{3} \cdot \frac{1}{3} + 1 \cdot 1 = \frac{1}{9} + \frac{1}{9} + 1 = 1 \\ \|u_3\|^2 &= \langle u_3, u_3 \rangle = \frac{1}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}} + -\frac{1}{\sqrt{2}} \cdot -\frac{1}{\sqrt{2}} + 0 \cdot 0 = \frac{1}{2} + \frac{1}{2} = 1 \end{aligned}$$

Since the squares of norms are 1, the norms are 1. Thus, u_1, u_2, u_3 are orthonormal. To find the projection of x onto S ...

Question 4

Let $v_1 := (1, 2, 0, -1)^T$ $v_2 := (1, -1, 0, 0)^T$ $v_3 := (0, 1, 0, -1)^T$. Find the angle between v_1, v_2 , v_2, v_3 , and v_1, v_3 . Find the norm of each of these vectors. Find the projection of v_1 onto v_2 and onto v_3 .

Question 5

Let A be an $m \times n$ matrix. Show that $A^T A$ and AA^T is a symmetric matrix. Assume that $m \geq n$ and $\text{rank}(A) = n$. Show that if $P = A(A^T A)^{-1}A^T$ then

$$P^2 = P$$

Solution: A matrix is symmetric if it is equal to its transpose. Using the properties of the matrix transpose, we can show that $A^T A$ and AA^T are symmetric.

$$\text{Transpose of } A^T A: (A^T A)^T = (A^T)^T A^T = AA^T = A^T A$$

$$\text{Transpose of } AA^T: (AA^T)^T = A^T (A^T)^T = A^T A = AA^T$$

As we can see here, applying the transpose to either matrix results in the same matrix. Proving that $P^2 = P$ is a bit more involved.

Proof. We have that $P = A(A^T A)^{-1}A^T$. We need to find

$$P^2 = A(A^T A)^{-1}A^T A(A^T A)^{-1}A^T$$

and show that it is equal to P . Using the associative property of matrix multiplication, we can change our order of multiplication to get

$$P^2 = A(A^T A)^{-1}(A^T A)(A^T A)^{-1}A^T$$

Since it is given that $\text{rank}(A) = n$, and $A^T A$ must be an $n \times n$ matrix, $A^T A$ is invertible. Since $A^T A$ is invertible, we can multiply it by its inverse to get the identity matrix. Now, we have

$$P^2 = AI(A^T A)^{-1}A^T = A(A^T A)^{-1}A^T = P$$

Thus, $P^2 = P$. □