

# Math2033 TA note 7

Yang Yunfei, Chen Yipei, Liu Ping  
March 24, 2019

## 1 LIMIT

**Example 1.** Let  $\{a_n\}$  be a sequence satisfying  $\lim_{n \rightarrow \infty} (a_{n+1} - \frac{a_n}{2}) = 0$ . Prove that  $\lim_{n \rightarrow \infty} a_n = 0$ .

*Solution:* Because

$$\lim_{n \rightarrow \infty} (a_{n+1} - \frac{a_n}{2}) = 0.$$

Then for  $\epsilon > 0, \exists K_1 \in \mathbb{N}, \forall n \geq K_1, |a_{n+1} - \frac{a_n}{2}| < \frac{\epsilon}{4}$ . Then

$$\forall n \geq K_1, |a_{n+1}| < \frac{|a_n|}{2} + \frac{\epsilon}{4}.$$

Then we prove by mathematical induction principle that

$$\forall n \geq K_1, |a_n| \leq \frac{|a_{K_1}|}{2^{n-K_1}} + \frac{\epsilon}{2}.$$

Firstly,  $|a_{K_1+1}| \leq \frac{|a_{K_1}|}{2} + \frac{\epsilon}{2}$ . If when  $n = k, |a_k| \leq \frac{|a_{K_1}|}{2^{k-K_1}} + \frac{\epsilon}{2}$ . Then for  $n = k + 1$ ,

$$\begin{aligned} |a_{k+1}| &\leq \frac{|a_k|}{2} + \frac{\epsilon}{4} \\ &\leq \frac{\frac{|a_{K_1}|}{2^{k-K_1}} + \frac{\epsilon}{2}}{2} + \frac{\epsilon}{4} \\ &= \frac{|a_{K_1}|}{2^{k+1-K_1}} + \frac{\epsilon}{2}. \end{aligned}$$

So by mathematical induction principle, the statement is true that

$$\forall n \geq K_1, |a_n| \leq \frac{|a_{K_1}|}{2^{n-K_1}} + \frac{\epsilon}{2}.$$

There exist  $K_2$  such that  $\forall n \geq K_2, \frac{|a_{K_1}|}{2^{n-K_1}} < \frac{\epsilon}{2}$ . So for  $\epsilon > 0, \exists K = \max\{K_1, K_2\}, \forall n \geq K$ ,

$$|a_n| \leq \frac{|a_{K_1}|}{2^{n-K_1}} + \frac{\epsilon}{2} < \epsilon.$$

That is  $\lim_{n \rightarrow \infty} a_n = 0$ .

**Example 2.** If  $\{x_n\}$  is a sequence such that  $|x_{k+1} - x_k| < \frac{1}{2^k}$  for  $k = 1, 2, 3, \dots$ , then show that  $\{x_n\}$  is a Cauchy sequence.

*Solution:* For every  $\epsilon > 0$ , there is a integer  $K = 2 - \lfloor \frac{\ln \epsilon}{\ln 2} \rfloor$ . Then  $n \geq K, p \in \mathbb{N}$  implies

$$\begin{aligned} |x_{n+p} - x_n| &= |(x_{n+p} - x_{n+p-1}) + (x_{n+p-1} - x_{n+p-2}) + \dots + (x_{n+1} - x_n)| \\ &\leq |x_{n+p} - x_{n+p-1}| + |x_{n+p-1} - x_{n+p-2}| + \dots + |x_{n+1} - x_n| \\ &< \frac{1}{2^{n+p-1}} + \frac{1}{2^{n+p-2}} + \dots + \frac{1}{2^n} \\ &< \sum_{j=n}^{\infty} \frac{1}{2^j} = \frac{1}{2^{n-1}} \\ &\leq \frac{1}{2^{K-1}} < \epsilon \end{aligned}$$

Therefore,  $\{x_n\}$  is a Cauchy sequence.

**Example 3.** We can find a divergent sequence  $\{a_n\}$  such that for any  $p \in \mathbb{N}$ ,  $\lim_{n \rightarrow \infty} (a_{n+p} - a_n) = 0$ .

*Solution:* Take

$$a_n = 1 + \frac{1}{2} + \dots + \frac{1}{n}.$$

It is divergent. However, for any  $p \in \mathbb{N}$ ,

$$a_{n+p} - a_n = \frac{1}{n+1} + \dots + \frac{1}{n+p} \leq \frac{p}{n+1} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

**Remark 4.** The Cauchy sequence,  $p$  is independent of  $n$  while the previous one is dependent of  $n$ .

**Definition 5.**  $f(x)$  converges to  $L$  as  $x$  tends to  $x_0$  in  $S$  iff for every  $\epsilon > 0$ , there exists a  $\delta > 0$  such that  $x \in S$  and  $0 < |x - x_0| < \delta$  imply  $|f(x) - L| < \epsilon$ .

**Example 6.** (1) Let  $f : (1, 3) \rightarrow \mathbb{R}$  be defined by  $f(x) = x^2 + \frac{1}{x}$ . Prove that  $\lim_{x \rightarrow 2} f(x) = \frac{9}{2}$  by checking the definition.

(2) Let  $f : (1, 4) \rightarrow \mathbb{R}$  be defined by  $f(x) = |x^2 - 9|$ . Prove that  $\lim_{x \rightarrow 2} f(x) = 5$  by checking the definition.

*Solution:* (1) For every  $\epsilon > 0$ , take  $\delta = 2\epsilon/11 > 0$ . If  $0 < |x - 2| < \delta$  and  $x \in (1, 3)$ , then

$$\begin{aligned} \left| f(x) - \frac{9}{2} \right| &= \left| x^2 + \frac{1}{x} - \frac{9}{2} \right| = \left| (x^2 - 4) + \left( \frac{1}{x} - \frac{1}{2} \right) \right| \leq |x^2 - 4| + \left| \frac{1}{x} - \frac{1}{2} \right| \\ &= |x+2||x-2| + \frac{|x-2|}{2|x|} \leq 5|x-2| + \frac{1}{2}|x-2| = \frac{11}{2}|x-2| < \frac{11}{2}\delta = \epsilon. \end{aligned}$$

(2) For every  $\epsilon > 0$ , take  $\delta = \min\{1, \epsilon/5\} > 0$ . If  $0 < |x - 2| < \delta$  and  $x \in (1, 4)$ , then  $x \in (1, 3)$  and  $f(x) = |x^2 - 9| = 9 - x^2$ . Therefore

$$|f(x) - 5| = |9 - x^2 - 5| = |4 - x^2| = |2 - x||2 + x| < 5|x - 2| < 5\delta \leq \epsilon.$$