

MATH2023 Multivariable Calculus 2013

From the textbook [Calculus of Several Variables \(5th\)](#) by R. Adams, Addison Wesley.

Homework 8

(Total: 12 questions)

Ex. 15.1

- 7 Sketch the plane vector field $\mathbf{F}(x, y) = \nabla \ln(x^2 + y^2)$ and determine its field lines.
- 9 Describe the streamlines of the velocity fields $\mathbf{v}(x, y, z) = y\mathbf{i} - y\mathbf{j} - y\mathbf{k}$.
- 16 Describe the streamlines of the velocity fields $\mathbf{v}(x, y) = x\mathbf{i} + (x + y)\mathbf{j}$. (Hint: let $y = xv(x)$.)

Ex. 15.3

- 2 Let C be the conical helix with parametric equations $x = t \cos t$, $y = t \sin t$, $z = t$, $(0 \leq t \leq 2\pi)$.
Find $\int_C z \, ds$.
- 8 Find $\int_C \sqrt{1 + 4x^2 z^2} \, ds$, where C is the curve of intersection of the surfaces $x^2 + z^2 = 1$ and $y = x^2$.
- 15 Find $\int_C \frac{ds}{(2y^2 + 1)^{3/2}}$, where C is the parabola $z^2 = x^2 + y^2$, $x + z = 1$.

Ex. 15.4

- 3 Evaluate the line integral of the tangential component of the vector field

$$\mathbf{F}(x, y, z) = y\mathbf{i} + z\mathbf{j} - x\mathbf{k}$$

along the straight line from $(0, 0, 0)$ to $(1, 1, 1)$.

- 5 Evaluate the line integral of the tangential component of the vector field $\mathbf{F}(x, y, z) = yz\mathbf{i} + xz\mathbf{j} + xy\mathbf{k}$ from $(-1, 0, 0)$ to $(1, 0, 0)$ along either direction of the curve of intersection of the cylinder $x^2 + y^2 = 1$ and the plane $z = y$.
- 11 Determine the values of A and B for which the vector field

$$\mathbf{F} = Ax \ln z \mathbf{i} + By^2 z \mathbf{j} + \left(\frac{x^2}{z} + y^3 \right) \mathbf{k}$$

is conservative. If C is the straight line from $(1, 1, 1)$ to $(2, 1, 2)$, find

$$\int_C 2x \ln z \, dx + 2y^2 z \, dy + y^3 \, dz.$$

- 13 If C is the intersection of $z = \ln(1 + x)$ and $y = x$ from $(0, 0, 0)$ to $(1, 1, \ln 2)$, evaluate

$$\int_C (2x \sin(\pi y) - e^z) \, dx + (\pi x^2 \cos(\pi y) - 3e^z) \, dy - xe^z \, dz.$$

- 14 Is each of the following sets a domain? a connected domain? a simply connected domain?

- (a) the set of points (x, y) in the plane such that $x > 0$ and $y \geq 0$
- (b) the set of points (x, y) in the plane such that $x = 0$ and $y \geq 0$
- (c) the set of points (x, y) in the plane such that $x \neq 0$ and $y > 0$
- (d) the set of points (x, y, z) in 3-space such that $x^2 > 1$
- (e) the set of points (x, y, z) in 3-space such that $x^2 + y^2 > 1$
- (f) the set of points (x, y, z) in 3-space such that $x^2 + y^2 + z^2 > 1$

- 22 Evaluate $\frac{1}{2\pi} \oint_C \frac{-y \, dx + x \, dy}{x^2 + y^2}$

- (a) counterclockwise around the circle $x^2 + y^2 = a^2$,
- (b) clockwise around the square with vertices $(-1, -1)$, $(-1, 1)$, $(1, 1)$, and $(1, -1)$,
- (c) counterclockwise around the boundary of the region $1 \leq x^2 + y^2 \leq 2$, $y \geq 0$.

Homework 9

(Total: 9 questions)

Ex. 15.2

- 5 Determine whether the vector field

$$\mathbf{F}(x, y, z) = (2xy - z^2)\mathbf{i} + (2yz + x^2)\mathbf{j} - (2zx - y^2)\mathbf{k}$$

is conservative and find a potential if it is conservative.

- 7 Find the three-dimensional vector field with potential $\phi(\mathbf{r}) = \frac{1}{\|\mathbf{r} - \mathbf{r}_0\|^2}$.

- 9 Show that the vector field

$$\mathbf{F}(x, y, z) = \frac{2x}{z}\mathbf{i} + \frac{2y}{z}\mathbf{j} - \frac{x^2 + y^2}{z^2}\mathbf{k}$$

is conservative, and find its potential. Describe the equipotential surfaces. Find the field lines of \mathbf{F} .

Ex. 15.5

- 10 Find the area of the part of the cylinder $x^2 + z^2 = a^2$ that lies inside the cylinder $y^2 + z^2 = a^2$.
- 14 Find $\iint_S y \, dS$, where S is the part of the cone $z = \sqrt{2(x^2 + y^2)}$ that lies below the plane $z = 1 + y$.

Ex. 15.6

- 1 Find the flux of $\mathbf{F} = x\mathbf{i} + z\mathbf{j}$ out of the tetrahedron bounded by the coordinate planes and the plane $x + 2y + 3z = 6$.
- 6 Find the flux of $\mathbf{F} = x\mathbf{i} + x\mathbf{j} + \mathbf{k}$ upward through the part of the surface $z = x^2 - y^2$ lying inside the cylinder $x^2 + y^2 = a^2$.
- 10 Find the flux of $\mathbf{F} = 2x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ upward through the surface $\mathbf{r} = u^2v\mathbf{i} + uv^2\mathbf{j} + v^3\mathbf{k}$, ($0 \leq u \leq 1$, $0 \leq v \leq 1$).
- 15 Define the flux of a *plane* vector field across a piecewise smooth *curve*. Find the flux of $\mathbf{F} = x\mathbf{i} + y\mathbf{j}$ outward across
- the circle $x^2 + y^2 = a^2$.
 - the boundary of the square $-1 \leq x, y \leq 1$.

Homework 10

(Total: 9 questions)

Ex. 16.4

- 4 Use the Divergence Theorem to calculate the flux of the vector field

$$\mathbf{F} = x^3\mathbf{i} + 3yz^2\mathbf{j} + (3y^2z + x^2)\mathbf{k}$$

out of the sphere S with equation $x^2 + y^2 + z^2 = a^2$, where $a > 0$.

- 8 Evaluate the flux of $\mathbf{F} = x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}$ outward across the boundary of the solid cylinder $x^2 + y^2 \leq 2y$, $0 \leq z \leq 4$.
- 11 A conical domain with vertex $(0, 0, b)$ and axis along the z -axis has as base a disk of radius a in the xy -plane. Find the flux of

$$\mathbf{F} = (x + y^2)\mathbf{i} + (3x^2y + y^3 - x^3)\mathbf{j} + (z + 1)\mathbf{k}$$

upward through the conical part of the surface of the domain.

- 23 If \mathbf{F} is a smooth vector field on D , show that

$$\iiint_D \phi \nabla \cdot \mathbf{F} \, dV + \iiint_D \nabla \phi \cdot \mathbf{F} \, dV = \iint_S \phi \mathbf{F} \cdot \hat{\mathbf{n}} \, dS.$$

- 24 If $\nabla^2 \phi = 0$ in D and $\phi(x, y, z) = 0$ on S , show that $\phi(x, y, z) = 0$ in D .

Ex. 16.5

- 2 Evaluate $\oint_C y \, dx - x \, dy + z^2 \, dz$ around the curve C of intersection of the cylinders $z = y^2$ and $x^2 + y^2 = 4$, oriented counterclockwise as seen from a point high on the z -axis.

- 3 Evaluate $\iint_S \nabla \times \mathbf{F} \cdot \hat{\mathbf{n}} \, dS$, where S is the hemisphere $x^2 + y^2 + z^2 = a^2$, $z \geq 0$ with outward normal, and $\mathbf{F} = 3y\mathbf{i} - 2xz\mathbf{j} + (x^2 - y^2)\mathbf{k}$.

- 8 Evaluate $\oint_C \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F} = ye^x\mathbf{i} + (x + e^x)\mathbf{j} + z^2\mathbf{k}$ and C is the curve

$$\mathbf{r} = (1 + \cos t)\mathbf{i} + (1 + \sin t)\mathbf{j} + (1 - \sin t - \cos t)\mathbf{k},$$

where $0 \leq t \leq 2\pi$.

- 9 Let C_1 be a straight line joining $(-1, 0, 0)$ to $(1, 0, 0)$ and let C_2 be the semicircle $x^2 + y^2 = 1$, $z = 0$, $y \geq 0$. Let S be a smooth surface joining C_1 to C_2 having upward normal, and let

$$\mathbf{F} = (\alpha x^2 - z)\mathbf{i} + (xy + y^3 + z)\mathbf{j} + \beta y^2(z + 1)\mathbf{k}.$$

Find the values of α and β for which $I = \iint_S \mathbf{F} \cdot \hat{\mathbf{n}} \, dS$ is independent of the choice of S , and find the value of I for these values of α and β .