MATH 102

Second Midterm Examination

Multivariable and Vector Calculus

15 Dec 2006

Answer ALL 8 questions

Time allowed - 180 minutes

Problem 1

(a) Find an equation of the plane through (-1, 4, -3) and perpendicular to the line

$$x = t + 2$$
, $y = 2t - 3$, $z = -t$.

- (b) Find a rectangular equation for the surface whose spherical equation is $\rho = 2\sin\theta\sin\phi$. Describe the surface.
- (c) Show that the two lines $\mathbf{r} = \mathbf{a} + \mathbf{v}t$ and $\mathbf{r} = \mathbf{b} + \mathbf{u}t$, where t is a parameter and \mathbf{a} , \mathbf{b} , \mathbf{u} and \mathbf{v} are constant vectors, will intersect if $(\mathbf{a} \mathbf{b}) \cdot (\mathbf{u} \times \mathbf{v}) = 0$.

Problem2

- (a) Describe the graph of the equation $\mathbf{r}_1(t) = -2\mathbf{i} + t\mathbf{j} + (t^2 1)\mathbf{k}$. Find also the vector equation of the tangent line to the curve $\mathbf{r}_1(t)$ such that it is parallel to the line $\mathbf{r}_2(t) = \mathbf{i} + (2 + 2t)\mathbf{j} + (3 + 4t)\mathbf{k}$.
- (b) Sketch the surfaces x + y = 4 and $\frac{y^2}{4^2} + \frac{z^2}{2^2} = 1$ in the first octant. Find the parametric equations of the curve C of intersection of the two surfaces above. Find the parametric equation of the projection curve C onto the xz-plane. Describe the projection curve.

Problem 3

- (a) Sketch the domain of the function $f(x,y) = \frac{\ln(x+y+1)}{x^2-1}$.
- (b) Determine the largest set on which the function

$$f(x,y) = \begin{cases} \frac{x^2y}{x^2 + y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0). \end{cases}$$

is continuous.

(c) Describe the level surfaces of the function $f(x, y, z) = (x - 2)^2 + y^2$.

Problem 4

- (a) Let $f(x,y) = \sqrt{3x + 2y}$.
 - (i) Find the slope of the surface z = f(x, y) in the x-direction at the point (4, 2).
 - (ii) Find the slope of the surface z = f(x, y) in the y-direction at the point (4, 2).
- (b) Let $g(x,y) = (x^2 + y^3)^{\frac{2}{3}}$. Find $g_x(x,y)$, at all points (x,y) in the xy-plane (include the point (0,0)).
- (c) Find $\frac{\partial^3}{\partial t^2 \partial s} f(s^2 t, s + t^2)$ in terms of partial derivatives of f. Assume that f has continuous partial derivatives of all orders.

Problem 5

- (a) If $f(x, y, z) = (\mathbf{r} \times \mathbf{A}) \cdot (\mathbf{r} \times \mathbf{B})$, where $\mathbf{r} = x \mathbf{i} + y \mathbf{j} + z \mathbf{k}$ and \mathbf{A} and \mathbf{B} are constant vectors, show that $\nabla f(x, y, z) = \mathbf{P} \times (\mathbf{r} \times \mathbf{A}) + \mathbf{Q} \times (\mathbf{r} \times \mathbf{B})$. Find \mathbf{P} and \mathbf{Q} in terms of \mathbf{A} , \mathbf{B} and \mathbf{r} .
- (b) Let $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and let $r = ||\mathbf{r}||$. If **A** and **B** are constant vectors, show that:
 - (i) $\mathbf{A} \cdot \nabla \left(\frac{1}{r}\right) = \frac{\mathbf{C}}{r^3}$. Find \mathbf{C} in terms of \mathbf{A} and \mathbf{r} .
 - (ii) $\mathbf{B} \cdot \nabla \left(\mathbf{A} \cdot \nabla \left(\frac{1}{r} \right) \right) = \frac{\mathbf{D}}{r^5} \frac{\mathbf{A} \cdot \mathbf{B}}{r^3}$. Find \mathbf{D} in terms of \mathbf{A} , \mathbf{B} and \mathbf{r} .

Problem 6

A three dimensional surface whose equation is y = f(x) is tangent to the surface $z^2 + 2xz + y = 0$ at all points common to the two surfaces. (i) Find f(x). (ii) Find all common points.

Problem 7

- (a) Let f be a nonconstant scalar field, differentiable everywhere in the plane, and let c be a constant. Assume the Cartesian equation f(x,y) = c describes a curve C having a tangent at each of its points. Prove that f has the following properties at each point of C:
 - (i) The gradient vector ∇f is normal to C.
 - (ii) The directional derivative of f is zero along C.
 - (iii) The directional derivative of f has its largest value in a direction normal to C.
- (b) Find the directional derivative of the scalar field $f(x,y) = x^2 3xy$ along the parabola $y = x^2 x + 2$ at the point (1,2).

Problem 8

A manufacturer is planning to sell a new product at the price of \$150 per unit and estimates that if x thousand dollars is spent on development and y thousand dollars is spent on promotion, approximately $\frac{320y}{y+2} + \frac{160x}{x+4}$ units of the product will be sold. The cost of manufacturing the product is \$50 per unit. If the manufacturer has a total \$8,000 to spend on development and promotion, how should this money be allocated to generate the largest possible profit?

Suppose the manufacturer in the above exercise decides to spend \$8,100 instead of \$8,000 on the development and promotion of the new product. Use the Lagrange multiplier λ to estimate how this change will affect the maximum possible profit.