

M10 TOBACCO CESSATION | O (MAX : 3 PT)

Intent : Reduce the use of tobacco through interventions that support tobacco cessation and prevent the sale and advertisement of tobacco products.

Summary : This WELL feature requires projects that sell retail goods to restrict sale and marketing of tobacco products and supports employee access to tobacco cessation support programs.

Issue : Tobacco is responsible for an estimated six million deaths per year globally among direct users, and serves as the cause of death for up to half of its users.^{1,2} In addition to those deaths caused by direct use, an estimated 890,000 annual deaths can be attributed to non-user exposure to second-hand smoke.¹ In the workplace, employees that smoke incur greater absences, take more sick days and have higher health care costs than non-smoking employees.³ Despite tobacco's impact, national comprehensive health services that fully or partially cover services to support tobacco cessation are only available in 24 countries, benefiting just 15% of the world's population.^{1,4} Although 70% of U.S. adult smokers are interested in quitting and 40% of smokers attempt to quit each year, only 8% of smoking employees report that their workplace offers smoking cessation assistance.⁵⁻⁷

Solutions : Employers can play a key role in supporting employee tobacco cessation efforts.⁷ Among those who attempt to quit, counseling and medication more than double the chance of quitting success.¹ Worksite-based incentives and competitions, when combined with additional interventions to support individual cessation efforts, can be effective in reducing tobacco use among workers.⁸ Another influencing factor on tobacco use is an individual's proximity to outlets where it is sold.⁹ Restricting the sale of tobacco on-site is a key strategy for preventing or curbing use of tobacco products, as well as providing support to those trying to quit.^{9,10}

PART 1 PROVIDE TOBACCO CESSATION RESOURCES (MAX : 2 PT)

For All Spaces:

1: Incentive program and cessation resources

The following requirements are met:

- a. The project or organization implements a tobacco cessation program for all eligible employees that meets the following:
 1. Focuses on increasing or improving motivation or action to quit, or maintaining quit effort.^{8,11}
 2. Includes incentives or rewards (e.g., direct financial payments, lottery for prizes) provided for participation in quit effort or success in abstaining from tobacco use.^{8,11}
- b. Tobacco cessation resources are available to all eligible employees at no cost or are subsidized and meet the following:
 1. Resources referring tobacco users to tobacco cessation telephone quit lines or online quitting resources.¹¹
 2. Tobacco cessation counseling. Programs may be provided in-person or virtually; in group or individual settings; through vendors, on-site staff, health insurance plans or programs, community groups or other qualified professionals (e.g., tobacco cessation specialist).¹¹
 3. Prescription tobacco cessation medications and nicotine replacement products (e.g., inhalers, nasal sprays, bupropion, varenicline).¹¹
 4. Over-the-counter nicotine replacement products (e.g., nicotine gum, patches, lozenges).¹¹

PART 2 LIMIT TOBACCO AVAILABILITY (MAX : 1 PT)

For Retail Spaces:

The following requirements are met for projects where retail products are sold on a daily basis:

- a. Sale of tobacco products (including e-cigarettes) is prohibited.¹¹
- b. Tobacco products (including e-cigarettes) are not marketed or promoted.¹²

REFERENCES

1. World Health Organization. Tobacco. <http://www.who.int/mediacentre/factsheets/fs339/en/>. Published 2017. Accessed January 9, 2018.
2. World Health Organization. WHO global report on trends in prevalence of tobacco smoking 2015. WHO Mag. 2015;1-359. doi:978 92 4 156492 2
3. Smith DR. Workplace tobacco control: The nexus of public and occupational health. Public Health. 2009;123(12):817-819. doi:10.1016/j.puhe.2009.10.014
4. World Health Organization. Global status report on noncommunicable diseases 2010. World Heal Organ. 2011;176. doi:ISBN 978 92 4 156422 9
5. Centers for Disease Control and Prevention. Cigarette Smoking Among Adults --- United States, 2000. MMWR Wkly. 2002;51(29):642-645. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5129a3.htm>.
6. Centers for Disease Control and Prevention. Cigarette Smoking Among Adults --- United States, 2007. MMWR Wkly. 2008;57(45):1221-1226. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5745a2.htm>.

7. Halpern MT, Taylor H. Employee and Employer Support for Workplace-based Smoking Cessation: Results from an International Survey. *J Occup Health*. 2010;52(6):375-382. doi:10.1539/joh.L10075
8. The Community Preventive Services Task Force. Reducing Tobacco Use and Secondhand Smoke Exposure: Worksite-based Incentives and Competitions When Combined with Additional Interventions. 2014;38.
9. Reitzel LR, Cromley EK, Li Y, et al. The Effect of Tobacco Outlet Density and Proximity on Smoking Cessation. *Am J Public Health*. 2011;101(2):315-320. doi:10.2105/AJPH.2010.191676
10. Polinski J, Howell B, Gagnon M, Kymes S, Brennan T, Shrank W. Impact of CVS Pharmacy's Discontinuance of Tobacco Sales on Cigarette Purchasing (2012-2014). *Am J Public Health*. 2017;107(4):556-562. doi:10.2105/AJPH.2016.303612
11. Centers for Disease Control and Prevention. The CDC Worksite Health ScoreCard: An Assessment Tool for Employers to Prevent Heart Disease, Stroke, and Related Health Conditions. Atlanta; 2014. https://www.cdc.gov/dhdsp/pubs/docs/hsc_manual.pdf.
12. Calo WA, Krasny S. Environmental Determinants of Smoking Behaviors: The Role of Policy and Environmental Interventions in Preventing Smoking Initiation and Supporting Cessation. *Curr Cardiovasc Risk Rep*. 2013;7(6):446-452. doi:10.1007/s12170-013-0344-7