

# EFFECTIVE APPROXIMATION OF HEAT FLOW EVOLUTION OF THE RIEMANN XI FUNCTION, AND AN UPPER BOUND FOR THE DE BRUIJN-NEWMAN CONSTANT

D.H.J. POLYMATH

ABSTRACT. For each  $t \in \mathbb{R}$ , define the entire function

$$H_t(z) := \int_0^\infty e^{tu^2} \Phi(u) \cos(zu) du$$

where  $\Phi$  is the super-exponentially decaying function

$$\Phi(u) := \sum_{n=1}^\infty (2\pi^2 n^4 e^{9u} - 3\pi n^2 e^{5u}) \exp(-\pi n^2 e^{4u}).$$

Newman showed that there exists a finite constant  $\Lambda$  (the *de Bruijn-Newman constant*) such that the zeroes of  $H_t$  are all real precisely when  $t \geq \Lambda$ . The Riemann hypothesis is the equivalent to the assertion  $\Lambda \leq 0$ ; recently, Rodgers and Tao established the matching lower bound  $\Lambda \geq 0$ . Ki, Kim and Lee established the upper bound  $\Lambda < \frac{1}{2}$ .

In this paper we establish several effective estimates on  $H_t(x + iy)$ , including some that are accurate for small or medium values of  $x$ . By combining these estimates with numerical computations, we are able to obtain a new upper bound  $\Lambda \leq 0.28$ ; we also obtain some new estimates controlling the asymptotic behavior of zeroes of  $H_t(x + iy)$  as  $x \rightarrow \infty$ .

## 1. INTRODUCTION

Let  $H_0: \mathbb{C} \rightarrow \mathbb{C}$  denote the function

$$(1) \quad H_0(z) := \frac{1}{8} \xi\left(\frac{1}{2} + \frac{iz}{2}\right),$$

where  $\xi$  denotes the Riemann xi function

$$(2) \quad \xi(s) := \frac{s(s-1)}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s)$$

(after removing all singularities) and  $\zeta$  is the Riemann zeta function. Then  $H_0$  is an entire even function with functional equation  $H_0(\bar{z}) = \overline{H_0(z)}$ , and the Riemann hypothesis is equivalent to the assertion that all the zeroes of  $H_0$  are real.

It is a classical fact (see [23, p. 255]) that  $H_0$  has the Fourier representation

$$H_0(z) = \int_0^\infty \Phi(u) \cos(zu) du$$

where  $\Phi$  is the super-exponentially decaying function

$$(3) \quad \Phi(u) := \sum_{n=1}^\infty (2\pi^2 n^4 e^{9u} - 3\pi n^2 e^{5u}) \exp(-\pi n^2 e^{4u}).$$

The sum defining  $\Phi(u)$  converges absolutely for negative  $u$  also. From Poisson summation one can verify that  $\Phi$  satisfies the functional equation  $\Phi(u) = \Phi(-u)$  (i.e.,  $\Phi$  is even).

De Bruijn [4] introduced the more general family of functions  $H_t: \mathbb{C} \rightarrow \mathbb{C}$  for  $t \in \mathbb{R}$  by the formula

$$(4) \quad H_t(z) := \int_0^\infty e^{tu^2} \Phi(u) \cos(zu) du.$$

As noted in [8, p.114], one can view  $H_t$  as the evolution of  $H_0$  under the backwards heat equation  $\partial_t H_t(z) = -\partial_{zz} H_t(z)$ . As with  $H_0$ , each of the  $H_t$  are entire even functions with functional equation  $H_t(\bar{z}) = \overline{H_t(z)}$ . De Bruijn showed that the zeroes of  $H_t$  are purely real for  $t \geq 1/2$ , and if  $H_t$  has purely real zeroes for some  $t$ , then  $H_{t'}$  has purely real zeroes for all  $t' > t$ . Newman [13] strengthened this result by showing that there is an absolute constant  $-\infty < \Lambda \leq 1/2$ , now known as the *De Bruijn-Newman constant*, with the property that  $H_t$  has purely real zeroes if and only if  $t \geq \Lambda$ . The Riemann hypothesis is then clearly equivalent to the upper bound  $\Lambda \leq 0$ . Recently in [18] the complementary bound  $\Lambda \geq 0$  was established, answering a conjecture of Newman [13]. Furthermore, Ki, Kim, and Lee [9] sharpened the upper bound  $\Lambda \leq 1/2$  of de Bruijn [4] slightly to  $\Lambda < 1/2$ .

## 2. NOTATION

Unless otherwise specified,  $\log$  denotes the standard branch of the complex logarithm, thus the branch cut is on the negative real axis and imaginary part in  $(-\pi, \pi]$ . We then define the standard complex powers  $z^w := \exp(w \log z)$ , and in particular define the standard square root  $\sqrt{z} := z^{1/2}$ . We record the standard gaussian identity

$$(5) \quad \int_{\mathbb{R}} \exp(-(au^2 + bu + c)) du = \sqrt{\frac{\pi}{a}} \exp\left(\frac{b^2}{4a} - c\right)$$

for any complex numbers  $a, b, c$  with  $\operatorname{Re} a > 0$ .

To obtain effective estimates, it is convenient to use the notation  $O_{\leq}(X)$  to denote any quantity that is bounded in magnitude by  $X$ . Any expression of the form  $A = B$  using this notation should be interpreted as the assertion that any quantity of the form  $A$  is also of the form  $B$ , thus for instance  $O_{\leq}(1) + O_{\leq}(1) = O_{\leq}(3)$ . (In particular, the equality relation is no longer symmetric with this notation.)

If  $F$  is a meromorphic function, we use  $F'$  to denote its derivative. We also use  $F^*$  to denote the reflection  $F^*(s) := \overline{F(\bar{s})}$  of  $F$ . Observe from analytic continuation that if  $F: \mathbb{C} \rightarrow \mathbb{C}$  is meromorphic and is real-valued on  $\mathbb{R}$  then it is equal to its own reflection:  $F = F^*$ .

We use  $x_+ := \max(x, 0)$  to denote the positive part of a real number  $x$ .

## 3. DYNAMICS OF ZEROES

**Proposition 3.1** (Dynamics of a single zero). *Let  $t_0 \in \mathbb{R}$ , and let  $(z_k(t_0))_{k \in \mathbb{Z} \setminus \{0\}}$  be an enumeration of the zeroes of  $H_{t_0}$  (counting multiplicity), with the symmetry condition  $z_{-k}(t_0) = -z_k(t_0)$ .*

- (i) *If  $j \in \mathbb{Z} \setminus \{0\}$  is such that  $z_j(t_0)$  is a simple zero of  $H_{t_0}$ , then there exists a neighbourhood  $U$  of  $z_j(t_0)$ , a neighbourhood  $I$  of  $t_0$  in  $\mathbb{R}$ , and a smooth map  $z_j: I \rightarrow U$  such that for*

every  $t \in I$ ,  $z_j(t)$  is the unique zero of  $H_t$  in  $U$ . Furthermore one has the equation

$$(6) \quad \left( \frac{\partial}{\partial t} z_j \right) (t_0) = 2 \sum'_{k \neq j} \frac{1}{z_j(t_0) - z_k(t_0)}$$

where the sum is over those  $k \in \mathbb{Z} \setminus \{0\}$  with  $k \neq j$ , and the prime means that the  $k$  and  $-k$  terms are summed together (assuming  $k \neq \pm j$ ) in order to make the sum convergent.

- (ii) If  $j \in \mathbb{Z} \setminus \{0\}$  is such that  $z_j(t_0)$  is a repeated zero of  $H_{t_0}$  of order  $m \geq 2$ , then there is a neighbourhood  $U$  of  $z_j(t_0)$  such that for  $t$  sufficiently close to  $t_0$ , there are precisely  $m$  zeroes of  $H_t$  in  $U$ , and they take the form

$$z_j(t_0) + \sqrt{2}(t - t_0)^{1/2} x_j + O(|t - t_0|)$$

for  $j = 1, \dots, m$  as  $t \rightarrow t_0$ , where  $x_1 < \dots < x_m$  are the roots of the  $m^{\text{th}}$  Hermite polynomial

$$(7) \quad \text{He}_m(z) := (-1)^m \exp\left(\frac{z^2}{2}\right) \frac{d^m}{dz^m} \exp\left(-\frac{z^2}{2}\right)$$

$$(8) \quad = \sum_{0 \leq l \leq m/2} \frac{m!}{l!(m-2l)!} (-1)^l \frac{z^{m-2l}}{2^l}$$

and the implied constant in the  $O()$  notation can depend on  $t_0$ ,  $j$ , and  $m$ .

The differential equation (6) was previously derived in [8, Lemma 2.4] in the case  $t > \Lambda$  (in which all zeroes are real and simple). The  $x_1, \dots, x_m$  can be given explicitly as

$$x_1 = -1; \quad x_2 = +1$$

when  $m = 2$ ,

$$x_1 = -\sqrt{3}; \quad x_2 = 0; \quad x_3 = +\sqrt{3}$$

when  $m = 3$ , and

$$x_1 = -\sqrt{3 + \sqrt{6}}; \quad x_2 = -\sqrt{3 - \sqrt{6}}; \quad x_3 = \sqrt{3 - \sqrt{6}}; \quad \sqrt{3 + \sqrt{6}}$$

when  $m = 4$ . From (7) and iterating Rolle's theorem we see that all the zeroes  $x_1, \dots, x_m$  of  $\text{He}_m$  are real; from the Hermite equation  $\left(\frac{d^2}{dz^2} - z\frac{d}{dz} + m\right)\text{He}_m(z) = 0$  and the Picard uniqueness theorem for ODE we see that the zeroes are all simple.

*Proof.* First suppose we are in the situation of (i). As  $z_j(t_0)$  is simple,  $\frac{\partial}{\partial z} H_t$  is non-zero at  $z_0$ ; since  $H_t(z)$  is a smooth function of both  $t$  and  $z$ , we conclude from the implicit function theorem that there is a unique solution  $z_j(t) \in U$  to the equation

$$H_t(z_j(t)) = 0$$

with  $z_j(t)$  in a sufficiently small neighbourhood  $U$  of  $z_j(t_0)$ , if  $t$  is a sufficiently small neighbourhood of  $I$ ; furthermore,  $z_j(t)$  depends smoothly on  $t$ . Differentiating the above equation at  $t_0$ , we obtain

$$\left( \frac{\partial}{\partial t} H_t \right) (z_j(t_0)) + \left( \frac{\partial}{\partial t} z_j \right) (t_0) H'_t(z_j(t_0)) = 0,$$

where the primes denote differentiation in the  $z$  variable. On the other hand, from (4) and differentiation under the integral sign (which can be justified using the rapid decrease of  $\Phi$ ) we have the backwards heat equation

$$(9) \quad \frac{\partial}{\partial t} H_t = -H_t''$$

and hence

$$(10) \quad \left( \frac{\partial}{\partial t} z_j \right) (t_0) = \frac{H_t''}{H_t'} (z_j(t_0)).$$

Henceforth we omit the dependence on  $t_0$  for brevity. From Taylor expansion of  $H_t$ ,  $H_t'$ , and  $H_t''$  around the simple zero  $z_j$  we see that

$$(11) \quad \frac{H_t''}{H_t'} (z_j) = 2 \lim_{z \rightarrow z_j} \left( \frac{H_t'}{H_t} (z_j) - \frac{1}{z - z_j} \right).$$

On the other hand, as  $H_t$  is even, zero, and entire of order 1 (as can be easily verified from (4)) we see from the Hadamard factorization theorem that

$$H_t(z) = C_t \prod_k' \left( 1 - \frac{z}{z_k} \right)$$

for some nonzero complex number  $C_t$ , where the prime indicates that the  $k$  and  $-k$  factors are multiplied together. Taking logarithmic derivatives, we conclude that

$$\frac{H_t'}{H_t}(z) = \sum_k' \frac{1}{z - z_k}.$$

Inserting this into (10), (11) and using the dominated convergence theorem (noting from the order one property of  $H_t$  that the number of zeroes  $z_k$  in a large disk  $D(0, R)$  will grow by at most  $O(R^{1+o(1)})$ ), we obtain the claim (i).

Now we prove (ii). We abbreviate  $z_j(t_0)$  as  $z_j$ . By Taylor expansion we have

$$\frac{\partial^k}{\partial z^k} H_{t_0}(z) = m(m-1) \dots (m-k+1) a_m (z - z_j)^{m-k} + O(|z - z_j|^{\max(m-k+1, 0)})$$

as  $z \rightarrow z_k$  for any fixed integer  $j \geq 0$  and some non-zero complex number  $a_m = a_m(z_j, t_0)$ ; applying the backwards heat equation (9) we thus have

$$\frac{\partial^k}{\partial t^k} H_{t_0}(z) = (-1)^k m(m-1) \dots (m-2k+1) a_m (z - z_j)^{m-2k} + O(|z - z_j|^{\max(m-2k+1, 0)}).$$

Performing Taylor expansion in time and using (8), we conclude that in the regime  $z - z_j = O(|t - t_0|^{1/2})$ , one has the bound

$$H_t(z) = a_m ((t - t_0)^{1/2})^m \left( \text{He}_m \left( \sqrt{2} \frac{z - z_j}{(t - t_0)^{1/2}} \right) + O(|t - t_0|^{1/2}) \right)$$

as  $t \rightarrow t_0$ , where we use some branch of the square root. By the inverse function theorem (and the simple nature of the zeroes of  $\text{He}_m$ ), we conclude that for  $t$  sufficiently close but not equal to  $t_0$ , we have  $m$  zeroes of  $H_t$  of the form

$$z_j + (t - t_0)^{1/2} x_j + O(|t - t_0|).$$

By Rouché's theorem, if  $U$  is a sufficiently small neighborhood of  $z_j$  then these are the only zeroes of  $H_t$  in  $U$  for  $t$  sufficiently close to  $t_0$ . The claim follows.  $\square$

We recall the following bound of de Bruijn:

**Theorem 3.2.** *Suppose that  $t_0 \in \mathbb{R}$  and  $y_0 > 0$  is such that there are no zeroes  $H_{t_0}(x + iy) = 0$  with  $x \in \mathbb{R}$  and  $y > y_0$ . Then for any  $t > t_0$ , there are no zeroes  $H_t(x + iy) = 0$  with  $x \in \mathbb{R}$  and  $y > \max(y_0^2 - 2(t - t_0), 0)^{1/2}$ . In particular one has  $\Lambda \leq t_0 + \frac{1}{2}y_0^2$ .*

*Proof.* See [4, Theorem 13].  $\square$

We can combine this with the preceding proposition to obtain the following criterion for upper bounding the de Bruijn-Newman constant.

**Theorem 3.3** (Upper bound criterion). *Suppose that  $t_0 > 0, X, y_0 > 0$  obey the following hypotheses:*

- (i) *There are no zeroes  $H_0(x + iy) = 0$  with  $0 \leq x \leq X$  and  $y \geq y_0$ .*
- (ii) *There are no zeroes  $H_{t_0}(x + iy) = 0$  with  $x \geq X + \sqrt{1 - y_0^2}$  and  $y \geq y_0$ .*
- (iii) *There are no zeroes  $H_t(x + iy) = 0$  with  $X \leq x \leq X + \sqrt{1 - y_0^2}$ ,  $y \geq y_0$ , and  $0 \leq t \leq t_0$ .*

*Then there are no zeroes  $H_{t_0}(x + iy) = 0$  with  $x \in \mathbb{R}$  and  $y \geq y_0$ . In particular, from Theorem 3.2 we have  $\Lambda \leq t_0 + \frac{1}{2}y_0^2$ .*

*Proof.* By hypotheses (ii), (iii) and the symmetry properties of  $H_t$ , it suffices to show that for every  $0 \leq t \leq t_0$ , there are no zeroes  $H_t(x + iy) = 0$  with  $0 \leq x \leq X$  and  $y \geq y_0$ . By hypothesis (i), this is true at time  $t = 0$ . Suppose the claim failed for some time  $0 < t \leq t_0$ . Let  $t_1 \in (0, t_0]$  be the minimal time in which this occurred. From Rouché's theorem (or Proposition 3.1) we conclude that there is a zero  $H_{t_1}(x + iy) = 0$  with  $x + iy$  on the boundary of the region  $\{x + iy : 0 \leq x \leq X, y \geq y_0\}$ . The right side  $x = X$  of this boundary is ruled out by hypothesis (ii), and the left side  $x = 0$  is ruled out by (4) and the positivity of  $\Phi$ . Thus by the symmetry properties of  $H_{t_1}$  we must have

$$H_{t_1}(x + iy_0) = 0$$

for some  $0 < x < X$ .

Suppose first that  $H_{t_1}$  has a repeated zero at  $x + iy_0$ . Using Proposition 3.1(ii) and observing (from the symmetry of  $\text{He}_m$ ) that at least one of the roots  $x_1, \dots, x_m$  is positive, we then see that for  $t < t_1$  sufficiently close to  $t_1$ ,  $H_t$  has a zero in the region  $\{x + iy : 0 \leq x \leq X, y \geq y_0\}$ , contradicting the minimality of  $t_1$ . Thus the zero  $x + iy_0$  of  $H_{t_1}$  must be simple. In particular, by Proposition 3.1(i) we can write  $x + iy_0 = z_j(t_1)$  for some smooth function  $z_j$  in a neighbourhood of  $t_1$  obeying (6), such that  $z_j(t)$  is a zero of  $H_t$  for all  $t$  close to  $t_1$ . We will prove that

$$(12) \quad \text{Im} \frac{\partial}{\partial t} z_j(t_1) < 0,$$

which implies that there is a zero of  $H_t$  in the region  $\{x + iy : 0 \leq x \leq X, y \geq y_0\}$  for  $t < t_1$  sufficiently close to  $t_1$ , giving the required contradiction.

By Proposition 3.1(i), the left-hand side of (12) is

$$2 \sum'_{k \neq j} \frac{y_0 - y_k}{(x - x_k)^2 + (y_0 - y_k)^2}$$

where we write  $z_k = x_k + iy_k$ . Clearly any zero  $x_k + iy_k$  with imaginary part  $y_k$  in  $[-y_0, y_0]$  gives a non-positive contribution to this sum, and at least one of the contributions is negative (e.g., the contribution of  $x - iy_0$ ). Grouping the remaining zeroes with their complex conjugates, it then suffices to show that

$$\frac{y_0 - y_k}{(x - x_k)^2 + (y_j - y_k)^2} + \frac{y_0 + y_k}{(x - x_k)^2 + (y_0 + y_k)^2} \leq 0$$

whenever  $y_k > y_0$ . Cross-multiplying and canceling like terms, this inequality eventually simplifies to

$$y_k^2 \leq (x - x_k)^2 + y_0^2.$$

But from the hypothesis (iii) and the assumption  $y_k > y_0$ , we have  $|x_k| \geq X + \sqrt{1 - y_0^2}$ , so  $(x - x_k)^2 \geq 1 - y_0^2$ . On the other hand from Theorem 3.2 one has  $y_k < 1$ , giving the required contradiction.  $\square$

#### 4. APPLYING THE FUNDAMENTAL SOLUTION FOR THE HEAT EQUATION

We can write  $H_t$  in terms of  $H_0$  using the fundamental solution to the heat equation. Namely, for any  $t > 0$ , we have from (5) that

$$e^{tu^2} = \int_{\mathbb{R}} e^{\pm 2\sqrt{t}vu} \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

for any complex  $u$  and either choice of sign  $\pm$ . Multiplying by  $e^{\pm izu}$  and averaging, we conclude that

$$e^{tu^2} \cos(zu) = \int_{\mathbb{R}} \cos((z - 2i\sqrt{t}v)u) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

for any complex  $z, u$ . Multiplying by  $\Phi(u)$  and using Fubini's theorem, we conclude that

$$H_t(z) = \int_{\mathbb{R}} H_0(z - 2i\sqrt{t}v) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

for any complex  $z$ . Using (1), we thus have

$$(13) \quad H_t(z) = \int_{\mathbb{R}} \frac{1}{8} \xi\left(\frac{1+iz}{2} + \sqrt{t}v\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv.$$

**Remark 4.1.** We have found numerically that the formula (13) gives a fast and accurate means to compute  $H_t(z)$  when  $z$  is of moderate size, e.g. if  $z = x + iy$  with  $|x| \leq 10^6$  and  $|y| \leq 1$ . However, we will not need to directly compute the right-hand side of (13) for our application to bounding  $\Lambda$ , as we will only need to control  $H_t(x + iy)$  for large values of  $x$ , and we will shortly develop tractable approximations of Riemann-Siegel type that are suitable for this regime.

We now combine this formula with expansions of the Riemann  $\xi$ -function. From [23, (2.10.6)] we have the Riemann-Siegel formula

$$(14) \quad \frac{1}{8} \xi(s) = R_{0,0}(s) + R_{0,0}^*(1-s)$$

for any complex  $s$  that is not an integer (in order to avoid the poles of the Gamma function), where  $R_{0,0}(s)$  is the contour integral

$$R_{0,0}(s) := \frac{1}{8} \frac{s(s-1)}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \int_{0 \wedge 1} \frac{w^{-s} e^{i\pi w^2}}{e^{\pi i w} - e^{-\pi i w}} dw$$

with  $0 \not\prec 1$  any infinite line oriented in the direction  $e^{5\pi i/4}$  that crosses the interval  $[0, 1]$ . From the residue theorem (and the gaussian decrease of  $e^{i\pi w^2}$  along the  $e^{\pi i/4}$  and  $e^{5\pi i/4}$  directions) we may expand

$$R_{0,0}(s) = \sum_{n=1}^N r_{0,n}(s) + R_{0,N}(s)$$

for any non-negative integer  $N$ , where  $r_{0,n}, R_{0,N}$  are the meromorphic functions

$$(15) \quad r_{0,n}(s) := \frac{1}{8} \frac{s(s-1)}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) n^{-s},$$

$$(16) \quad R_{0,N}(s) := \frac{1}{8} \frac{s(s-1)}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \int_{N \not\prec N+1} \frac{w^{-s} e^{i\pi w^2}}{e^{\pi i w} - e^{-\pi i w}}$$

and  $N \not\prec N+1$  denotes any infinite line oriented in the direction  $e^{5\pi i/4}$  that crosses the interval  $[N, N+1]$ . For any  $z$  that is not purely imaginary, we see from Stirling's approximation that the functions  $r_{0,n}(\frac{1+iz}{2} + \sqrt{t}v)$  and  $R_{0,N}(\frac{1+iz}{2} + \sqrt{t}v)$  grow slower than gaussian as  $v \rightarrow \pm\infty$  (indeed they grow like  $\exp(O(|v| \log |v|))$ , where the implied constants depend on  $t, z$ ). From this and (13), (14) we conclude that

$$(17) \quad H_t(z) = \sum_{n=1}^N r_{t,n}\left(\frac{1+iz}{2}\right) + \sum_{n=1}^N r_{t,n}^*\left(\frac{1-iz}{2}\right) + R_{t,N}\left(\frac{1+iz}{2}\right) + R_{t,N}^*\left(\frac{1-iz}{2}\right)$$

for any  $t > 0$ , any  $z$  that is not purely imaginary, and any non-negative integer  $N$ , where  $r_{t,n}(s), R_{t,N}(s)$  are defined for non-real  $s$  by the formulae

$$r_{t,n}(s) := \int_{\mathbb{R}} r_{0,n}(s + \sqrt{t}v) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

$$R_{t,N}(s) := \int_{\mathbb{R}} R_{0,N}(s + \sqrt{t}v) \frac{1}{\sqrt{\pi}} e^{-v^2} dv;$$

these can be thought of as the evolutions of  $r_{0,n}, R_{0,N}$  respectively under the forward heat equation.

The functions  $r_{0,n}(s), R_{0,N}(s)$  grow slower than gaussian as long as the imaginary part of  $s$  is bounded and bounded away from zero. As a consequence, we may shift contours (replacing  $v$  by  $v + \frac{\sqrt{t}}{2}\alpha_n$ ) and write

$$(18) \quad r_{t,n}(s) = \exp\left(-\frac{t}{4}\alpha_n^2\right) \int_{\mathbb{R}} \exp(-\sqrt{t}v\alpha_n) r_{0,n}\left(s + \sqrt{t}v + \frac{t}{2}\alpha_n\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

for any complex number  $\alpha_n$  with  $\text{Im}(s), \text{Im}(s + \frac{t}{2}\alpha_n)$  having the same sign. Similarly we may write

$$(19) \quad R_{t,N}(s) = \exp\left(-\frac{t}{4}\beta_N^2\right) \int_{\mathbb{R}} \exp(-\sqrt{t}v\beta_N) R_{0,N}\left(s + \sqrt{t}v + \frac{t}{2}\beta_N\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

for any complex number  $\beta_N$  with  $\text{Im}(s), \text{Im}(s + \frac{t}{2}\beta_N)$  having the same sign. In the spirit of the saddle point method, we will select the parameters  $\alpha_n, \beta_N$  later in the paper in order to make the phases in  $r_{0,n}, R_{0,N}$  close to stationary, in order to obtain good estimates and approximations for these terms.

## 5. ELEMENTARY ESTIMATES

We have the following elementary estimates:

**Lemma 5.1** (Elementary estimates). *Let  $x > 0$ .*

(i) *If  $a > 0$  and  $b \geq 0$  are such that  $x > b/a$ , then*

$$O_{\leq}\left(\frac{a}{x}\right) + O_{\leq}\left(\frac{b}{x^2}\right) = O_{\leq}\left(\frac{a}{x - b/a}\right).$$

*More generally, if  $a > 0$  and  $b, c \geq 0$  are such that  $x > b/a$ ,  $\sqrt{c/a}$ , then*

$$O_{\leq}\left(\frac{a}{x}\right) + O_{\leq}\left(\frac{b}{x^2}\right) + O_{\leq}\left(\frac{c}{x^3}\right) = O_{\leq}\left(\frac{a}{x - \max(b/a, \sqrt{c/a})}\right).$$

(ii) *If  $x > 1$ , then*

$$\log\left(1 + O_{\leq}\left(\frac{1}{x}\right)\right) = O_{\leq}\left(\frac{1}{x-1}\right).$$

*or equivalently*

$$1 + O_{\leq}\left(\frac{1}{x}\right) = \exp\left(O_{\leq}\left(\frac{1}{x-1}\right)\right).$$

(iii) *If  $x > 1/2$ , then*

$$\exp\left(O_{\leq}\left(\frac{1}{x}\right)\right) = 1 + O_{\leq}\left(\frac{1}{x-0.5}\right).$$

(iv) *We have*

$$\exp(O_{\leq}(x)) = 1 + O_{\leq}(e^x - 1).$$

(v) *If  $z$  is a complex number with  $|\operatorname{Im}(z)| \geq 1$  or  $\operatorname{Re} z \geq 1$ , then*

$$\Gamma(z) = \sqrt{2\pi} \exp\left(\left(z - \frac{1}{2}\right)\log z - z + O_{\leq}\left(\frac{1}{12(|z| - 0.33)}\right)\right).$$

(vi) *If  $a, b > 0$  and  $x \geq x_0 \geq \exp(a/b)$  and  $x_0 > c \geq 0$ , then*

$$\frac{\log^a x}{(x-c)^b} \leq \frac{\log^a x_0}{(x_0-c)^b}.$$

*Proof.* Claim (i) follows from the geometric series formula

$$\frac{a}{x-t} = \frac{a}{x} + \frac{at}{x^2} + \frac{at^2}{x^3} + \dots$$

whenever  $0 \leq t < x$ .

For Claim (ii), we use the Taylor expansion of the logarithm to note that

$$\log\left(1 + O_{\leq}\left(\frac{1}{x}\right)\right) = O_{\leq}\left(\frac{1}{x} + \frac{1}{2x^2} + \frac{1}{3x^3} + \dots\right)$$

which on comparison with the geometric series formula

$$\frac{1}{x-1} = \frac{1}{x} + \frac{1}{x^2} + \frac{1}{x^3} + \dots$$

gives the claim. Similarly for Claim (iii), we may compare the Taylor expansion

$$\exp\left(O_{\leq}\left(\frac{1}{x}\right)\right) = 1 + O_{\leq}\left(\frac{1}{x} + \frac{1}{2!x^2} + \frac{1}{3!x^3} + \dots\right)$$



with the geometric series formula

$$\frac{1}{x-0.5} = \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{2^2x^3} + \dots$$

and note that  $k! \geq 2^k$  for all  $k \geq 2$ .

Claim (iv) follows from the trivial identity  $e^x = 1 + (e^x - 1)$  and the elementary inequality  $e^{-x} \geq 1 - (e^x - 1)$ . For Claim (v), we may use the functional equation  $\Gamma = \Gamma^*$  to assume that  $\text{Im}(z) \geq 0$ . We use equations (1.13), (3.1), (3.14) and (3.15) of [3] to obtain the Stirling approximation

$$\Gamma(z) = \sqrt{2\pi} \exp\left(\left(z - \frac{1}{2}\right) \log z - z\right) \left(1 + \frac{1}{12z} + R_2(z)\right)$$

where the remainder  $R_2(z)$  obeys the bound

$$|R_2(z)| \leq (2\sqrt{2} + 1) \frac{C_2 \Gamma(2)}{(2\pi)^3 |z|^2}$$

for  $\text{Re}(z) \geq 0$  and

$$|R_2(z)| \leq (2\sqrt{2} + 1) \frac{C_2 \Gamma(2)}{(2\pi)^3 |z|^2 |1 - e^{2\pi iz}|}$$

for  $\text{Re}(z) \leq 0$ , where  $C_2$  is the constant

$$C_2 := \frac{1}{2} (1 + \zeta(2)) = \frac{1}{2} \left(1 + \frac{\pi^2}{6}\right).$$

In the latter case, we have  $\text{Im}(z) \geq 1$  by hypothesis, and hence  $|1 - e^{2\pi iz}| \geq 1 - e^{-2\pi}$ . We conclude that in all ranges of  $z$  of interest, we have

$$|R_2(z)| \leq (2\sqrt{2} + 1) \frac{C_2 \Gamma(2)}{(2\pi)^3 |z|^2 (1 - e^{-2\pi})} \leq \frac{0.0205}{|z|^2}$$

and hence by Claim (i)

$$\Gamma(z) = \sqrt{2\pi} \exp\left(\left(z - \frac{1}{2}\right) \log z - z\right) \left(1 + O_{\leq}\left(\frac{1}{12(|z| - 0.246)}\right)\right)$$

and the claim then follows by Claim (ii).

For Claim (vi), it suffices to show that the function  $x \mapsto \frac{\log^a x}{(x-c)^b}$  is non-increasing for  $x \geq \exp(a/b)$ . Taking logarithms and differentiating, we wish to show that  $\frac{a}{x \log x} - \frac{b}{x-c} \leq 0$ . But this is clear since  $\frac{b}{x-c} \geq \frac{b}{x}$  and  $\log x \geq a/b$ .  $\square$

## 6. ESTIMATES FOR LARGE $x$

The objective of this section is to obtain effective approximations for the expression  $H_t(x + iy)$  in the region

$$(20) \quad 0 < t \leq \frac{1}{2}; \quad 0 \leq y \leq 1; \quad x \geq 200.$$

The bounds in this region are somewhat arbitrary; one could easily replace the constants  $\frac{1}{2}$ , 1, 200 here by other positive numbers, at the expense of altering many of the numerical constants in the effective estimates.

The strategy is to use the expansion (17), which turns out to be an effective approximation as long as  $x$  stays away from zero, so that quantities such as  $s + \sqrt{t}v + \frac{t}{2}\alpha_n$  or  $s + \sqrt{t}v + \frac{t}{2}\beta_N$ , with

$s = \frac{1+i(x+iy)}{2}$ , stay away from the real axis where the poles of  $\Gamma$  are located (and also where the error terms in the Riemann-Siegel approximation deteriorate).

Accordingly, we will need effective estimates on the functions  $r_{t,n}, R_{t,N}$  appearing in Section 4. We will treat these two functions separately.

**6.1. Estimation of  $r_{t,n}$ .** We will need the function

$$(21) \quad M_0(s) := \frac{1}{8} \frac{s(s-1)}{2} \pi^{-s/2} \sqrt{2\pi} \exp\left(\left(\frac{s}{2} - \frac{1}{2}\right) \log \frac{s}{2} - \frac{s}{2}\right)$$

defined for all  $s$  away from the negative axis. Clearly this function is non-vanishing for all such  $s$ . We may compute the logarithmic derivative

$$(22) \quad \alpha := \frac{M'_0}{M_0}$$

of this function as

$$(23) \quad \begin{aligned} \alpha(s) &= \frac{1}{s} + \frac{1}{s-1} - \frac{1}{2} \log \pi + \frac{1}{2} \log \frac{s}{2} - \frac{1}{2s} \\ &= \frac{1}{2s} + \frac{1}{s-1} + \frac{1}{2} \log \frac{s}{2\pi}. \end{aligned}$$

We can also compute one further derivative:

$$(24) \quad \alpha'(s) = -\frac{1}{2s^2} - \frac{1}{(s-1)^2} + \frac{1}{2s}.$$

If  $\text{Im}(s) > 3$ , we conclude in particular that

$$(25) \quad \begin{aligned} \alpha'(s) &= O_{\leq}\left(\frac{1}{2\text{Im}(s)^2}\right) + O_{\leq}\left(\frac{1}{\text{Im}(s)^2}\right) + O_{\leq}\left(\frac{1}{2\text{Im}(s)}\right) \\ &= O_{\leq}\left(\frac{1}{2(\text{Im}(s) - 3)}\right) \end{aligned}$$

thanks to Lemma 5.1(i). Finally, we introduce the more general functions

$$(26) \quad M_t(s) := \exp\left(\frac{t}{4}\alpha(s)^2\right)M_0(s)$$

for any  $t \geq 0$ , as well as the sequence

$$(27) \quad b_n^t := \exp\left(\frac{t}{4} \log^2 n\right).$$

**Proposition 6.1** (Estimate for  $r_{t,n}$ ). *Let  $\sigma$  be real, let  $T > 10$ , let  $n$  be a positive integer, and let  $0 < t \leq 1/2$ . Then*

$$r_{t,n}(\sigma + iT) = M_t(\sigma + iT) \frac{b_n^t}{n^{\sigma+iT+\frac{t}{2}\alpha(\sigma+iT)}} (1 + O_{\leq}(\varepsilon_{t,n}(\sigma + iT)))$$

where

$$(28) \quad \varepsilon_{t,n}(\sigma + iT) := \exp\left(\frac{\frac{t^2}{8}|\alpha(\sigma + iT) - \log n|^2 + \frac{t}{4} + \frac{1}{6}}{T - 3.33}\right) - 1,$$

*Proof.* From (15), (21) and Lemma 5.1(v) one has

$$r_{0,n}(s) = M_0(s)n^{-s} \exp\left(O_{\leq}\left(\frac{1}{6(|s| - 0.66)}\right)\right)$$

whenever  $\text{Im}(s) > 2$ . Let  $\alpha_n$  denote the quantity

$$(29) \quad \alpha_n := \alpha(\sigma + iT) - \log n;$$

this is the logarithmic derivative of  $M(s)n^{-s}$  at  $s = \sigma + iT$ . From (18) we have

$$\begin{aligned} r_{t,n}(\sigma + iT) &= \exp\left(-\frac{t}{4}\alpha_n^2\right) \int_{\mathbb{R}} \exp(-\sqrt{t}v\alpha_n) M_0\left(\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n\right) \times \\ &\quad \times \exp\left(-\left(\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n\right) \log n + O_{\leq}\left(\frac{1}{6(|\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n| - 0.66)}\right)\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv. \end{aligned}$$

By (23) and the hypothesis  $T \geq 10$ , the imaginary part of  $\alpha_n$  may be lower bounded by

$$\text{Im}(\alpha_n) \geq -\frac{1}{2T} - \frac{1}{T} \geq -0.15;$$

since  $t \leq 1/2$ , we conclude that  $\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n$  has imaginary part at least  $T - 0.08$ . Thus

$$\begin{aligned} r_{t,n}(s) &= \exp\left(-\frac{t}{4}\alpha_n^2\right) \int_{\mathbb{R}} \exp(-\sqrt{t}v\alpha_n) M_0\left(\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n\right) \times \\ &\quad \times \exp\left(-\left(\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n\right) \log n + O_{\leq}\left(\frac{1}{6(T - 0.74)}\right)\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv. \end{aligned}$$

From (25) we have

$$\alpha'(s) = O_{\leq}\left(\frac{1}{2(T - 3.08)}\right)$$

for all  $s$  on the line segment between  $\sigma + iT$  and  $\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n$ . Applying Taylor's theorem with remainder to a branch of the complex logarithm of  $M_0$ , we conclude that

$$M_0(\sigma + iT + \sqrt{t}v + \frac{t}{2}\alpha_n) = M_0(\sigma + iT) \exp\left(\alpha(\sigma + iT)(\sqrt{t}v + \frac{t}{2}\alpha_n) + O_{\leq}\left(\frac{|\sqrt{t}v + \frac{t}{2}\alpha_n|^2}{4(T - 3.08)}\right)\right).$$

Inserting this estimate, writing  $\alpha(\sigma + iT) = \alpha_n + \log n$ , estimating  $\frac{1}{6(T - 0.74)}$  by  $\frac{1}{6(T - 3.08)}$  and  $|\sqrt{t}v + \frac{t}{2}\alpha_n|^2$  by  $2tv^2 + \frac{t^2}{2}|\alpha_n|^2$ , and simplifying, we conclude that

$$\begin{aligned} r_{t,n}(s) &= M_0(\sigma + iT) \exp\left(\frac{t}{4}\alpha_n^2 - (\sigma + iT) \log n\right) \times \\ &\quad \times \int_{\mathbb{R}} \exp\left(O_{\leq}\left(\frac{\frac{t}{2}v^2 + \frac{t^2}{8}|\alpha_n|^2 + \frac{1}{6}}{T - 3.08}\right)\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv. \end{aligned}$$

Using (29), (26), (27) we see that

$$M_0(\sigma + iT) \exp\left(\frac{t}{4}\alpha_n^2 - (\sigma + iT) \log n\right) = M_t(\sigma + iT) \frac{b_n^t}{n^{\sigma + iT + \frac{t}{2}\alpha(\sigma + iT)}}$$

and so it suffices to show that

$$\int_{\mathbb{R}} \exp\left(O_{\leq}\left(\frac{\frac{t}{2}v^2 + \frac{t^2}{8}|\alpha_n|^2 + \frac{1}{6}}{T - 3.08}\right)\right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv = 1 + O\left(\exp\left(\frac{\frac{t^2}{8}|\alpha_n|^2 + \frac{t}{4} + \frac{1}{6}}{T - 3.33}\right) - 1\right).$$

Since  $\frac{1}{\sqrt{\pi}}e^{-v^2} dv$  integrates to one, and  $\frac{1}{T-3.08} \leq \frac{1}{T-3.33}$ , it suffices by Lemma 5.1(iv) to show that

$$(30) \quad \int_{\mathbb{R}} \exp\left(\frac{tv^2}{2(T-3.08)}\right) \frac{1}{\sqrt{\pi}}e^{-v^2} dv \leq \exp\left(\frac{t}{4(T-3.33)}\right).$$

Using (5), the left-hand side may be calculated exactly as

$$\left(1 - \frac{t}{2(T-3.08)}\right)^{-1/2}.$$

Applying Lemma 5.1(ii) and using the hypotheses  $t \leq 1/2$ ,  $T \geq 10$ , one has

$$1 - \frac{t}{2(T-3.08)} = \exp\left(O_{\leq}\left(\frac{t}{2(T-3.33)}\right)\right)$$

and the claim follows.  $\square$

**6.2. Estimation of  $R_{t,N}$ .** We begin with the following estimates of Arias de Reyna [1] on the term  $\int_{N \swarrow N+1} \frac{w^{-s} e^{i\pi w^2}}{e^{\pi i w} - e^{-\pi i w}}$  appearing in (16):

**Proposition 6.2.** *Let  $\sigma$  be real and  $T' > 0$ , and define the quantities*

$$(31) \quad s := \sigma + iT'$$

$$(32) \quad a := \sqrt{\frac{T'}{2\pi}}$$

$$(33) \quad N := \lfloor a \rfloor$$

$$(34) \quad p := 1 - 2(a - N)$$

$$(35) \quad U := \exp\left(-i\left(\frac{T'}{2} \log \frac{T'}{2\pi} - \frac{T'}{2} - \frac{\pi}{8}\right)\right).$$

*Let  $K$  be a positive integer. Then we have an expansion*

$$\int_{N \swarrow N+1} \frac{w^{-s} e^{i\pi w^2}}{e^{\pi i w} - e^{-\pi i w}} = (-1)^{N-1} U a^{-\sigma} \left( \sum_{k=0}^K \frac{C_k(p, \sigma)}{a^k} + RS_K(s) \right)$$

*where  $C_0(p, \sigma) = C_0(p)$  is independent of  $\sigma$  and is given explicitly by the formula*

$$(36) \quad C_0(p) := \frac{e^{\pi i(\frac{p^2}{2} + \frac{3}{8})} - i\sqrt{2} \cos \frac{\pi p}{2}}{2 \cos(\pi p)}$$

*(removing the singularities at  $p = \pm 1/2$ ), while for  $k \geq 1$  the  $C_k(p, \sigma)$  are quantities obeying the bounds*

$$(37) \quad |C_k(p, \sigma)| \leq \frac{\sqrt{2} 9^{\sigma} \Gamma(k/2)}{2\pi 2^k}$$

*for  $\sigma > 0$  and*

$$(38) \quad |C_k(p, \sigma)| \leq \frac{2^{\frac{1}{2}-\sigma}}{2\pi} \frac{\Gamma(k/2)}{2\pi((3-2\log 2)\pi)^{k/2}}$$

*for  $\sigma \leq 0$ , while the error term  $RS_K(s)$  is a quantity obeying the bounds*

$$(39) \quad |RS_K(s)| \leq \frac{1}{7} 2^{3\sigma/2} \frac{\Gamma((K+1)/2)}{(a/1.1)^{K+1}}$$

for  $\sigma \geq 0$ , and

$$(40) \quad |RS_K(s)| \leq \frac{1}{2} \left( \frac{9}{10} \right)^{\lceil -\sigma \rceil} \frac{\Gamma((K+1)/2)}{(a/1.1)^{K+1}}$$

if  $\sigma < 0$  and  $K + \sigma \geq 2$ .

*Proof.* This follows from [1, Theorems 3.1, 4.1, 4.2] combined with [1, (3.2), (5.2)]. The dependence of  $C_k(p, \sigma)$ ,  $k \geq 1$  on  $\sigma$  and the dependence of  $RS_K(s)$  on  $s$  is suppressed in [1], but can be discerned from the definitions of these quantities (and the related quantities  $g(\tau, z)$ ,  $P_k(z) = P_k(z, \sigma)$ ,  $Rg_K(\tau, z)$ ) in [1, (3.9), (3.10), (3.7), (3.6)].  $\square$

Note that  $p$  ranges in the interval  $[-1, 1]$ . One can show that

$$(41) \quad |C_0(p)| \leq \frac{1}{2}$$

for all  $p \in [-1, 1]$ ; this follows for instance from the  $n = 0$  case of [1, Theorem 6.1].

**Proposition 6.3** (Estimate for  $R_{i,N}$ ). *Let  $0 \leq \sigma \leq 1$ , let  $T \geq 100$ , and let  $0 < t \leq 1/2$ . Set*

$$T' := T + \frac{\pi t}{8}$$

*and then define  $a, N, p, U, C_0(p)$  using (32), (33), (35), (36). Then*

$$R_{i,N}(\sigma + iT) = (-1)^{N-1} U e^{\pi i \sigma / 4} \exp\left(\frac{t\pi^2}{64}\right) M_0(iT') (C_0(p) + O_{\leq}(\tilde{\varepsilon}(\sigma + iT)))$$

where

$$(42) \quad \tilde{\varepsilon}(\sigma + iT) := \left( \frac{0.397 \times 9^\sigma}{a - 0.125} + \frac{5}{3(T' - 3.33)} \right) \exp\left(\frac{3.49}{T' - 3.33}\right).$$

*Proof.* We apply (19) with  $\beta_N := \pi i / 4$  to obtain

$$R_{i,N}(\sigma + iT) = \exp\left(\frac{t\pi^2}{64}\right) \int_{\mathbb{R}} \exp\left(-\frac{\sqrt{t}\nu\pi i}{4}\right) R_{0,N}(\sigma + iT' + \sqrt{t}\nu) \frac{1}{\sqrt{\pi}} e^{-\nu^2} d\nu.$$

From (16) we have

$$R_{0,N}(\sigma + iT' + \sqrt{t}\nu) = \frac{1}{8} \frac{s_\nu(s_\nu - 1)}{2} \pi^{-s_\nu/2} \Gamma\left(\frac{s_\nu}{2}\right) (-1)^{N-1} U a^{-\sigma - \sqrt{t}\nu} \left( \sum_{k=0}^{K_\nu} \frac{C_k(p, \sigma + \sqrt{t}\nu)}{a^k} + RS_{K_\nu}(s_\nu) \right)$$

for any positive integer  $K_\nu$  that we permit to depend (in a measurable fashion) on  $\nu$ , where  $s_\nu := \sigma + iT' + \sqrt{t}\nu$ . From (21) and Lemma 5.1(v) we thus have

$$R_{0,N}(\sigma + iT' + \sqrt{t}\nu) = M_0(s_\nu) \exp\left(O_{\leq}\left(\frac{1}{12(T' - 0.33)}\right)\right) (-1)^{N-1} U a^{-\sigma - \sqrt{t}\nu} \left( \sum_{k=0}^{K_\nu} \frac{C_k(p, \sigma + \sqrt{t}\nu)}{a^k} + RS_{K_\nu}(s_\nu) \right).$$

From (25) and Taylor expansion of a logarithm of  $M$ , we have

$$M_0(s_\nu) = M_0(iT') \exp\left(\alpha(iT')(\sigma + \sqrt{t}\nu) + O_{\leq}\left(\frac{(\sigma + \sqrt{t}\nu)^2}{4(T' - 0.33)}\right)\right).$$

From (23), (32) one has

$$\alpha(iT') = O_{\leq}\left(\frac{1}{2T'}\right) + O_{\leq}\left(\frac{1}{T'}\right) + \frac{1}{2} \log \frac{iT'}{2\pi} = \log a + \frac{i\pi}{4} + O_{\leq}\left(\frac{3}{2T'}\right)$$

and hence (bounding  $\frac{3}{2T'}$  by  $\frac{6}{4(T'-0.33)}$ )

$$\alpha(iT')(\sigma + \sqrt{t}v) = (\sigma + \sqrt{t}v) \log a + \frac{\pi i \sigma}{4} + \frac{\sqrt{t}v \pi i}{4} + O_{\leq} \left( \frac{6|\sigma + \sqrt{t}v|}{4(T' - 0.33)} \right).$$

We conclude that

$$\begin{aligned} \exp \left( -\frac{\sqrt{t}v \pi i}{4} \right) R_{0,N}(\sigma + iT' + \sqrt{t}v) &= M_0(iT') \exp \left( O_{\leq} \left( \frac{(\sigma + \sqrt{t}v)^2 + 6|\sigma + \sqrt{t}v| + \frac{1}{3}}{4(T' - 0.33)} \right) \right) \times \\ &\times (-1)^{N-1} U e^{\pi i \sigma / 4} \left( \sum_{k=0}^{K_v} \frac{C_k(p, \sigma + \sqrt{t}v)}{a^k} + RS_{K_v}(s_v) \right). \end{aligned}$$

Bounding  $6|\sigma + \sqrt{t}v| \leq 3(\sigma + \sqrt{t}v)^2 + 3$ , we have

$$\frac{(\sigma + \sqrt{t}v)^2 + 6|\sigma + \sqrt{t}v| + \frac{1}{3}}{4(T' - 0.33)} \leq \frac{(\sigma + \sqrt{t}v)^2 + \frac{5}{6}}{T' - 0.33}.$$

Putting all this together, we obtain

$$\begin{aligned} R_{t,N}(\sigma + iT) &= (-1)^{N-1} U e^{\pi i \sigma / 4} \exp \left( \frac{t \pi^2}{64} \right) M_0(iT') \times \\ &\times \int_{\mathbb{R}} \exp \left( O_{\leq} \left( \frac{(\sigma + \sqrt{t}v)^2 + \frac{5}{6}}{T' - 0.33} \right) \right) \left( \sum_{k=0}^{K_v} \frac{C_k(p, \sigma + \sqrt{t}v)}{a^k} + RS_{K_v}(s_v) \right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv. \end{aligned}$$

We separate the  $k = 0$  term from the rest. By Lemma 5.1(iv) and the fact that  $\frac{1}{\sqrt{\pi}} e^{-v^2}$  integrates to one, we can write the above expression as

$$(43) \quad R_{t,N}(\sigma + iT) = (-1)^{N-1} U e^{\pi i \sigma / 4} \exp \left( \frac{t \pi^2}{64} \right) M_0(iT') (C_0(p)(1 + O_{\leq}(\epsilon)) + O_{\leq}(\delta))$$

where

$$\epsilon := \int_{\mathbb{R}} \left( \exp \left( \frac{(\sigma + \sqrt{t}v)^2 + \frac{5}{6}}{T' - 0.33} \right) - 1 \right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv$$

and

$$\delta := \int_{\mathbb{R}} \exp \left( \frac{(\sigma + \sqrt{t}v)^2 + \frac{5}{6}}{T' - 0.33} \right) \left( \sum_{k=1}^{K_v} \frac{|C_k(p, \sigma + \sqrt{t}v)|}{a^k} + |RS_{K_v}(s_v)| \right) \frac{1}{\sqrt{\pi}} e^{-v^2} dv.$$

Bounding  $(\sigma + \sqrt{t}v)^2 \leq 2\sigma^2 + 2tv^2$  and using (5) we obtain

$$\epsilon \leq \exp \left( \frac{2\sigma^2 + \frac{5}{6}}{T' - 0.33} \right) \left( 1 - \frac{2t}{T' - 0.33} \right)^{-1/2} - 1.$$

Applying Lemma 5.1(ii) and using the hypotheses  $t \leq 1/2$ ,  $T \geq 100$ , one has

$$1 - \frac{2t}{T' - 0.33} = \exp \left( O_{\leq} \left( \frac{2t}{T' - 3.33} \right) \right)$$

and hence

$$\epsilon \leq \exp \left( \frac{2\sigma^2 + t + \frac{5}{6}}{T' - 3.33} \right) - 1.$$

With  $t \leq 1/2$  and  $0 \leq \sigma \leq 1$ , one has  $2\sigma^2 + t + \frac{5}{6} \leq \frac{10}{3}$ . By the mean value theorem we then have

$$(44) \quad \epsilon \leq \frac{10}{3(T' - 3.33)} \exp\left(\frac{10}{3(T' - 3.33)}\right).$$

Now we work on  $\delta$ . Making the change of variables  $u := \sigma + \sqrt{t}v$ , we have

$$\delta = \int_{\mathbb{R}} \exp\left(\frac{u^2 + \frac{5}{6}}{T' - 0.33}\right) \left( \sum_{k=1}^{\tilde{K}_u} \frac{|C_k(p, u)|}{a^k} + |RS_{\tilde{K}_u}(u + iT')| \right) \frac{1}{\sqrt{\pi t}} e^{-(u-\sigma)^2/t} du,$$

where  $\tilde{K}_u$  is a positive integer parameter that can depend arbitrarily on  $u$  (as long as it is measurable, of course).

We choose  $\tilde{K}_u$  to equal 1 when  $u \geq 0$  and  $\max(\lfloor -\sigma \rfloor + 3, \lfloor \frac{T'}{\pi} \rfloor)$  when  $u < 0$ , so that Proposition 6.2 applies. The expression

$$\sum_{k=1}^{\tilde{K}_u} \frac{|C_k(p, u)|}{a^k} + |RS_{\tilde{K}_u}(u + iT')|$$

is then bounded by

$$(45) \quad \frac{\sqrt{2}}{2\pi} \frac{9^u \Gamma(1/2)}{2a} + \frac{1}{7} 2^{3u/2} \frac{\Gamma(1)}{(a/1.1)^2} \leq \frac{0.200 \times 9^u}{a} + \frac{0.173 \times 2^{3u/2}}{a^2}$$

for  $u \geq 0$  and

$$(46) \quad \sum_{1 \leq k \leq \tilde{K}_u} \frac{2^{\frac{1}{2}-u}}{2\pi} \frac{\Gamma(k/2)}{2\pi((3-2\log 2)\pi)^{k/2} a^k} + \frac{1}{2} (9/10)^{\lceil -u \rceil} \frac{\Gamma((\tilde{K}_u + 1)/2)}{(a/1.1)^{\tilde{K}_u+1}}$$

for  $u < 0$ . One can calculate that

$$\frac{2^{\frac{1}{2}}}{2\pi} \frac{1}{2\pi} \leq 0.036 \leq \frac{1}{2}$$

and

$$\frac{1}{((3-2\log 2)\pi)^{1/2}} \leq 0.445 \leq 1.1$$

and hence we can bound (46) by

$$0.0362^{-u} \sum_{1 \leq k \leq \frac{T'}{\pi}} (0.445)^k \frac{\Gamma(k/2)}{a^k} \frac{1}{2} 2^{-u} \sum_{\frac{T'}{\pi} \leq k \leq -u+4} \frac{\Gamma(k/2)}{(a/1.1)^k}.$$

For  $u \geq 0$ , we can estimate (45) by

$$0.2 \times 9^u \left( \frac{1}{a} + \frac{0.865}{a^2} \right) \leq \frac{0.2 \times 9^u}{a - 0.865}$$

thanks to Lemma 5.1(i). For  $u < 0$ , we observe that if  $k \leq 2a^2 = \frac{T'}{\pi}$  then

$$\frac{\Gamma(k+2/2)}{a^{k+2}} = \frac{k}{2a^2} \frac{\Gamma(k/2)}{a^k} \leq \frac{\Gamma(k/2)}{a^k}$$

and hence by the geometric series formula

$$\sum_{2 \leq k \leq \frac{T'}{\pi}, k \text{ even}} (0.445)^k \frac{\Gamma(k/2)}{a^k} \leq \frac{(0.445)^2}{1 - (0.445)^2} \frac{\Gamma(2/2)}{a} \leq \frac{0.247}{a^2}$$

and similarly

$$\sum_{3 \leq k \leq \frac{T'}{\pi}, k \text{ odd}} (0.445)^k \frac{\Gamma(k/2)}{a^k} \leq \frac{(0.445)^3}{1 - (0.445)^2} \frac{\Gamma(3/2)}{(a/1.1)^3} \leq \frac{0.098}{a^3}$$

and hence we can bound (46) by

$$0.0362^{-u} \left( \frac{0.445 \sqrt{\pi}}{a} + \frac{0.247}{a^2} + \frac{0.098}{a^3} \right) + \frac{1}{2} 2^{-u} \sum_{\frac{T'}{\pi} \leq k \leq -u+4} \frac{\Gamma(k/2)}{(a/1.1)^k}.$$

By Lemma 5.1(i) we have

$$0.036 \left( \frac{0.445 \sqrt{\pi}}{a} + \frac{0.247}{a^2} + \frac{0.098}{a^3} \right) \leq \frac{0.029}{a - 0.353}$$

and thus we can bound (46) by

$$\frac{0.029 \times 2^{-u}}{a - 0.353} + \frac{1}{2} 2^{-u} \sum_{\frac{T'}{\pi} \leq k \leq -u+4} (1.1)^k \frac{\Gamma(k/2)}{a^k}.$$

Putting this together, we conclude that

$$\sum_{k=1}^{\tilde{K}_u} \frac{|C_k(p, u)|}{a^k} + |RS_{\tilde{K}_u}(u + iT')| \leq \frac{0.2 \times 9^u}{a - 0.865} + \frac{0.029 \times 2^{-u}}{a - 0.353} + \frac{2^{-u}}{2} \sum_{\frac{T'}{\pi} \leq k \leq -u+4} (1.1)^k \frac{\Gamma(k/2)}{a^k}$$

for all  $u$  (positive or negative). We conclude that  $\delta \leq \delta_1 + \delta_2 + \delta_3$ , where

$$\begin{aligned} \delta_1 &:= \int_{\mathbb{R}} \exp\left(\frac{u^2 + \frac{5}{6}}{T' - 0.33}\right) \frac{0.2 \times 9^u}{a - 0.865} \frac{1}{\sqrt{\pi t}} e^{-(u-\sigma)^2/t} du \\ \delta_2 &:= \int_{\mathbb{R}} \exp\left(\frac{u^2 + \frac{5}{6}}{T' - 0.33}\right) \frac{0.029 \times 2^{-u}}{a - 1.25} \frac{1}{\sqrt{\pi t}} e^{-(u-\sigma)^2/t} du \\ (47) \quad \delta_3 &:= \int_{\mathbb{R}} \exp\left(\frac{u^2 + \frac{5}{6}}{T' - 0.33}\right) \frac{2^{-u}}{2} \sum_{\frac{T'}{\pi} \leq k \leq -u+4} (1.1)^k \frac{\Gamma(k/2)}{a^k} \frac{1}{\sqrt{\pi t}} e^{-(u-\sigma)^2/t} du. \end{aligned}$$

For  $\delta_1$ , we translate  $u$  by  $\sigma$  to obtain

$$\delta_1 = \frac{0.2 \times 9^\sigma}{a - 0.865} \int_{\mathbb{R}} \exp\left(\frac{u^2 + 2\sigma u + \sigma^2 + \frac{5}{6}}{T' - 0.33} + 2u \log 3\right) \frac{1}{\sqrt{\pi t}} e^{-u^2/t} du$$

and hence by (5)

$$(48) \quad \delta_1 = \frac{0.2 \times 9^\sigma}{a - 0.865} \exp\left(\frac{\sigma^2 + \frac{5}{6}}{T' - 0.33} + \frac{t(\log 3 + \frac{\sigma}{T' - 0.33})^2}{1 - \frac{t}{T' - 0.33}}\right) \left(1 - \frac{t}{T' - 0.33}\right)^{-1/2}.$$

One can write

$$(49) \quad \frac{1}{1 - \frac{t}{T' - 0.33}} = 1 + \frac{t}{T' - 0.33 - t} \leq 1 + \frac{t}{T' - 0.83}$$

while by Lemma 5.1(ii) we have

$$(50) \quad 1 - \frac{t}{T' - 0.33} = \exp\left(O_{\leq}\left(\frac{t}{T' - 0.33 - t}\right)\right) = \exp\left(O_{\leq}\left(\frac{t}{T' - 0.83}\right)\right).$$



We conclude that

$$\delta_1 \leq \frac{0.2 \times 9^\sigma}{a - 0.865} \exp \left( \frac{5 + 3t + 6\sigma^2}{6(T' - 0.83)} + t \left( \log 3 + \frac{\sigma}{T' - 0.33} \right)^2 \left( 1 + \frac{t}{T' - 0.83} \right) \right).$$

From Lemma 5.1(i) and the hypothesis  $0 \leq \sigma \leq 1$ , we have

$$\begin{aligned} \left( \log 3 + \frac{\sigma}{T' - 0.33} \right)^2 &\leq (\log^2 3) \left( 1 + \frac{2\sigma / \log 3}{T' - 0.33 - \frac{\sigma}{2 \log 3}} \right) \\ &\leq (\log^2 3) \left( 1 + \frac{2\sigma / \log 3}{T' - 0.83} \right) \end{aligned}$$

and therefore by a further application of Lemma 5.1(i)

$$\begin{aligned} \left( \log 3 + \frac{\sigma}{T' - 0.33} \right)^2 \left( 1 + \frac{t}{T' - 0.83} \right) &\leq \log^2 3 \left( 1 + \frac{\frac{2\sigma}{\log 3} + t}{T' - 0.83 - \frac{2\sigma t / \log 3}{2\sigma / \log 3 + t}} \right) \\ &\leq \log^2 3 \left( 1 + \frac{\frac{2\sigma}{\log 3} + t}{T' - 0.83 - t} \right) \\ &\leq \log^2 3 \left( 1 + \frac{\frac{2\sigma}{\log 3} + t}{T' - 1.33} \right) \end{aligned}$$

and thus

$$\delta_1 \leq \frac{0.2 \times 9^\sigma \exp(t \log^2 3)}{a - 0.865} \exp \left( \frac{5 + 3t + 6\sigma^2 + 12t\sigma \log 3 + 6t^2 \log^2 3}{6(T' - 1.33)} \right).$$

By repeating the proof of (48), we have

$$\delta_2 = \frac{0.029 \times 2^{-\sigma}}{a - 0.353} \exp \left( \frac{\sigma^2 + \frac{5}{6}}{T' - 0.33} + \frac{t \left( -\log \sqrt{2} + \frac{\sigma}{T' - 0.33} \right)^2}{1 - \frac{t}{T' - 0.33}} \right) \left( 1 - \frac{t}{T' - 0.33} \right)^{-1/2}.$$

We can bound  $(-\log \sqrt{2} + \frac{\sigma}{T' - 0.33})^2$  by  $\log^2 \sqrt{2}$ . Using (49), (50) we thus have

$$\delta_2 \leq \frac{0.029 \times 2^{-\sigma} \exp(t \log^2 \sqrt{2})}{a - 0.353} \exp \left( \frac{5 + 3t + 6\sigma^2}{6(T' - 1.33)} \right).$$

With  $t \leq 1/2$  and  $0 \leq \sigma \leq 1$  one has

$$0.2 \exp(t \log^2 3) \leq 0.366$$

$$0.029 \exp(t \log^2 \sqrt{2}) \leq 0.031$$

$$\frac{5 + 3t + 6\sigma^2}{6} \leq \frac{5 + 3t + 6\sigma^2 + 12t\sigma \log 3 + 6t^2 \log^2 3}{6} \leq 3.49$$

and hence

$$\delta_1 \leq \frac{0.366 \times 9^\sigma}{a - 0.865} \exp \left( \frac{3.49}{T' - 1.33} \right)$$

and

$$\delta_2 \leq \frac{0.031 \times 2^{-\sigma}}{a - 0.353} \exp \left( \frac{3.49}{T' - 1.33} \right).$$

Now we turn to  $\delta_3$ , which will end up being extremely small compared to  $\delta_1$  or  $\delta_2$ . By (47) and the Fubini-Tonelli theorem, we have

$$\delta_3 = \frac{1}{2\sqrt{\pi t}} \sum_{k \geq \frac{T'}{2.2\pi}} (1.1)^k \frac{\Gamma(k/2)}{a^k} \int_{-\infty}^{4-k} \exp\left(\frac{u^2 + \frac{5}{6}}{T' - 0.33} - \frac{(u - \sigma)^2}{t} - u \log 2\right) du.$$

Since  $u \leq 4 - k$ ,  $k \geq \frac{T'}{2.2\pi}$ , and  $T' \geq T \geq 100$ , we have  $k \geq 14$  and  $u \leq -10$ ; since  $\sigma \geq 0$ , we may thus lower bound  $(u - \sigma)^2/t$  by  $u^2/t$ . Since  $t \leq 1/2$ , we can upper bound  $\frac{u^2 + \frac{5}{6}}{T' - 0.33} - \frac{u^2}{t}$  by (say)  $-\frac{u^2}{2t}$ , thus

$$\delta_3 \leq \frac{1}{2\sqrt{\pi t}} \sum_{k \geq \frac{T'}{2.2\pi}} (1.1)^k \frac{\Gamma(k/2)}{a^k} \int_{-\infty}^{4-k} e^{-\frac{u^2}{2t} - u \log 2} du.$$

We can bound  $e^{-\frac{u^2}{2t}} \leq e^{\frac{(k-4)u}{2t}}$ , in the range of integration and thus

$$\int_{-\infty}^{4-k} e^{-\frac{u^2}{2t} - u \log 2} du \leq \frac{1}{\frac{k-4}{2t} - \log 2} e^{-\frac{(k-4)^2}{2t} + (k-4) \log 2} \leq \frac{1}{\frac{k-4}{2t} - \log 2} e^{-(k-4)^2 + (k-4) \log 2},$$

bounding

$$\frac{k-4}{2t} - \log 2 = \frac{k-4-2t \log 2}{2t} \geq \frac{k-6}{2t}$$

we conclude that

$$\delta_3 \leq \frac{\sqrt{t}}{\sqrt{\pi}} \sum_{k \geq \frac{T'}{2.2\pi}} (1.1)^k \frac{\Gamma(k/2)}{(k-6)a^k} e^{-(k-4)^2 + (k-4) \log 2}.$$

For  $k \geq 14$  one can easily verify that  $(1.1)^k \Gamma(k/2) e^{-(k-4)^2 + (k-4) \log 2} \leq 10^{-30}$ ; discarding the  $\frac{\sqrt{t}}{\sqrt{\pi}}$  and  $\frac{1}{k-6}$  factors we thus have

$$\delta_3 \leq \sum_{k \geq 14} \frac{10^{-30}}{a^k} \leq \frac{2 \times 10^{-30}}{a^{14}}$$

(say). Since

$$\frac{0.031 \times 2^{-\sigma}}{a - 0.353} + \frac{2 \times 10^{-30}}{a^{14}} \leq \frac{0.031 \times 2^{-\sigma}}{a - 0.865}$$

we thus have

$$\delta \leq \delta_1 + \delta_2 + \delta_3 \leq \frac{0.366 \times 9^\sigma + 0.031 \times 2^{-\sigma}}{a - 0.865} \exp\left(\frac{3.49}{T' - 1.33}\right).$$

Inserting this and (44), (41) into (43), and crudely bounding  $2^{-\sigma}$  by  $9^\sigma$ , we obtain the claim.  $\square$

**6.3. Combining the estimates.** Combining Propositions 6.1, 6.3 with (17) and the triangle inequality (and noting that  $M_0 = M_0^*$ ,  $M_t = M_t^*$  and  $\alpha = \alpha^*$ , and that  $U$  has magnitude 1), we conclude the following “ $A + B - C$  approximation to  $H_t$ ”:

**Corollary 6.4** ( $A + B - C$  approximation). *Let  $t, x, y$  obey (20). Set*

$$T' := \frac{x}{2} + \frac{\pi t}{8}$$

and then define  $a, N, p, U, C_0(p)$  using (32), (33), (35), (36). Define the quantities

$$\begin{aligned} A(x+iy) &:= M_t \left( \frac{1-y+ix}{2} \right) \sum_{n=1}^N \frac{b_n^t}{n^{\frac{1-y+ix}{2} + \frac{t}{2}\alpha(\frac{1-y+ix}{2})}} \\ B(x+iy) &:= M_t \left( \frac{1+y-ix}{2} \right) \sum_{n=1}^N \frac{b_n^t}{n^{\frac{1+y-ix}{2} + \frac{t}{2}\alpha(\frac{1+y-ix}{2})}} \\ C(x+iy) &:= 2e^{-\pi i y/8} (-1)^N \exp\left(\frac{t\pi^2}{64}\right) \operatorname{Re}(M_0(iT')C_0(p)Ue^{\pi i/8}) \end{aligned}$$

where  $M_0, b_n^t$  were defined in (26), (27). Then

$$H_t(x+iy) = A(x+iy) + B(x+iy) - C(x+iy) + O_{\leq}(E_A(x+iy) + E_B(x+iy) + E_C(x+iy))$$

where

$$\begin{aligned} E_A(x+iy) &:= |M_t \left( \frac{1-y+ix}{2} \right)| \sum_{n=1}^N \frac{b_n^t}{n^{\frac{1-y}{2} + \frac{t}{2}\operatorname{Re}\alpha(\frac{1-y+ix}{2})}} \varepsilon_{t,n} \left( \frac{1-y+ix}{2} \right) \\ E_B(x+iy) &:= |M_t \left( \frac{1+y+ix}{2} \right)| \sum_{n=1}^N \frac{b_n^t}{n^{\frac{1+y}{2} + \frac{t}{2}\operatorname{Re}\alpha(\frac{1+y+ix}{2})}} \varepsilon_{t,n} \left( \frac{1+y+ix}{2} \right) \\ E_C(x+iy) &:= \exp\left(\frac{t\pi^2}{64}\right) |M_0(iT')| (\tilde{\varepsilon} \left( \frac{1-y+ix}{2} \right) + \tilde{\varepsilon} \left( \frac{1+y+ix}{2} \right)) \end{aligned}$$

and  $\varepsilon_{t,n}, \tilde{\varepsilon}$  were defined in (28), (42).

In many cases one can use the cruder “ $A+B$ ” approximation that is immediate from the above corollary and (41):

**Corollary 6.5** ( $A+B$  approximation). *With the notation and hypotheses as in Corollary 6.4, we have*

$$H_t(x+iy) = A(x+iy) + B(x+iy) + O_{\leq}(E_A(x+iy) + E_B(x+iy) + E_{C,0}(x+iy))$$

where

$$E_{C,0}(x+iy) := \exp\left(\frac{t\pi^2}{64}\right) |M_0(iT')| (1 + \tilde{\varepsilon} \left( \frac{1-y+ix}{2} \right) + \tilde{\varepsilon} \left( \frac{1+y+ix}{2} \right)).$$

It will be convenient to normalize by the first term

$$(51) \quad B_0(x+iy) := M_t \left( \frac{1+y-ix}{2} \right)$$

of  $B(x+iy)$ . This expression is non-zero, and we conclude (under the hypotheses of Corollary 6.5) that

$$(52) \quad \frac{H_t(x+iy)}{B_0(x+iy)} = \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{s_*+\kappa}} + \frac{C(x+iy)}{B_0(x+iy)} + O_{\leq}(e_A + e_B + e_C)$$

and

$$(53) \quad \frac{H_t(x+iy)}{B_0(x+iy)} = \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{s_*+\kappa}} + O_{\leq}(e_A + e_B + e_{C,0})$$

where

$$(54) \quad \gamma := \gamma(x + iy) := \frac{M_t(\frac{1-y+ix}{2})}{M_t(\frac{1+y-ix}{2})}$$

$$(55) \quad s_* := s_*(x + iy) := \frac{1+y-ix}{2} + \frac{t}{2}\alpha\left(\frac{1+y-ix}{2}\right)$$

$$(56) \quad \kappa := \kappa(x + iy) := \frac{t}{2}\left(\alpha\left(\frac{1-y+ix}{2}\right) - \alpha\left(\frac{1+y+ix}{2}\right)\right)$$

$$(57) \quad e_A := e_A(x + iy) := |\gamma| \sum_{n=1}^N n^y \frac{b_n^t}{n^{\operatorname{Re}(s)+\operatorname{Re}(\kappa)}} \varepsilon_{t,n}\left(\frac{1-y+ix}{2}\right)$$

$$(58) \quad e_B := e_B(x + iy) := \sum_{n=1}^N \frac{b_n^t}{n^{\operatorname{Re}(s)}} \varepsilon_{t,n}\left(\frac{1+y+ix}{2}\right)$$

$$(59) \quad e_C := e_C(x + iy) := \frac{\exp\left(\frac{t^2}{64}\right) |M_0(iT')|}{|M_t(\frac{1+y+ix}{2})|} \left(\tilde{\varepsilon}\left(\frac{1-y+ix}{2}\right) + \tilde{\varepsilon}\left(\frac{1+y+ix}{2}\right)\right).$$

$$(60) \quad e_{C,0} := e_{C,0}(x + iy) := \frac{\exp\left(\frac{t^2}{64}\right) |M_0(iT')|}{|M_t(\frac{1+y+ix}{2})|} \left(1 + \tilde{\varepsilon}\left(\frac{1-y+ix}{2}\right) + \tilde{\varepsilon}\left(\frac{1+y+ix}{2}\right)\right).$$

It will be convenient to bound these quantities by expressions that depend on  $N$  rather than  $x$ . Observe that

$$N = \lfloor \sqrt{\frac{T'}{2\pi}} \rfloor = \lfloor \sqrt{\frac{x}{4\pi} + \frac{t}{16}} \rfloor$$

and hence

$$(61) \quad x_N \leq x < x_{N+1}$$

where

$$(62) \quad x_N := 4\pi N^2 - \frac{\pi t}{4}.$$

Similarly one has

$$2\pi N^2 \leq T' < 2\pi(N+1)^2.$$

**Proposition 6.6** (Estimates). *Let the notation and hypotheses be as in Corollary 6.4, and let  $\gamma, \kappa, e_A, e_B, e_{C,0}, e_C$  be defined by (54)-(60).*

(i) *One has*

$$|\gamma| \leq e^{0.02y} \left(\frac{x}{4\pi}\right)^{-y/2}$$

(ii) *One has*

$$\operatorname{Re} s_* \geq \frac{1+y}{2} + \frac{t}{4} \log \frac{x}{4\pi} - \frac{(1-3y + \frac{4y(1+y)}{x^2})_+ t}{2x^2}.$$

(iii) *One has*

$$\kappa = O_{\leq} \left( \frac{ty}{2(x-6)} \right).$$

(iv) *One has*

$$e_A \leq |\gamma| N^{|k|} \sum_{n=1}^N n^y \frac{b_n^t}{n^{\operatorname{Re}(s_*)}} \left( \exp \left( \frac{\frac{t^2}{32} \log^2 \frac{x}{4\pi n^2} + 0.313}{\frac{x}{2} - 3.33} \right) - 1 \right).$$

(v) *One has*

$$e_B \leq \sum_{n=1}^N \frac{b_n^t}{n^{\operatorname{Re}(s_*)}} \left( \exp \left( \frac{\frac{t^2}{32} \log^2 \frac{x}{4\pi n^2} + 0.313}{\frac{x}{2} - 3.33} \right) - 1 \right).$$

(vi) *One has*

$$e_C \leq \left( \frac{x}{4\pi} \right)^{-\frac{1+y}{4}} \exp \left( -\frac{t}{16} \log^2 \frac{x}{4\pi} + \frac{3|\log \frac{x}{4\pi} + i\frac{\pi}{2}| + 3.58}{x - 8.52} \right) \left( \frac{1.24 \times (3^y + 3^{-y})}{N - 0.125} + \frac{6.92}{x - 6.66} \right).$$

and

$$e_{C,0} \leq \left( \frac{x}{4\pi} \right)^{-\frac{1+y}{4}} \exp \left( -\frac{t}{16} \log^2 \frac{x}{4\pi} + \frac{3|\log \frac{x}{4\pi} + i\frac{\pi}{2}| + 3.58}{x - 8.52} \right) \left( 1 + \frac{1.24 \times (3^y + 3^{-y})}{N - 0.125} + \frac{6.92}{x - 6.66} \right).$$

*Proof.* From the mean value theorem we have

$$\log \gamma = -y \frac{d}{d\sigma} \log \left| M_t \left( \sigma + \frac{ix}{2} \right) \right|$$

for some  $\frac{1-y}{2} \leq \sigma \leq \frac{1+y}{2}$ . From (22), (26) we have

$$\frac{d}{d\sigma} \log \left| M_t \left( \sigma + \frac{ix}{2} \right) \right| = \operatorname{Re} \left( \frac{t}{2} \alpha \left( \sigma + \frac{ix}{2} \right) \alpha' \left( \sigma + \frac{ix}{2} \right) + \alpha \left( \sigma + \frac{ix}{2} \right) \right).$$

From (25) one has

$$(63) \quad \alpha' \left( \sigma + \frac{ix}{2} \right) = O_{\leq} \left( \frac{1}{x-6} \right)$$

and from Taylor expansion we also have

$$\alpha \left( \sigma + \frac{ix}{2} \right) = \alpha \left( \frac{ix}{2} \right) + O_{\leq} \left( \frac{\sigma}{x-6} \right);$$

from (23) one has

$$\alpha \left( \frac{ix}{2} \right) = O_{\leq} \left( \frac{1}{x} \right) + O_{\leq} \left( \frac{1}{x} \right) + \frac{1}{2} \log \frac{ix}{4\pi} = \frac{1}{2} \log \frac{x}{4\pi} + i\frac{\pi}{4} + O_{\leq} \left( \frac{2}{x} \right)$$

and hence

$$(64) \quad \alpha \left( \sigma + \frac{ix}{2} \right) = \frac{1}{2} \log \frac{x}{4\pi} + i\frac{\pi}{4} + O_{\leq} \left( \frac{2+\sigma}{x-6} \right).$$

Inserting these bounds, we conclude that

$$\log \gamma = -y \operatorname{Re} \left( \left( \frac{1}{2} \log \frac{x}{4\pi} + i\frac{\pi}{4} + O_{\leq} \left( \frac{2+\sigma}{x-6} \right) \right) \left( 1 + O_{\leq} \left( \frac{t}{2(x-6)} \right) \right) \right).$$

Expanding this out, we have

$$\log \gamma = -y \left( \frac{1}{2} \log \frac{x}{4\pi} + O_{\leq} \left( \frac{2+\sigma + \frac{t}{4} \log \frac{x}{4\pi} + \frac{t\pi}{8} + \frac{t(2+\sigma)}{2(x-6)}}{x-6} \right) \right).$$

In the region (20), which implies that  $0 \leq \sigma \leq 1$ , we have

$$2 + \sigma + \frac{t\pi}{8} + \frac{t(2 + \sigma)}{2(x - 6)} \leq 3.21$$

and thus

$$\log \gamma \leq -\frac{y}{2} \log \frac{x}{4\pi} + y \frac{\frac{t}{4} \log \frac{x}{4\pi} + 3.21}{x - 6}.$$

The function  $x \mapsto \frac{\log \frac{x}{4\pi}}{x-6}$  is decreasing for  $x \geq 200$  thanks to Lemma 5.1(vi), hence

$$y \frac{\frac{t}{4} \log \frac{x}{4\pi} + 3.21}{x - 6} \leq y \frac{\frac{t}{4} \log \frac{200}{4\pi} + 3.21}{200 - 6} \leq 0.02y.$$

Claim (i) follows. We remark that one can improve the  $e^{0.02y}$  factor here by Taylor expanding  $\alpha$  to second order rather than first order, but we will not need to do so here.

To prove claim (ii), it suffices by (31) to show that

$$\operatorname{Re} \alpha \left( \frac{1 + y - ix}{2} \right) \geq \frac{1}{2} \log \frac{x}{4\pi} - \frac{(1 - 3y)_+}{x^2} - \frac{4y(1 + y)}{x^4}.$$

By (23) one has

$$\operatorname{Re} \alpha \left( \frac{1 + y - ix}{2} \right) = \frac{1 + y}{(1 + y)^2 + x^2} - \frac{2(1 - y)}{(1 - y)^2 + x^2} + \frac{1}{2} \log \frac{\sqrt{(1 + y)^2 + x^2}}{4\pi}.$$

We bound  $\sqrt{(1 + y)^2 + x^2} \geq x$  and calculate

$$\begin{aligned} \frac{1 + y}{(1 + y)^2 + x^2} - \frac{2(1 - y)}{(1 - y)^2 + x^2} &= -\frac{1 - 3y}{(1 + y)^2 + x^2} - \frac{4y(1 + y)}{((1 + y)^2 + x^2)((1 - y)^2 + x^2)} \\ &\geq -\frac{1 - 3y + \frac{4y(1 + y)}{x^2}}{(1 + y)^2 + x^2}. \end{aligned}$$

Lower bounding the numerator by its nonnegative part and then lower bounding  $(1 + y)^2 + x^2$  by  $x^2$ , we obtain the claim.

Claim (iii) is immediate from (63) and the fundamental theorem of calculus. Now we turn to (iv), (v). From (64) one has

$$\alpha \left( \frac{1 \pm y + ix}{2} \right) - \log n = \frac{1}{2} \log \frac{x}{4\pi n^2} + i \frac{\pi}{4} + O_{\leq} \left( \frac{3}{x - 6} \right)$$

for either choice of sign  $\pm$ . In particular, we have

$$(65) \quad \left| \alpha \left( \frac{1 \pm y + ix}{2} \right) - \log n \right|^2 = \frac{1}{4} \log^2 \frac{x}{4\pi n^2} + \frac{\pi^2}{16} + O_{\leq} \left( \frac{3 \left| \log \frac{x}{4\pi n^2} + i \frac{\pi}{2} \right|}{x - 6} + \frac{9}{(x - 6)^2} \right).$$

For any  $1 \leq n \leq N$ , we have

$$1 \leq n^2 \leq N^2 \leq a^2 = \frac{x + \frac{\pi t}{16}}{4\pi};$$

in the region (20), the right-hand side is certainly bounded by  $(\frac{x}{4\pi})^2$ , so that

$$\frac{4\pi}{x} \leq \frac{x}{4\pi n^2} \leq \frac{x}{4\pi}$$

and hence

$$\left| \log \frac{x}{4\pi n^2} + i \frac{\pi}{2} \right| \leq \left| \log \frac{x}{4\pi} + i \frac{\pi}{2} \right|.$$

In the region (20) we have  $x \geq 200$ , we see from Lemma 5.1(vi) (after squaring) that  $\frac{|\log \frac{x}{4\pi} + i\frac{\pi}{2}|}{x-6}$  is decreasing in  $x$ . Thus

$$\frac{\pi^2}{16} + \frac{3|\log \frac{x}{4\pi} + i\frac{\pi}{2}|}{x-6} + \frac{9}{(x-6)^2} \leq \frac{\pi^2}{16} + \frac{3|\log \frac{200}{4\pi} + i\frac{\pi}{2}|}{200-6} + \frac{9}{(200-6)^2} \leq 0.667.$$

Similarly, in (20) we also have

$$\frac{t^2}{8} \times 0.667 + \frac{t}{4} + \frac{1}{6} \leq 0.313.$$

We conclude from (28) that

$$\varepsilon_{t,n} \left( \frac{1 \pm y + ix}{2} \right) \leq \exp \left( \frac{\frac{t^2}{32} \log^2 \frac{x}{4\pi} + 0.313}{T - 3.33} \right) - 1.$$

Inserting this bound into (57), (58), we obtain claims (iv), (v).

Now we establish (vi). From (26) we have

$$\frac{\exp \left( \frac{t\pi^2}{64} \right) |M_0(iT')|}{|M_t(\frac{1+y+ix}{2})|} = \exp \left( \frac{t\pi^2}{64} - \frac{t}{4} \operatorname{Re} \alpha \left( \frac{1+y+ix}{2} \right) \right) \frac{|M_0(iT')|}{|M_0(\frac{1+y+ix}{2})|}.$$

Note that  $\frac{1+y+ix}{2} = iT' + \frac{1+y}{2} - \frac{\pi it}{8}$ . From (25) we see that  $|\alpha'(s)| \leq \frac{1}{x-6}$  for any  $s$  on the line segment between  $iT'$  and  $\frac{1+y+ix}{2}$ . From Taylor's theorem with remainder applied to a branch of  $\log M_0$ , we conclude that

$$\frac{|M_0(iT')|}{|M_0(\frac{1+y+ix}{2})|} = \exp \left( \operatorname{Re} \left( \left( -\frac{1+y}{2} + \frac{\pi it}{8} \right) \alpha(iT') \right) + O_{\leq} \left( \left| -\frac{1+y}{2} + \frac{\pi it}{8} \right|^2 \right) \right).$$

For  $0 \leq y \leq 1$  and  $0 < t \leq \frac{1}{2}$  we have

$$\frac{\left| -\frac{1+y}{2} + \frac{\pi it}{8} \right|^2}{2} \leq 0.52$$

and from (23) one has

$$\alpha(iT') = O_{\leq} \left( \frac{1}{2T'} \right) + O_{\leq} \left( \frac{1}{T'} \right) + \frac{1}{2} \log \frac{iT'}{2\pi} = \frac{1}{2} \log \frac{T'}{2\pi} + \frac{i\pi}{4} + O_{\leq} \left( \frac{3}{2T'} \right)$$

and hence

$$\frac{|M_0(iT')|}{|M_0(\frac{1+y+ix}{2})|} = \exp \left( -\frac{1+y}{4} \log \frac{T'}{2\pi} - \frac{t\pi^2}{32} + O_{\leq} \left( \frac{3\left| -\frac{1+y}{2} + \frac{\pi it}{8} \right|}{2T'} + \frac{0.52}{x-6} \right) \right).$$

Bounding  $\frac{1}{2T'} \leq \frac{1}{x-6}$  and  $\left| -\frac{1+y}{2} + \frac{\pi it}{8} \right| \leq 1.02$ , this becomes

$$\frac{|M_0(iT')|}{|M_0(\frac{1+y+ix}{2})|} = \left( \frac{T'}{2\pi} \right)^{-\frac{1+y}{4}} \exp \left( -\frac{t\pi^2}{32} + O_{\leq} \left( \frac{3.58}{x-6} \right) \right)$$

and hence

$$\frac{\exp\left(\frac{t\pi^2}{64}\right)|M_0(iT')|}{|M_t\left(\frac{1+y+ix}{2}\right)|} = \left(\frac{T'}{2\pi}\right)^{-\frac{1+y}{4}} \exp\left(-\frac{t\pi^2}{64} - \frac{t}{4}\operatorname{Re}\alpha\left(\frac{1+y+ix}{2}\right)^2 + O_\leq\left(\frac{3.58}{x-6}\right)\right).$$

By repeating the proof of (65) we have

$$\operatorname{Re}\alpha\left(\frac{1+y+ix}{2}\right)^2 = \frac{1}{4}\log^2\frac{x}{4\pi} - \frac{\pi^2}{16} + O_\leq\left(\frac{3|\log\frac{x}{4\pi} + i\frac{\pi}{2}|}{x-6} + \frac{9}{(x-6)^2}\right).$$

As before, in the region (20) we have

$$\frac{3|\log\frac{x}{4\pi} + i\frac{\pi}{2}|}{x-6} + \frac{9}{(x-6)^2} \leq \frac{3|\log\frac{x}{4\pi} + i\frac{\pi}{2}|}{x-6} + \frac{9}{(x-6)^2}$$

and thus

$$\begin{aligned} \frac{\exp\left(\frac{t\pi^2}{64}\right)|M_0(iT')|}{|M_t\left(\frac{1+y+ix}{2}\right)|} &= \left(\frac{T'}{2\pi}\right)^{-\frac{1+y}{4}} \exp\left(-\frac{t}{16}\log^2\frac{x}{4\pi} + O_\leq\left(\frac{3|\log\frac{x}{4\pi} + i\frac{\pi}{2}| + 3.58}{x-6} + \frac{9}{(x-6)^2}\right)\right) \\ &= \left(\frac{T'}{2\pi}\right)^{-\frac{1+y}{4}} \exp\left(-\frac{t}{16}\log^2\frac{x}{4\pi} + O_\leq\left(\frac{3|\log\frac{x}{4\pi} + i\frac{\pi}{2}| + 3.58}{x-8.52}\right)\right) \end{aligned}$$

thanks to Lemma 5.1(i). Finally, since  $T' \geq \frac{x}{2} \geq 100$  in (20), one has

$$\exp\left(\frac{3.49}{T' - 3.33}\right) \leq 1.037$$

and hence by (42)

$$\tilde{\varepsilon}\left(\frac{1+y+ix}{2}\right) \leq \frac{1.24 \times 3^{\pm y}}{a - 0.125} + \frac{1.73}{T' - 3.33}.$$

Hence

$$\tilde{\varepsilon}\left(\frac{1-y+ix}{2}\right) + \tilde{\varepsilon}\left(\frac{1+y+ix}{2}\right) \leq \frac{1.24 \times (3^y + 3^{-y})}{a - 0.125} + \frac{3.46}{T' - 3.33}$$

giving the claim (substituting  $T' = x/2$  and  $a \geq N$ ).  $\square$

### give table illustrating accuracy of approximation

## 7. BOUNDING DIRICHLET SERIES

In view of the approximation (53), it is of interest to obtain lower bounds for the quantity

$$(66) \quad X := \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{s_*+\kappa}}$$

for  $x, y, t$  in the region (20). By the triangle inequality (and the trivial identity  $|z| = |\bar{z}|$ ), we obtain the lower bound

$$|X| \geq \left( \left| \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} \right| - |\gamma| \left| \sum_{n=1}^N n^y \frac{b_n^t}{n^{s_*+\kappa}} \right| \right)_+.$$

It is thus of interest to obtain lower bounds for differences

$$(67) \quad \Delta := \left( \left| \sum_{n=1}^N \frac{\beta_n}{n^s} \right| - \left| \sum_{n=1}^N \frac{\alpha_n}{n^s} \right| \right)_+$$



of magnitudes of Dirichlet series for various coefficients  $\beta_n, \alpha_n$ . Our tools for this will be as follows.

**Lemma 7.1.** *Let  $N$  be a natural number, let  $s = \sigma + iT$  be a complex number for some real  $\sigma, T$ , and let  $\alpha_n, \beta_n$  be complex numbers for  $n = 1, \dots, N$  with  $\beta_1 = 1$ . Let  $\Delta$  denote the quantity (67).*

(i) *(Triangle inequality) We have*

$$\Delta \geq 1 - \alpha_1 - \sum_{n=2}^N \frac{|\alpha_n| + |\beta_n|}{n^\sigma}.$$

(ii) *(Refined triangle inequality) If the  $\alpha_n, \beta_n$  are all real and  $0 \leq \alpha_1 < 1$ , then we have*

$$\Delta \geq 1 - \alpha_1 - \sum_{n=2}^N \frac{\max(|\beta_n - \alpha_n|, \frac{1-\alpha_1}{1+\alpha_1}|\beta_n + \alpha_n|)}{n^\sigma}.$$

(iii) *(Dirichlet mollifier) If  $\lambda_1, \dots, \lambda_D$  are complex numbers, not all zero, then*

$$\Delta \geq \frac{\tilde{\Delta}}{\sum_{d=1}^D \frac{|\lambda_d|}{d^\sigma}}$$

where

$$\tilde{\Delta} := \left( \left| \sum_{n=1}^{DN} \frac{\tilde{\beta}_n}{n^s} \right| - \left| \sum_{n=1}^{DN} \frac{\tilde{\alpha}_n}{n^s} \right| \right)_+$$

and  $\tilde{\alpha}_n, \tilde{\beta}_n$  are the Dirichlet convolutions of  $\alpha_n, \beta_n$  with the  $\lambda_d$ :

$$\begin{aligned} \tilde{\alpha}_n &:= \sum_{1 \leq d \leq D: d|n} \lambda_d \alpha_{n/d} \\ \tilde{\beta}_n &:= \sum_{1 \leq d \leq D: d|n} \lambda_d \beta_{n/d}. \end{aligned}$$

*Proof.* The claim (i) is immediate from the triangle inequality.

Now we prove (ii). We may assume that the right-hand side is positive, as the claim is trivial otherwise. By a continuity argument (replacing  $\beta_n, \alpha_n$  for  $n \geq 2$  by  $t\beta_n, t\alpha_n$  with  $t$  increasing continuously from zero to one, noting that this only increases the right-hand side of the inequality) it suffices to verify the claim when  $\Delta$  is positive. In this case, we may write

$$\Delta = \left| \sum_{n=1}^N \frac{\beta_n - e^{i\theta} \alpha_n}{n^s} \right|$$

for some phase  $\theta$ . By the triangle inequality, we then have

$$\Delta \geq |1 - e^{i\theta} \alpha_1| - \sum_{n=2}^N \frac{|\beta_n - e^{i\theta} \alpha_n|}{n^\sigma}.$$

We factor out  $|1 - e^{i\theta} \alpha_1|$ , which is at least  $1 - \alpha_1$ , to obtain the lower bound

$$\Delta \geq (1 - \alpha_1) \left( 1 - \sum_{n=2}^N \frac{|\beta_n - e^{i\theta} \alpha_n| / |1 - e^{i\theta} \alpha_1|}{n^\sigma} \right).$$

By the cosine rule, we have

$$|\beta_n - e^{i\theta}\alpha_n|/|1 - e^{i\theta}\alpha_1|^2 = \frac{\beta_n^2 + \alpha_n^2 - 2\alpha_n\beta_n \cos \theta}{1 + \alpha_1^2 - 2\alpha_1 \cos \theta}.$$

This is a fractional linear function of  $\cos \theta$  with no poles in the range  $[-1, 1]$  of  $\cos \theta$ . Thus this function is monotone on this range and attains its maximum at either  $\cos \theta = +1$  or  $\cos \theta = -1$ . We conclude that

$$\frac{|\beta_n - e^{i\theta}\alpha_n|}{|1 - e^{i\theta}\alpha_1|} \leq \max\left(\frac{|\beta_n - \alpha_n|}{1 - \alpha_1}, \frac{|\beta_n + \alpha_n|}{1 + \alpha_1}\right)$$

and the claim follows.

For claim (iii), we recall the well-known relationship

$$\begin{aligned} \sum_{n=1}^{DN} \frac{\tilde{\alpha}_n}{n^s} &= \left( \sum_{d=1}^D \frac{\lambda_d}{d^s} \right) \left( \sum_{n=1}^N \frac{\alpha_n}{n^s} \right) \\ \sum_{n=1}^{DN} \frac{\tilde{\beta}_n}{n^s} &= \left( \sum_{d=1}^D \frac{\lambda_d}{d^s} \right) \left( \sum_{n=1}^N \frac{\beta_n}{n^s} \right) \end{aligned}$$

between Dirichlet convolution and Dirichlet series, which implies that

$$\tilde{\Delta} = \left| \sum_{d=1}^D \frac{\lambda_d}{d^s} \right| \Delta.$$

Since  $\tilde{\Delta}, \Delta$  are non-negative and

$$\left| \sum_{d=1}^D \frac{\lambda_d}{d^s} \right| \leq \sum_{d=1}^D \frac{|\lambda_d|}{d^{\sigma}},$$

the claim follows.  $\square$

Returning to the estimation of (66), we conclude from Lemma 7.1(i) with  $s$  replaced by  $s_*$ ,  $\beta_n$  replaced by  $b_n^t$ , and  $\alpha_n$  replaced by  $|\gamma|n^{y-\bar{\kappa}}b_n^t$  that

$$(68) \quad |X| \geq 1 - |\gamma| - \sum_{n=2}^N \frac{b_n^t}{n^{\sigma}} (1 + |\gamma|n^{y-\operatorname{Re}(\kappa)}),$$

where  $\sigma := \operatorname{Re} s_*$ . This rather crude bound will suffice when  $x$  is very large, particularly when combined with the estimates in Proposition 6.6. For smaller values of  $x$ , we would like to use parts (ii) and (iii) of Lemma 7.1. A technical difficulty arises because the quantity  $|\lambda|n^{y-\bar{\kappa}}b_n^t$  quantity need not be real, so that Lemma 7.1(ii) is not directly available. However, by writing

$$n^{-\bar{\kappa}} = 1 + O_{\leq}(n^{|\kappa|} - 1)$$

we see from the triangle inequality that

$$|X| \geq \left( \left| \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} \right| - \left| \sum_{n=1}^N \frac{|\gamma|b_n^t n^y}{n^{s_*}} \right| \right)_+ - |\gamma| \sum_{n=1}^N \frac{b_n^t (n^{|\kappa|} - 1)}{n^{\sigma-y}}.$$

Assuming for now that  $|\gamma| < 1$  (which in practice will follow from Proposition 6.6(i)), we can then apply Lemma 7.1(iii) follows by Lemma 7.1(ii) to conclude that

$$(69) \quad |X| \geq \frac{1 - \tilde{\alpha}_1 - \sum_{n=2}^N \frac{\max(|\tilde{\beta}_n - \tilde{\alpha}_n|, \frac{1-\tilde{\alpha}_1}{1+\tilde{\alpha}_1} |\tilde{\beta}_n + \tilde{\alpha}_n|)}{n^\sigma}}{\sum_{d=1}^D \frac{|\lambda_d|}{d^\sigma}} - |\gamma| \sum_{n=1}^N \frac{b_n^t (n^{|\kappa|} - 1)}{n^{\sigma-y}}$$

for any real numbers  $\lambda_1, \dots, \lambda_D$  with  $\lambda_1 = 1$ , where

$$\begin{aligned} \tilde{\alpha}_n &:= \sum_{1 \leq d \leq D: d|n} \lambda_d b_{n/d}^t |\gamma| n^y \\ \tilde{\beta}_n &:= \sum_{1 \leq d \leq D: d|n} \lambda_d b_{n/d}^t. \end{aligned}$$

In practice, it has proven convenient to use this estimate with Dirichlet mollifiers  $\sum_{d=1}^D \frac{\lambda_d}{d^s}$  that are Euler products of the form

$$\sum_{d=1}^D \frac{\lambda_d}{d^s} = \prod_{p \leq P} \left( 1 - \frac{b_p^t}{p^s} \right)$$

for some small prime  $P$ , where the product is over primes  $p$  up to  $P$ . For instance, if  $P = 3$ , then we would take  $D = 6$ ,  $\lambda_1 = 1$ ,  $\lambda_2 = -b_2^t$ ,  $\lambda_3 = -b_3^t$ ,  $\lambda_6 = b_2^t b_3^t$ , and all other  $\lambda_d$  vanishing. This choice achieves a substantial amount of cancellation in the  $\tilde{\beta}_n$  coefficients, which we have found to make the lower bound in (69) favorable. (For instance, it makes  $\tilde{\beta}_p$  vanish for all primes  $p \leq P$ ).

## 8. ESTIMATION FOR SMALL $x$

**the results in this section are currently unused.**

We now turn to the problem of estimating  $H_t(x + iy)$  (and its derivative) accurately when  $x$  is small (e.g.,  $x \leq 200$ ). Here, the approximations based on the Riemann-Siegel formula (14) do not seem to be particularly efficient. Instead, we return to the original definition (4). From the functional equation  $\Phi(-u) = -\Phi(u)$  we have

$$H_t(z) = \frac{1}{2} \int_{\mathbb{R}} e^{tu^2} \Phi(u) e^{izu} du$$

and hence on differentiation under the integral sign (which can be justified using for instance the Cauchy integral formula and Fubini's theorem) we have

$$H'_t(z) = \frac{i}{2} \int_{\mathbb{R}} e^{tu^2} u \Phi(u) e^{izu} du.$$

For any parameter  $0 \leq \theta < \pi/8$ , we may shift the contour to  $i\theta + \mathbb{R}$  and then use the even nature of  $\Phi$  to reflect the left half of that contour around the origin to obtain the identities

$$\begin{aligned} (70) \quad H_t(z) &= \frac{1}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) e^{izu} du + \frac{1}{2} \int_{i\theta-\infty}^{i\theta} e^{tu^2} \Phi(u) e^{izu} du \\ &= \frac{1}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) e^{izu} du + \frac{1}{2} \int_{-i\theta}^{-i\theta+\infty} e^{tu^2} \Phi(u) e^{-izu} du \\ &= \frac{1}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) e^{izu} du + \frac{1}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) e^{i\bar{z}u} du \end{aligned}$$

and similarly

$$(71) \quad H'_t(z) = \frac{i}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) u e^{izu} du - \frac{i}{2} \int_{i\theta}^{i\theta+\infty} e^{tu^2} \Phi(u) u e^{i\bar{z}u} du.$$

Since

$$\Phi(u) = \sum_{n=1}^{\infty} (2\pi^2 n^4 e^{9u} - 3\pi n^2 e^{5u}) \exp(-\pi n^2 e^{4u}),$$

we thus have for  $z = x + iy$  that

$$(72) \quad \begin{aligned} H_t(z) &= \frac{1}{2} \sum_{n=1}^{\infty} 2\pi^2 n^4 I_{t,\theta}(z - 9i, \pi n^2) - 3\pi n^2 I_{t,\theta}(z - 5i, \pi n^2) + 2\pi^2 n^4 \overline{I_{t,\theta}(\bar{z} - 9i, \pi n^2)} - 3\pi n^2 \overline{I_{t,\theta}(\bar{z} - 5i, \pi n^2)} \\ &= \frac{1}{2} \sum_{n=1}^{\infty} 2\pi^2 n^4 I_{t,\theta}(x - (9 - y)i, \pi n^2) - 3\pi n^2 I_{t,\theta}(x - (5 - y)i, \pi n^2) \\ &\quad + 2\pi^2 n^4 \overline{I_{t,\theta}(x - (9 + y)i, \pi n^2)} - 3\pi n^2 \overline{I_{t,\theta}(x - (5 + y)i, \pi n^2)} \end{aligned}$$

and similarly

$$(73) \quad \begin{aligned} H'_t(z) &= \frac{i}{2} \sum_{n=1}^{\infty} 2\pi^2 n^4 J_{t,\theta}(z - 9i, \pi n^2) - 3\pi n^2 J_{t,\theta}(z - 5i, \pi n^2) - 2\pi^2 n^4 \overline{J_{t,\theta}(\bar{z} - 9i, \pi n^2)} + 3\pi n^2 \overline{J_{t,\theta}(\bar{z} - 5i, \pi n^2)} \\ &= \frac{i}{2} \sum_{n=1}^{\infty} 2\pi^2 n^4 J_{t,\theta}(x - (9 - y)i, \pi n^2) - 3\pi n^2 J_{t,\theta}(x - (5 - y)i, \pi n^2) \\ &\quad - 2\pi^2 n^4 \overline{J_{t,\theta}(x - (9 + y)i, \pi n^2)} + 3\pi n^2 \overline{J_{t,\theta}(x - (5 + y)i, \pi n^2)} \end{aligned}$$

where  $I_{t,\theta}(b, \beta)$ ,  $J_{t,\theta}(b, \beta)$  denote the integrals

$$(74) \quad I_{t,\theta}(b, \beta) := \int_{i\theta}^{i\theta+\infty} \exp(tu^2 - \beta e^{4u} + ibu) du$$

$$(75) \quad J_{t,\theta}(b, \beta) := \int_{i\theta}^{i\theta+\infty} \exp(tu^2 - \beta e^{4u} + ibu) u du.$$

It is thus of interest to find good estimates for  $I_{t,\theta}(b, \beta)$ ,  $J_{t,\theta}(b, \beta)$  when  $\beta = \pi n^2$  for some  $n$ , and  $b = x - ia$  for  $a = 9 + y, 9 - y, 5 + y, 5 - y$ . It will be convenient to split

$$I_{t,\theta}(b, \beta) = I_{t,\theta,\leq X}(b, \beta) + I_{t,\theta,\geq X}(b, \beta)$$

for any cutoff  $X \geq 0$ , where

$$I_{t,\theta,\leq X}(b, \beta) := \int_{i\theta}^{i\theta+X} \exp(tu^2 - \beta e^{4u} + ibu) du$$

and

$$I_{t,\theta,\geq X}(b, \beta) := \int_{i\theta+X}^{i\theta+\infty} \exp(tu^2 - \beta e^{4u} + ibu) du.$$

One can similarly split

$$J_{t,\theta}(b, \beta) = J_{t,\theta,\leq X}(b, \beta) + J_{t,\theta,\geq X}(b, \beta)$$

**Proposition 8.1.** *Let  $b = x - ia$  for some  $x, a > 0$ , let  $0 \leq \theta < \frac{\pi}{8}$ , and let  $\beta > 0$  and  $X, t \geq 0$ .*

(i) *One has the upper bounds and*

$$(76) \quad |I_{t,\theta,\leq X}(b, \beta)| \leq \exp(-t\theta^2 - \theta x) \int_0^X \exp(tv^2 - \beta e^{4v} \cos(4\theta) + av) dv$$

and

$$(77) \quad |J_{t,\theta,\leq X}(b, \beta)| \leq \exp(-t\theta^2 - \theta x) \int_0^X \exp(tv^2 - \beta e^{4v} \cos(4\theta) + av)(\theta^2 + v^2)^{1/2} dv.$$

(ii) *If one has the hypothesis*

$$(78) \quad \beta e^{4X} \cos(4\theta) > \max\left(\frac{t}{2}, \frac{a + 2tX}{4}\right)$$

then one has

$$(79) \quad |I_{t,\theta,\geq X}(b, \beta)| \leq \frac{\exp(-t\theta^2 + tX^2 - \beta e^{4X} \cos(4\theta) - \theta x + aX)}{4\beta e^{4X} \cos(4\theta) - a - 2tX}$$

and similarly

$$(80) \quad |J_{t,\theta,\geq X}(b, \beta)| \leq \exp(-t\theta^2 + tX^2 - \beta e^{4X} \cos(4\theta) - \theta x + aX) \left( \frac{|X + i\theta|}{4\beta e^{4X} \cos(4\theta) - a - 2tX} + \frac{1}{(4\beta e^{4X} \cos(4\theta) - a - 2tX)^2} \right).$$

In particular, setting  $X = 0$ , we see that if

$$(81) \quad \beta \cos(4\theta) > \max\left(\frac{t}{2}, \frac{a}{4}\right)$$

then

$$(82) \quad |I_{t,\theta}(b, \beta)| \leq \frac{\exp(-t\theta^2 - \beta \cos(4\theta) - \theta x)}{4\beta \cos(4\theta) - a}$$

and similarly

$$(83) \quad |J_{t,\theta}(b, \beta)| \leq \exp(-t\theta^2 - \beta \cos(4\theta) - \theta x) \left( \frac{\theta}{4\beta \cos(4\theta) - a} + \frac{1}{(4\beta \cos(4\theta) - a)^2} \right).$$

In practice, the tail estimates (79), (80) become useful when  $\theta$  is close to  $\pi/8$  (to get the most gain out of the  $\exp(-\theta x)$  factor), and  $X$  is large. Similarly, (82), (83) become useful when  $\theta$  is close to  $\pi/8$  and  $\beta$  is large.

*Proof.* We first prove (i). Writing  $u = i\theta + v$  and  $b = x - ia$  we have

$$I_{t,\theta,\leq X}(b, \beta) = \int_0^X \exp(t(-\theta^2 + v^2 + 2i\theta v) - \beta e^{4v} \cos(4\theta) - i\beta e^{4v} \sin(4\theta) - \theta x + ivx + av + ia\theta) dv$$

and the claim (76) then follows from the triangle inequality. The claim (77) is proven similarly, with an additional factor of  $i\theta + v = O_{\leq}((\theta^2 + v^2)^{1/2})$  in the integrand.

Now we prove (ii). Writing  $u = i\theta + X + v$  and  $b = x - ia$ , we have

$$I_{t,\theta,\geq X}(b, \beta) = \int_0^\infty \exp(t(-\theta^2 + X^2 + 2Xv + v^2 + 2i\theta(X+v)) - \beta e^{4X} e^{4v} \cos(4\theta) - i\beta e^{4X} e^{4v} \sin(4\theta) - \theta x + ix(X+v) + aX + av + ia\theta) dv$$

and hence by the triangle inequality

$$|I_{t,\theta,\geq X}(b,\beta)| \leq \exp(-t\theta^2 + tX^2 - \beta e^{4X} \cos(4\theta) - \theta x + aX) \int_0^\infty \exp(tv^2 + 2tXv - \beta e^{4X}(e^{4v} - 1) \cos(4\theta) + av) dv.$$

Similarly one has

$$|J_{t,\theta,\geq X}(b,\beta)| \leq \exp(-t\theta^2 + tX^2 - \beta e^{4X} \cos(4\theta) - \theta x + aX) \int_0^\infty \exp(tv^2 + 2tXv - \beta e^{4X}(e^{4v} - 1) \cos(4\theta) + av)(|X + i\theta| + v) dv.$$

The exponent  $tv^2 + 2tXv - \beta e^{4X}(e^{4v} - 1) \cos(4\theta) + av$  equals 0 when  $v = 0$  and has derivative

$$2tv + 2tX - 4\beta e^{4X} \cos(4\theta) e^{4v} + a.$$

Note that  $e^{4v} \geq 1 + 4v$ . Hence by (81), this derivative will be bounded from below by the negative quantity  $-(4\beta e^{4X} \cos(4\theta) - a - 2tX)$ . This implies that

$$\exp(tv^2 - \beta e^{4X}(e^{4v} - 1) \cos(4\theta) + av) \leq \exp(-v(4\beta e^{4X} \cos(4\theta) - a - 2tX))$$

and the bounds (79), (80) follow from the standard integral identities

$$\begin{aligned} \int_0^\infty \exp(a - bv) dv &= \frac{\exp(a)}{b} \\ \int_0^\infty \exp(a - bv)v dv &= \frac{\exp(a)}{b^2}, \end{aligned}$$

valid for any  $b > 0$ . Setting  $X = 0$  we then obtain (82), (83).  $\square$

**Corollary 8.2.** *Let  $x, a, t > 0$  and  $0 \leq \theta < \frac{\pi}{4}$ , and let  $n_0$  be a natural number such that*

$$\pi n_0^2 \cos(4\theta) > \max\left(\frac{t}{2}, \frac{a}{4}\right).$$

*Then we have*

$$(84) \quad \left| \sum_{n \geq n_0} 2\pi^2 n^4 I_{t,\theta}(x - ia, \pi n^2) \right| \leq \frac{2\pi^2 \exp(-t\theta^2 - \theta x - \pi n_0^2 \cos(4\theta))}{4\pi n_0^2 \cos(4\theta) - a} \times (n_0^4 G_0(\alpha) + 4n_0^3 G_1(\alpha) + 6n_0^2 G_2(\alpha) + 4n_0 G_3(\alpha) + G_4(\alpha))$$

*and*

$$(85) \quad \left| \sum_{n \geq n_0} 3\pi n^2 I_{t,\theta}(x - ia, \pi n^2) \right| \leq \frac{3\pi \exp(-t\theta^2 - \theta x - \pi n_0^2 \cos(4\theta))}{4\pi n_0^2 \cos(4\theta) - a} \times (n_0^2 G_0(\alpha) + 2n_0 G_1(\alpha) + G_2(\alpha))$$

*where  $\alpha := e^{-\pi n_0 \cos(4\theta)}$ , and  $G_k(\alpha)$  are the generalised geometric series*

$$G_k(\alpha) := \sum_{m=0}^{\infty} m^k \alpha^m.$$

*Similarly one has*

$$(86) \quad \left| \sum_{n \geq n_0} 2\pi^2 n^4 J_{t,\theta}(x - ia, \pi n^2) \right| \leq 2\pi^2 \exp(-t\theta^2 - \pi n_0^2 \cos(4\theta) - \theta x) \left( \frac{\theta}{4\pi n_0^2 \cos(4\theta) - a} + \frac{1}{(4\pi n_0^2 \cos(4\theta) - a)^2} \right) \times (n_0^4 G_0(\alpha) + 4n_0^3 G_1(\alpha) + 6n_0^2 G_2(\alpha) + 4n_0 G_3(\alpha) + G_4(\alpha))$$

and

(87)

$$\left| \sum_{n \geq n_0} 3\pi n^2 J_{t,\theta}(x - ia, \pi n^2) \right| \leq 3\pi \exp(-t\theta^2 - \pi n_0^2 \cos(4\theta) - \theta x) \left( \frac{\theta}{4\pi n_0^2 \cos(4\theta) - a} + \frac{1}{(4\pi n_0^2 \cos(4\theta) - a)^2} \right) \\ \times (n_0^2 G_0(\alpha) + 2n_0 G_1(\alpha) + G_2(\alpha)).$$

We remark that  $G_k(\alpha)$  can be computed explicitly for small  $k$  and  $|\alpha| < 1$  as

$$\begin{aligned} G_0(\alpha) &= \frac{1}{1 - \alpha} \\ G_1(\alpha) &= \frac{\alpha}{(1 - \alpha)^2} \\ G_2(\alpha) &= \frac{\alpha^2 + \alpha}{(1 - \alpha)^3} \\ G_3(\alpha) &= \frac{\alpha^3 + 4\alpha^2 + \alpha}{(1 - \alpha)^4} \\ G_4(\alpha) &= \frac{\alpha^4 + 11\alpha^3 + 11\alpha^2 + \alpha}{(1 - \alpha)^5}. \end{aligned}$$

In general one has

$$G_k(\alpha) = \frac{\sum_{j=0}^{k-1} A(k, j) \alpha^{j+1}}{(1 - \alpha)^{k+1}}$$

for any  $k \geq 1$  and  $|\alpha| < 1$  where  $A(k, j)$  are the Eulerian numbers [16].

*Proof.* From (82) one has

$$\begin{aligned} \left| \sum_{n \geq n_0} 2\pi^2 n^4 I_{t,\theta}(x - ia, \pi n^2) \right| &\leq \sum_{n \geq n_0} 2\pi^2 n^4 \frac{\exp(-t\theta^2 - \pi n^2 \cos(4\theta) - \theta x)}{4\pi n^2 \cos(4\theta) - a} \\ &\leq \frac{2\pi^2 \exp(-t\theta^2 - \theta x)}{4\pi n_0^2 \cos(4\theta) - a - 2tX} \sum_{n \geq n_0} n^4 \exp(-\pi n n_0 \cos(4\theta)) \\ &= \frac{2\pi^2 \exp(-t\theta^2 - \theta x - \pi n_0^2 \cos(4\theta))}{4\pi n_0^2 \cos(4\theta) - a} \sum_{m=0}^{\infty} (n_0^4 + 4n_0^3 m + 6n_0^2 m^2 + 4n_0 m^3 + m^4) \alpha^m \\ &= \frac{2\pi^2 \exp(-t\theta^2 - \theta x - \pi n_0^2 \cos(4\theta))}{4\pi n_0^2 \cos(4\theta) - a} \\ &\quad \times (n_0^4 G_0(\alpha) + 4n_0^3 G_1(\alpha) + 6n_0^2 G_2(\alpha) + 4n_0 G_3(\alpha) + G_4(\alpha)) \end{aligned}$$

giving (84) as desired. The estimates (85), (86), (87) are proven similarly.  $\square$

## 9. A NEW UPPER BOUND FOR THE DE BRUIJN-NEWMAN CONSTANT

In this section we prove

**Theorem 9.1.** *We have  $\Lambda \leq 0.28$ .*

To prove this theorem, we apply Theorem 3.3 with  $t_0 := 0.2$ ,  $X := 5 \times 10^9$ , and  $y_0 = 0.4$ . It will thus suffice to verify the three conditions of the theorem. For the first claim - namely, that  $H_0(x + iy) \neq 0$  when  $0 \leq x \leq 5 \times 10^9$  and  $y \geq 0.4$  - this is immediate from this follows from (1), (2), and the result of Platt that all the non-trivial zeroes of  $\zeta$  with imaginary part between 0 and  $3.06 \times 10^{10}$  lie on the critical line  $\{\text{Re}(s) = 1/2\}$ .

For the remaining claims, one has to verify  $H_t(x + iy) \neq 0$  for various  $(x, y, t)$  in the region (20). To show that  $H_t(x + iy) \neq 0$ , it suffices by (53), (66) and the triangle inequality to show that

$$(88) \quad |X| > e_A + e_B + e_{C,0}$$

where

$$\begin{aligned} X &:= \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{\overline{s_*} + \kappa}} \\ b_n^t &:= \exp\left(\frac{t}{4} \log^2 n\right), \\ N &:= \lfloor \sqrt{\frac{x}{4\pi} + \frac{t}{16}} \rfloor \leq \left\lfloor \frac{x}{4\pi} \left(1 + \frac{\pi t}{4x}\right) \right\rfloor^{1/2} \end{aligned}$$

so in particular

$$(89) \quad \log \frac{x}{4\pi} \geq 2 \log N - \log\left(1 + \frac{\pi t}{4x}\right).$$

and  $s_*, \kappa, \gamma, e_A, e_B, e_{C_0}$  are given by (54)-(60).

Note that if  $x \geq 5 \times 10^9$  then  $N \geq 19947$ . We calculate a somewhat crude upper bound for the right-hand side of (88):

**Lemma 9.2.** *If  $x \geq 5 \times 10^5$ ,  $0 \leq t \leq 0.2$ , and  $0.4 \leq y \leq 1$ , then*

$$e_A + e_B + e_{C,0} \leq 4.37 \times 10^{-4} + 3.23 \times 10^{-10} \left( F_{N,0.2} \left( 0.7 + \frac{1}{20} \log \frac{x}{4\pi} \right) + |\gamma| F_{N,0.2} \left( 0.3 + \frac{1}{20} \log \frac{x}{4\pi} \right) \right).$$

where

$$(90) \quad F_{N,t}(\sigma) := \sum_{n=1}^N \frac{b_n^t}{n^\sigma}.$$

*Proof.* From Proposition 6.6(iv), (v) one has

$$(91) \quad e_A + e_B \leq (e^{\delta_1} - 1)(F_{N,t}(\text{Re}(s_*)) + |\gamma| F_{N,t}(\text{Re}(s_*) - y - |\kappa|))$$

where

$$\delta_1 := \frac{\frac{t^2}{32} \log^2 \frac{x}{4\pi} + 0.313}{\frac{x}{2} - 3.33}.$$

From Lemma 5.1(vi), the quantity  $\delta_1$  is monotone decreasing in  $x$  in the region (20). Thus we have

$$(92) \quad \delta_1 \leq \frac{\frac{t^2}{32} \log^2 \frac{x_0}{4\pi} + 0.313}{\frac{x_0}{2} - 3.33}$$

whenever  $x \geq x_0 \geq 200$ .



From Proposition 6.6(vi) one has

$$e_{C,0} \leq \left(\frac{x}{4\pi}\right)^{-\frac{1+y}{4}} \exp\left(-\frac{t}{16} \log^2 \frac{x}{4\pi} + \frac{3|\log \frac{x}{4\pi} + i\frac{\pi}{2}| + 3.58}{x - 8.52}\right) \left(1 + \frac{1.24 \times (3^y + 3^{-y})}{N - 0.125} + \frac{6.72}{x - 6.66}\right).$$

From Lemma 5.1(vi), the quantities  $\frac{\log^2 \frac{x}{4\pi} + \frac{\pi^2}{4}}{(x-8.52)^2}$  is monotone decreasing in  $x$  in (20), hence  $\frac{|\log \frac{x}{4\pi} + i\frac{\pi}{2}|}{x-8.52}$  is also monotone decreasing. Also the expression is monotone decreasing in  $y$ . We conclude that (93)

$$e_{C,0} \leq \left(\frac{x_0}{4\pi}\right)^{-\frac{1+y_0}{4}} \exp\left(-\frac{t}{16} \log^2 \frac{x_0}{4\pi} + \frac{3|\log \frac{x_0}{4\pi} + i\frac{\pi}{2}| + 3.58}{x_0 - 8.52}\right) \left(1 + \frac{1.24 \times (3^{y_0} + 3^{-y_0})}{N_0 - 0.125} + \frac{6.72}{x_0 - 6.66}\right)$$

whenever  $x \geq x_0 \geq 200$ ,  $y \geq y_0 > 0$ , and  $N \geq N_0$ . Setting  $x_0 = 5 \times 10^9$ ,  $y_0 = 0.4$ , and  $N_0 = 19947$  and using  $t \geq 0$  we conclude that From (93) one has

$$e_{C,0} \leq \left(\frac{5 \times 10^9}{4\pi}\right)^{-\frac{1+0.4}{4}} \exp\left(-\frac{t}{16} \log^2 \frac{5 \times 10^9}{4\pi} + \frac{3|\log \frac{5 \times 10^9}{4\pi} + i\frac{\pi}{2}| + 3.58}{5 \times 10^9 - 8.52}\right) \left(1 + \frac{1.24 \times (3^{0.4} + 3^{-0.4})}{19947 - 0.125} + \frac{6.92}{5 \times 10^9 - 6.66}\right)$$

From (92) one has

$$\begin{aligned} \delta_1 &\leq \frac{\frac{0.2^2}{32} \log^2 \frac{5 \times 10^9}{4\pi} + 0.313}{\frac{5 \times 10^9}{2} - 3.33} \\ &\leq 3.22 \times 10^{-10} \end{aligned}$$

and hence

$$e^{\delta_1} - 1 \leq 3.23 \times 10^{-10}.$$

From Proposition 6.6(ii) (and the observation that  $1 - 3y + \frac{4y(1+y)}{x^2}$  is negative when  $y \geq 0.4$  and  $x \geq 5 \times 10^9$ ) we have

$$\operatorname{Re}(s_*) \geq 0.7 + \frac{1}{20} \log \frac{x}{4\pi}.$$

Since  $F_{N,t}(\sigma)$  is non-decreasing in  $t$  and non-increasing in  $\sigma$ , the claim follows.  $\square$

It now suffices to establish the following claims:

- (i) (88) holds when  $5 \times 10^9 \leq x \leq 5 \times 10^9 + 1$ ,  $0 \leq t \leq 0.2$ , and  $y \geq 0.4$ .
- (ii) (88) holds when  $x \geq 5 \times 10^9$ ,  $19947 \leq N \leq 40000$ ,  $t = 0.2$ , and  $y \geq 0.4$ .
- (iii) (88) holds when  $40000 \leq N \leq 3 \times 10^5$ ,  $t = 0.2$ , and  $y \geq 0.4$ .
- (iv) (88) holds when  $N \geq 3 \times 10^5$ ,  $t = 0.2$ , and  $y \geq 0.4$ .

We begin with claim (i). We need some derivative estimates on the quantity  $X = X(t, x + iy)$ .

**Lemma 9.3.** *In the region (20), we have*

$$\left|\frac{\partial X}{\partial x}\right| = \left|\frac{\partial X}{\partial y}\right| \leq \sum_{n=1}^N \frac{b'_n}{n^{\operatorname{Re}(s_*)}} \left(\frac{\log n}{2} + \frac{t \log n}{4(x-6)}\right) + |\gamma| N^{|k|} \sum_{n=1}^N \frac{b'_n n^y}{n^{\operatorname{Re}(s_*)}} \left(\frac{t \log n}{4(x-6)} + \left(\log \frac{|1+y+ix|}{4\pi} + \pi + \frac{3}{x}\right) \left(\frac{1}{2} + \frac{t}{4(x-6)}\right)\right)$$

*Proof.* Write

$$s_{**} := \overline{s_*} - y + \kappa = \frac{1-y+ix}{2} + \frac{t}{2} \alpha \left(\frac{1-y+ix}{2}\right)$$

then

$$X = \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N \frac{b_n^t}{n^{s_{**}}}.$$

One can check that  $s_*$ ,  $s_{**}$ ,  $\gamma$  are holomorphic functions of  $x + iy$ , hence by the Cauchy-Riemann equations

$$\left| \frac{\partial X}{\partial x} \right| = \left| \frac{\partial X}{\partial y} \right|.$$

By the product and chain rules, we may calculate

$$\frac{\partial X}{\partial x} = - \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} \frac{\partial s_*}{\partial x} \log n + \gamma \sum_{n=1}^N \frac{b_n^t}{n^{s_{**}}} \left( \frac{\partial}{\partial x} \log \gamma - \frac{\partial s_{**}}{\partial x} \log n \right).$$

From (55), (25) we have

$$\begin{aligned} \frac{\partial s_*}{\partial x} &= -\frac{i}{2} - \frac{it}{4} \alpha' \left( \frac{1-y+ix}{2} \right) \\ &= -\frac{i}{2} + O_{\leq} \left( \frac{t}{4(x-6)} \right). \end{aligned}$$

Similarly we have

$$\frac{\partial s_{**}}{\partial x} = \frac{i}{2} + O_{\leq} \left( \frac{t}{4(x-6)} \right).$$

Writing  $s = \frac{1-y+ix}{2}$ , we have from (54), (26) that

$$\log \lambda = \frac{t}{4} (\alpha(s)^2 - \alpha(1-s)^2) + \log M_0(s) - \log M_0(1-s)$$

and hence by (22)

$$\frac{\partial \lambda}{\partial x} = \frac{it}{4} (\alpha(s) \alpha'(s) + \alpha(1-s) \alpha'(1-s)) + \frac{i}{2} \alpha(s) + \frac{i}{2} \alpha(1-s).$$

From the triangle inequality and (25), we thus have

$$\left| \frac{\partial X}{\partial x} \right| \leq \sum_{n=1}^N \frac{b_n^t}{n^{\operatorname{Re}(s_*)}} \left( \frac{\log n}{2} + \frac{t \log n}{4(x-6)} \right) + |\gamma| \sum_{n=1}^N \frac{b_n^t}{n^{\operatorname{Re}(s_{**})}} \left( \frac{t \log n}{4(x-6)} + \frac{|\alpha(s) + \alpha(1-s) - \log n|}{2} + \frac{t(|\alpha(s)| + |\alpha(1-s)|)}{4(x-6)} \right).$$

We have from (23) that

$$|\alpha(s)|, |\alpha(s) - \frac{1}{2} \log n| \leq \frac{1}{2} \log \frac{|1-y+ix|}{4\pi} + \frac{\pi}{2} + \frac{3}{2x}$$

since  $n \leq N \leq \frac{x}{4\pi} \leq \frac{|1-y+ix|}{4\pi}$ . Similarly

$$|\alpha(1-s)|, |\alpha(1-s) - \frac{1}{2} \log n| \leq \frac{1}{2} \log \frac{|1+y+ix|}{4\pi} + \frac{\pi}{2} + \frac{3}{2x}$$

and thus

$$|\alpha(s) + \alpha(1-s)|, |\alpha(s) + \alpha(1-s) - \log n| \leq \log \frac{|1+y+ix|}{4\pi} + \pi + \frac{3}{x}.$$

Writing  $\operatorname{Re}(s_{**}) = \operatorname{Re}(s_*) - y + \operatorname{Re}(\kappa)$ , we then have the claim.  $\square$

...

Now we prove claim (ii).

...

Now we prove claim (iii).

...

Finally, we prove claim (iv). In this regime we have

$$x \geq 4\pi N_0^2 - \frac{\pi t}{4} \geq 1.13 \times 10^{12}.$$

We use the triangle inequality bound

$$\begin{aligned} |X| &\geq (2 - F_{N,t}(\operatorname{Re}(s_*)) - |\gamma|F_{N,t}(\operatorname{Re}(s_*) - y))_+ \\ &\geq \left(2 - F_{N,0.2}(0.7 + \frac{1}{20} \log \frac{x}{4\pi}) - |\gamma|F_{N,0.2}(0.3 + \frac{1}{20} \log \frac{x}{4\pi})\right)_+ \end{aligned}$$

so to prove (88) here, it suffices by Lemma 9.2 to show that

$$(94) \quad F_{N,0.2}(0.7 + \frac{1}{20} \log \frac{x}{4\pi}) + |\gamma|F_{N,0.2}(0.3 + \frac{1}{20} \log \frac{x}{4\pi}) \leq 2 - 4.38 \times 10^{-4}.$$

From Proposition 6.6(i) we have

$$|\gamma| \leq e^{0.0016} \left(\frac{x}{4\pi}\right)^{-0.2}.$$

We can then bound the left-hand side of (94) by  $Q_1 + Q_2 + Q_3$ , where

$$\begin{aligned} Q_1 &:= F_{N_0,0.2}(0.7 + \frac{1}{20} \log \frac{x}{4\pi}) \\ Q_2 &:= e^{0.0016} \left(\frac{x}{4\pi}\right)^{-0.2} F_{N_0,0.2}(0.3 + \frac{1}{20} \log \frac{x}{4\pi}) \\ Q_3 &:= \sum_{N_0 < n \leq N} \frac{\exp(\frac{1}{20} \log^2 n)}{n^{0.7 + \frac{1}{20} \log \frac{x}{4\pi}}} (1 + e^{0.0016} (\frac{x}{4\pi n^2})^{-0.2}). \end{aligned}$$

We have

$$\begin{aligned} Q_1 &\leq F_{N_0,0.2}(0.7 + \frac{1}{20} \log \frac{1.13 \times 10^{12}}{4\pi}) \\ &\leq 1.898 \end{aligned}$$

and

$$\begin{aligned} Q_2 &\leq e^{0.0016} \left(\frac{1.13 \times 10^{12}}{4\pi}\right)^{-0.2} F_{N_0,0.2}(0.3 + \frac{1}{20} \log \frac{1.13 \times 10^{12}}{4\pi}) \\ &\leq 0.047. \end{aligned}$$

To estimate  $Q_3$ , we first observe that

$$\begin{aligned} \frac{x}{4\pi n^2} &\geq \frac{x}{4\pi N^2} \\ &\geq \left(1 + \frac{\pi t}{4x}\right)^{-1} \\ &\geq 1 - 1.4 \times 10^{-13} \end{aligned}$$

and hence

$$1 + e^{0.0016\left(\frac{x}{4\pi n^2}\right)^{-0.2}} \leq 2.0017.$$

Also from (89) we have

$$\frac{1}{n^{\frac{1}{20} \log \frac{x}{4\pi}}} \leq \frac{1}{n^{\frac{1}{20} \log N^2}} \exp\left(\log\left(1 + \frac{\pi t}{4x}\right) \log N\right).$$

Using Proposition 5.1(vi) we can easily bound

$$\log\left(1 + \frac{\pi t}{4x}\right) \log N \leq 10^{-6}$$

(say), and hence

$$Q_3 \leq 2.0018 \sum_{N_0 < n \leq N} \frac{\exp\left(\frac{1}{20} \log^2 n\right)}{n^{0.7 + \frac{1}{20} \log N^2}}.$$

The function  $n \mapsto \frac{\exp(\frac{1}{20} \log^2 n)}{n^{0.7 + \frac{1}{20} \log N^2}}$  is decreasing for  $1 \leq n < N$ , so by the integral test we have

$$Q_3 \leq 2.0018 \int_{N_0}^N \frac{\exp\left(\frac{1}{20} \log^2 a\right)}{a^{0.7 + \frac{1}{20} \log N^2}} da.$$

Making the change of variables  $a = e^u$ , this becomes

$$Q_3 \leq 2.0018 \int_{\log N_0}^{\log N} \exp\left(0.3u + \frac{1}{20}(u^2 - 2u \log N)\right) du.$$

The quadratic expression  $-0.3u + \frac{1}{20}(u^2 - 2u \log N)$  is convex in  $u$ , and is thus bounded by its maximum at the endpoints; thus

$$Q_3 \leq 2.0018 \left(\log \frac{N}{N_0}\right) \exp\left(\max\left(0.3 \log N_0 - \frac{1}{20} \log N_0 \log \frac{N^2}{N_0}, 0.3 \log N - \frac{1}{20} \log^2 N\right)\right).$$

Writing  $N = e^b N_0$  for some  $b > 0$ , this bound can be rewritten as

$$Q_3 \leq \frac{2.0018}{N_0^{\frac{1}{20} \log N_0 - 0.3}} \exp\left(\log b + \frac{1}{20}(-2b \log N_0 + (6b - b^2)_+)\right).$$

A routine numerical maximisation shows that

$$\log b + \frac{1}{20}(-2b \log N_0 + (6b - b^2)_+) \leq -1.009$$

and hence

$$Q_3 \leq 0.0113.$$

Combining these bounds we obtain (94).

## 10. ASYMPTOTIC RESULTS

In this section we use the effective estimates from Section 6 to obtain asymptotic information about the function  $H_t$ , which improves (and makes more effective) the results of Ki, Kim, and Lee [9]. We use the usual asymptotic notation  $X = O(Y)$  to denote the bound  $X = O_{\leq}(CY)$  for an absolute constant  $C$ .

**Proposition 10.1.** *Let  $0 < t \leq 1/2$ ,  $x \geq 200$ , and  $-10 \leq y \leq 10$ .*

(i) *If  $x \geq \exp(\frac{C}{t})$  for a sufficiently large absolute constant  $C$ , then*

$$H_t(x + iy) = (1 + O(x^{-ct}))M_t\left(\frac{1 + y - ix}{2}\right) + (1 + O(x^{-ct}))M_t\left(\frac{1 - y + ix}{2}\right)$$

*for an absolute constant  $c > 0$ , where  $M_t$  is defined in (26).*

(ii) *If instead we have  $3 \leq y \leq 4$  and  $x \geq C$  for a sufficiently large absolute constant  $C$ , then*

$$H_t(x + iy) = (1 + O_{\leq}(0.7))M_t\left(\frac{1 + y - ix}{2}\right).$$

(iii) *If  $x = x_0 + O(1)$  for some  $x_0 \geq 200$ , then*

$$H_t(x + iy) = O(x_0^{O(1)}|M_t(\frac{1 + y - ix}{2})|) = O(x_0^{O(1)}|M_t(\frac{ix_0}{2})|).$$

*Proof.* We begin with (i). Since  $H_t = H_t^*$  and  $M_t = M_t^*$ , we may assume without loss of generality that  $y \geq 0$ . Using (54), (51) we may write the desired estimate as

$$\frac{H_t(x + iy)}{B_0(x + iy)} = 1 + O(x^{-ct}) + \gamma.$$

We apply the decomposition (53) followed by Proposition 6.6. (Strictly speaking, the estimates there required  $y \leq 1$  rather than  $y \leq 10$ ; however, as remarked at the beginning of Section 6, all the estimates in that section would continue to hold under this weaker hypothesis if one adjusted all the numerical constants appropriately.) This gives

$$(95) \quad \frac{H_t(x + iy)}{B_0(x + iy)} = \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{s_* + \kappa}} + O_{\leq}(e_A + e_B + e_{C,0})$$

where

$$\begin{aligned} \gamma &= O(x^{-y/2}) \\ \kappa &= O(x^{-1}) \\ \operatorname{Re}(s_*) &\geq \frac{1+y}{2} + \frac{t}{2} \log \frac{x}{4\pi} - O(x^{-2}) \\ e_A &= O\left(x^{-y/2} \sum_{n=1}^N b_n^t n^{-\frac{1+y}{2} - \frac{t}{2} \log \frac{x}{4\pi} - O(x^{-1})} \frac{\log^2 x}{x}\right) \\ e_B &= O\left(\sum_{n=1}^N b_n^t n^{-\frac{1+y}{2} - \frac{t}{2} \log \frac{x}{4\pi} + O(x^{-1})} \frac{\log^2 x}{x}\right) \\ e_{C,0} &= O\left(x^{-\frac{1+y}{4}}\right) \end{aligned}$$

Since  $N = O(x^{1/2})$ , we have  $x^{-y/2}n^y = O(1)$  and  $n^{O(x^{-1})} = O(1)$  for all  $1 \leq n \leq N$ . We conclude that

$$\frac{H_t(x+iy)}{B_0(x+iy)} = 1 + \gamma + O\left(\frac{\log^2 x}{x} + \sum_{n=2}^N \frac{b_n^t}{n^{\frac{1+y}{2} + \frac{t}{2} \log \frac{x}{4\pi}}} + x^{-\frac{1+y}{4}}\right)$$

so it will suffice (for  $c$  small enough) to show that

$$\sum_{n=2}^N \frac{b_n^t}{n^{\frac{1+y}{2} + \frac{t}{2} \log \frac{x}{4\pi}}} = O(x^{-ct}).$$

By (27) we can write the left-hand side as

$$\sum_{n=2}^N \frac{1}{n^{\frac{1+y}{2} + \frac{t}{2} \log \frac{x}{4\pi\sqrt{n}}}} = O(x^{-ct}).$$

For  $2 \leq n \leq N$ , we have

$$\frac{1+y}{2} + \frac{t}{2} \log \frac{x}{4\pi\sqrt{n}} \geq ct \log x$$

for some absolute constant  $c > 0$ . By the integral test, the left-hand side is then bounded by

$$\frac{1}{2^{ct \log x}} + \int_2^\infty \frac{1}{u^{ct \log x}} du$$

which, for  $x \geq \exp(C/t)$  and  $C$  large, is bounded by  $O(2^{-ct \log x})$ . The claim then follows after adjusting  $c$  appropriately.

Now we prove (ii). As before we have the expansion (95). We have

$$\begin{aligned} \gamma \sum_{n=1}^N n^y \frac{b_n^t}{n^{\frac{y}{2} + \kappa}} &= O\left(x^{-y/2} \sum_{n=1}^N \frac{b_n^t}{n^{-\frac{1-y}{2} + \frac{t}{2} \log \frac{x}{4\pi}}}\right) \\ &= O\left(x^{-y/2} \sum_{n=1}^N n^{\frac{y-1}{2}}\right) \\ &= O(x^{-\frac{y-1}{4}}); \end{aligned}$$

similar arguments give  $e_A = O(\frac{\log^2 x}{x^{\frac{y-1}{4}}})$ , while

$$\begin{aligned} e_B &= O\left(\frac{\log^2 x}{x} \sum_{n=1}^N b_n^t n^{-\frac{1+y}{2} - \frac{t}{2} \log \frac{x}{4\pi}}\right) \\ &= O\left(\frac{\log^2 x}{x} \sum_{n=1}^N n^{-2}\right) \\ &= O\left(\frac{\log^2 x}{x}\right). \end{aligned}$$

We conclude that

$$\begin{aligned}
\frac{H_t(x+yi)}{B_0(x+yi)} &= \sum_{n=1}^N \frac{b_n^t}{n^{s_*}} + O(x^{-\frac{y-1}{4}}) \\
&= 1 + O_{\leq} \left( \sum_{n=2}^N n^{-\frac{1+y}{2} - \frac{t}{2} \log \frac{x}{4\pi} - O(x^{-1})} \right) + O(x^{\frac{y-1}{4}}) \\
&= 1 + O_{\leq} \left( \sum_{n=2}^N n^{-2} \right) + O(x^{-1/2}) \\
&= 1 + O_{\leq} \left( \frac{\pi^2}{6} - 1 \right) + O(x^{-1/2}) \\
&= 1 + O_{\leq}(0.7)
\end{aligned}$$

as claimed, if  $x \geq C$  for  $C$  large enough.

Finally we prove (iii). Again our starting point is (95). The right-hand side can be bounded crudely by  $O(x^{O(1)}) = O(x_0^{O(1)})$ , hence

$$H_t(x+iy) = O(x_0^{O(1)} |M_t(\frac{1+y+ix}{2})|).$$

However, from (26), (21), (23) it is not hard to see that the log-derivative of  $M_t(s)$  is of size  $O(\log x_0)$  in the region  $s = \frac{ix_0}{2} + O(1)$ . Thus

$$|M_t(\frac{1+y+ix}{2})| = O(x_0^{O(1)} |M_t(\frac{ix_0}{2})|),$$

giving the claim.  $\square$

To understand the behavior of  $M_t(x+iy)$  we make the following simple observations:

**Lemma 10.2.** *Let  $0 < t \leq 1/2$ , let  $x_* > 0$  be sufficiently large, and let  $x+iy = x_* + O(1)$ . Then*

$$M_t(\frac{1+y+ix}{2}) = M_t(\frac{1+ix_*}{2}) \exp \left( (i(x-x_*) + y) \left( \frac{1}{4} \log \frac{x_*}{4\pi} + \frac{\pi i}{8} \right) + O \left( \frac{\log x_*}{x_*} \right) \right).$$

*Also, there is a continuous branch of  $\arg M_t(\frac{1+ix_*}{2})$  for all large real  $x_*$  such that*

$$\arg M_t \left( \frac{1+ix_*}{2} \right) = \frac{t\pi}{16} \log \frac{x_*}{4\pi} + \frac{7\pi}{8} + \frac{x_*}{4} \log \frac{x_*}{4\pi} - \frac{x_*}{4} + O \left( \frac{\log x_*}{x_*} \right).$$

*Proof.* By (26), (22), the log-derivative of  $M_t$  is given by

$$(96) \quad \frac{M'_t}{M_t} = \alpha + \frac{t}{2} \alpha \alpha'.$$

For  $s = \frac{ix_*}{2} + O(1)$ , we have from (23) that

$$(97) \quad \alpha(s) = \frac{1}{2} \log \frac{x_*}{4\pi} + \frac{\pi i}{4} + O \left( \frac{1}{x_*} \right)$$

and from this and (25) we conclude that

$$\frac{M'_t(s)}{M_t(s)} = \frac{1}{2} \log \frac{x_*}{4\pi} + \frac{\pi i}{4} + O \left( \frac{\log x_*}{x_*} \right)$$

whenever  $s = \frac{ix_*}{2} + O(1)$ . The first claim then follows by applying the fundamental theorem of calculus to a branch of  $\log M_t$ .

For the second claim, we calculate

$$\begin{aligned} \arg M_t \left( \frac{1+ix_*}{2} \right) &= \frac{t}{4} \operatorname{Im} \alpha \left( \frac{1+ix_*}{2} \right)^2 + \pi - \frac{x_*}{4} \log \pi + \operatorname{Im} \left( \frac{-1+ix_*}{4} \log \frac{1+ix_*}{4} - \frac{1+ix_*}{4} \right) \\ &= \frac{t}{4} \left( \frac{\pi}{4} \log \frac{x_*}{4\pi} + O\left(\frac{\log x_*}{x_*}\right) \right) + \pi - \frac{x_*}{4} \log \pi + \operatorname{Im} \left( \frac{-1+ix_*}{4} \left( \log \frac{x_*}{4} + \frac{i\pi}{2} - \frac{i}{x_*} + O\left(\frac{1}{x_*^2}\right) \right) \right) - \frac{x_*}{4} \\ &= \frac{t\pi}{16} \log \frac{x_*}{4\pi} + \pi + \frac{x_*}{4} \log \pi + \frac{x_*}{4} \log \frac{x_*}{4} - \frac{\pi}{8} + O\left(\frac{\log x_*}{x_*}\right) \\ &= \frac{t\pi}{16} \log \frac{x_*}{4\pi} + \frac{7\pi}{8} + \frac{x_*}{4} \log \frac{x_*}{4\pi} - \frac{x_*}{4} + O\left(\frac{\log x_*}{x_*}\right) \end{aligned}$$

as desired.  $\square$

Roughly speaking, these estimates tell us that the zeroes of  $H_t$  behaves (on macroscopic scales) like those of  $H_0$  in the region  $x = O(\exp(O(1/t)))$ , and are very evenly spaced outside of this range. More precisely, we have

**Corollary 10.3** (Distribution of zeroes of  $H_t$ ). *Let  $0 < t \leq 1/2$ , let  $C > 0$  be a sufficiently large absolute constant, and let  $c > 0$  be a sufficiently small absolute constant. For all  $n \geq C$ , let  $x_n$  be the unique real number greater than  $4\pi$  such that*

$$(98) \quad \frac{x_n}{4\pi} \log \frac{x_n}{4\pi} - \frac{x_n}{4\pi} + \frac{5}{8} + \frac{t}{16} \log \frac{x_n}{4\pi} = n.$$

(This is well-defined since the left-hand side is an increasing function of  $x_n$  for  $x_n \geq 4\pi$ .)

(i) *If  $x \geq \exp(\frac{C}{t})$  and  $H_t(x+iy) = 0$ , then  $y = 0$ , and*

$$x = x_n + O(x^{-ct}).$$

(ii) *Conversely, for each  $n \geq \exp(\frac{C}{t})$  there is exactly one zero  $H_t$  in the disk  $\{x+iy : |x+iy - x_n| \leq \frac{c}{\log x_n}\}$  (and by part (i), this zero will be real and lie within  $O(x^{-ct})$  of  $x_n$ ).*

(iii) *If  $X \geq \exp(\frac{C}{t})$ , the number  $N_t(X)$  of zeroes with real part between 0 and  $X$  (counting multiplicity) is*

$$N_t(X) = \frac{X}{4\pi} \log \frac{X}{4\pi} - \frac{X}{4\pi} + \frac{t}{16} \log \frac{X}{4\pi} + O(1).$$

(iv) *For any  $X \geq 0$ , one has*

$$N_t(X+1) - N_t(X) \leq O(\log(2+X))$$

and

$$N_t(X) = \frac{X}{4\pi} \log \frac{X}{4\pi} - \frac{X}{4\pi} + O(\log(2+X)).$$

The implied constants in the  $O()$  notation are absolute (in particular, they are independent of  $t$  and  $C$ ).

These results refine Theorems 1.3 and 1.4 of [9], which gave similar results but with constants that depended on  $t$  in a non-uniform (and ineffective) fashion, and error terms that were of shape  $o(1)$  rather than  $O(x^{-ct})$  in the limit  $x \rightarrow \infty$  (holding  $t$  fixed). The results may be compared with



those in [2], who (in our notation) show that assuming RH, the zeroes of  $H_0$  are precisely the solutions  $x_n$  to the equation

$$\frac{1}{2\pi} \arg(-e^{2i\vartheta(x_n/2)} \frac{\zeta'(\frac{1-ix_n}{2})}{\zeta'(\frac{1+ix_n}{2})}) = n$$

for integer  $n$ , where  $-\vartheta(t)$  is the phase of  $\zeta(\frac{1}{2} + it)$  and one chooses a branch of the argument so that the left-hand side is  $-\frac{1}{2}$  when  $x_n = 0$ .

*Proof.* We begin with (ii). Let  $n \geq \exp(\frac{C}{t})$ , and suppose that  $x + iy = x_n + O(1)$ . By Proposition 10.1(i) and Lemma 10.2 we have

$$(99) \quad \begin{aligned} H_t(x + iy) &= \overline{M_t\left(\frac{1+ix_n}{2}\right)} \exp\left((-i(x-x_n) + y)\left(\frac{1}{4} \log \frac{x_n}{4\pi} - \frac{\pi i}{8}\right) + O(x_n^{-ct})\right) \\ &\quad + M_t\left(\frac{1+ix_n}{2}\right) \exp\left((i(x-x_n) - y)\left(\frac{1}{4} \log \frac{x_n}{4\pi} + \frac{\pi i}{8}\right) + O(x_n^{-ct})\right). \end{aligned}$$

From Lemma 10.2 and (98) one has

$$\arg M_t\left(\frac{1+ix_n}{2}\right) = \frac{\pi}{2} + O\left(\frac{\log x_n}{x_n}\right) \bmod \pi$$

and hence

$$(100) \quad \overline{M_t\left(\frac{1+ix_n}{2}\right)} = -\exp\left(O\left(\frac{\log x_n}{x_n}\right)\right) M_t\left(\frac{1+ix_n}{2}\right).$$

If we now make the further assumption  $y = O\left(\frac{1}{\log x_n}\right)$ , we can thus simplify the above approximation as

$$(101) \quad \begin{aligned} H_t(x + iy) &= -M_t\left(\frac{1+ix_n}{2}\right) e^{-\pi(x-x_n)/8} \exp\left((-i(x-x_n) + y)\frac{1}{4} \log \frac{x_n}{4\pi} + O(|y| \log x_n + x_n^{-ct})\right) \\ &\quad + M_t\left(\frac{1+ix_n}{2}\right) e^{-\pi(x-x_n)/8} \exp\left((i(x-x_n) - y)\frac{1}{4} \log \frac{x_n}{4\pi} + O(|y| \log x_n + x_n^{-ct})\right) \\ &= 2iM_t\left(\frac{1+ix_n}{2}\right) e^{-\pi(x-x_n)/8} \left(\sin\left(\frac{x+iy-x_n}{4} \log \frac{x_n}{4\pi}\right) + O(|y| \log x_n + x_n^{-ct})\right). \end{aligned}$$

In particular, if  $x + iy$  traverses the circle  $\{x_n + \frac{c}{\log n} e^{i\theta} : 0 \leq \theta \leq 2\pi\}$  once anti-clockwise and  $c$  is small enough, the quantity  $H_t(x + iy)$  will wind exactly once around the origin, and hence by the argument principle there is precisely one zero of  $H_t$  inside this circle. As the zeroes of  $H_t$  are symmetric around the real axis, this zero must be real. This proves (ii).

Now we prove (i). Suppose that  $H_t(x + iy) = 0$  and  $x \geq \exp(\frac{C}{t})$ . We can assume  $|y| \leq 1$  since it is known (e.g., from [4, Theorem 13]) that there are no zeroes with  $|y| > 1$ .

Let  $n$  be a natural number that minimises  $|x - x_n|$ , then  $x = x_n + O\left(\frac{1}{\log x_n}\right)$  since the derivative of the left-hand side of (98) in  $x_n$  is comparable to  $\log x_n$ . From (99) we have

$$0 = \overline{M_t\left(\frac{1+ix_n}{2}\right)} \exp\left((-i(x-x_n)+y)\left(\frac{1}{4}\log\frac{x_n}{4\pi} - \frac{\pi i}{8}\right) + O(x_n^{-ct})\right) \\ + M_t\left(\frac{1+ix_n}{2}\right) \exp\left((i(x-x_n)-y)\left(\frac{1}{4}\log\frac{x_n}{4\pi} + \frac{\pi i}{8}\right) + O(x_n^{-ct})\right).$$

Thus both summands on the right-hand side have the same magnitude, which on taking logarithms and canceling like terms implies that

$$y\frac{1}{4}\log\frac{x_n}{4\pi} + O(x_n^{-ct}) = -y\frac{1}{4}\log\frac{x_n}{4\pi} + O(x_n^{-ct})$$

and hence  $y = O\left(\frac{x_n^{-ct}}{\log x_n}\right)$ . We can now apply (101) to conclude that

$$\sin\left(\frac{x+iy-x_n}{4}\log\frac{x_n}{4\pi}\right) + O(x_n^{-ct}) = 0$$

which (when combined with the hypothesis that  $|x - x_n|$  is minimal) forces  $x - x_n = O\left(\frac{x_n^{-ct}}{\log x_n}\right)$ . This gives the claim.

Next, we prove (iii). In view of parts (i) and (ii), and adjusting  $C$  if necessary, we may assume that  $X$  takes the form  $X = x_n + \frac{c}{\log x_n}$  for some  $n \geq \exp(\frac{C}{t})$ . By the argument principle,  $N_t(X)$  is equal to  $\frac{-1}{2\pi}$  times the variation in the argument of  $H_t$  on the boundary of the rectangle  $\{x + iy : 0 \leq x \leq X; -3 \leq y \leq 3\}$  traversed clockwise, since there are no zeroes with imaginary part of magnitude greater than one. By compactness, the variation on the left edge  $\{iy : -3 \leq y \leq 3\}$  is  $O(1)$ , and similarly for any fixed portion  $\{x + 3i : 0 \leq y \leq C\}$  of the upper edge. From Proposition 10.1 (and (99)), we see that the variation of  $H_t(x + iy)/M_t(\frac{1+y-ix}{2})$  on the remaining upper edge  $\{x + 3i : C \leq x \leq X\}$  and on the top half  $\{X + iy : 0 \leq y \leq 3\}$  of the right edge are both equal to  $O(1)$ . Since  $H_t = H_t^*$ , the variation on the lower half of the rectangle is equal to that of the upper half. We thus conclude that

$$N_t(X) = -\frac{1}{\pi} \arg M_t\left(\frac{1-iX}{2}\right) + O(1)$$

where we use a continuous branch of the argument of  $M_t\left(\frac{1-iX}{2}\right)$  that is bounded at  $3i$ . The claim now follows from Lemma 10.2.

Finally, we prove (iv). From (4) and the rapid decrease of  $\Phi$  it is easy to verify that the entire function  $H_t$  has order 1, while from the positivity of  $\Phi$  we see that  $H_t$  has no zero at the origin. Thus by the Hadamard factorization theorem we have

$$H_t(z) = \exp(a + bz) \prod_n \left(1 - \frac{z}{z_n}\right) \exp\left(\frac{z}{z_n}\right)$$

for some complex numbers  $a, b$ , where  $z_n$  are the zeroes of  $H_t$  counting multiplicity; using the functional equation  $H_t(z) = H_t(-z)$  we can index the zeros as  $z_n = (z_n)_{n \in \mathbb{Z} \setminus \{0\}}$  with  $z_{-n} = -z_n$ , and conclude that  $b = 0$ , and

$$H_t(z) = \exp(a) \prod_{n>0} \left(1 - \frac{z^2}{z_n^2}\right).$$

Taking logarithmic derivatives, we conclude that

$$(102) \quad \frac{H'_t(z)}{H_t(z)} = \sum_{n>0} \left( \frac{1}{z - z_n} + \frac{1}{z + z_n} \right).$$

Setting  $z = X + 4i$ , we see from Proposition 10.1 and the generalized Cauchy integral formula that the logarithmic derivative of  $H_t(x + iy)/M_t\left(\frac{1+y-ix}{2}\right)$  is equal to  $O(1)$  at  $X + 4i$  for all sufficiently large  $X$ , and hence for all  $X$  by symmetry and compactness. On the other hand, from Stirling's formula (or the logarithmic growth of the digamma function) one easily verifies that the logarithmic derivative of  $M_t\left(\frac{1+y-ix}{2}\right)$  is equal to  $O(\log(2 + X))$  at  $X + 4i$ . Hence  $\frac{H'_t(X+4i)}{H_t(X+4i)} = O(\log(2 + X))$ . Taking imaginary parts, we conclude that

$$\sum_{n>0} \frac{(4 - y_n)}{(X - x_n)^2 + (4 - y_n)^2} + \frac{(4 - y_n)}{(X + x_n)^2 + (4 + y_n)^2} = O(\log(2 + X))$$

where we write  $z_n = x_n + iy_n$ ; equivalently one has

$$\sum_n \frac{(4 - y_n)}{(X - x_n)^2 + (4 - y_n)^2} = O(\log(2 + X))$$

where the sum now ranges over all zeroes, including any at the origin. Since  $|y_n| \leq 1$ , every zero in  $[X, X + 1]$  makes a contribution of at least  $\frac{1}{100}$  (say). As the summands are all positive, the first part of claim (iv) follows. To prove the second part, we may assume by compactness that  $x \geq C$ . Repeating the proof of (iii), and reduce to showing that the variation of  $\arg H_t$  on the short vertical interval  $\{X + iy : 0 \leq y \leq 3\}$  is  $O(\log X)$ . If we let  $\theta$  be a phase such that  $e^{i\theta} H_t(X + 3i)$  is real and positive, we see that this variation is at most  $\pi(m + 1)$ , where  $m$  is the number of zeroes of  $\operatorname{Re} e^{i\theta} H_t(X + yi)$  for  $0 \leq y \leq 3$ , since every increment of  $\pi$  in  $\arg e^{i\theta} H_t$  must be accompanied by at least one such zero. As  $H_t = H_t^*$ , this is also the number of zeroes of  $e^{i\theta} H_t(X + yi) + e^{-i\theta} H_t(2X - (X + yi))$ . On the other hand, from Proposition 10.1(ii), (iii) and Jensen's formula we see that the number of such zeroes is  $O(\log X)$ , and the claim follows.  $\square$

**Remark 10.4.** *The above proposition gives good control on  $H_t(x + iy)$  whenever  $x \geq \exp(C/t)$ . As a consequence (and assuming for sake of argument that the Riemann hypothesis holds), then for any  $\Lambda_0 > 0$ , the bound  $\Lambda \leq \Lambda_0$  should be numerically verifiable in time  $O(\exp(O(1/\Lambda_0)))$ , by applying the arguments of previous sections with  $t$  and  $y$  set equal to small multiples of  $\Lambda_0$ . We leave the details to the interested reader.*

**Remark 10.5.** *Our discussion here will be informal. In view of the results of [8], it is expected that the zeroes  $z_j(t)$  of  $H_t(x + iy)$  should evolve according to the system of ordinary differential equations*

$$\partial_t z_k(t) = 2 \sum_{j \neq k}^{} \frac{1}{z_k(t) - z_j(t)}$$

where the sum is evaluated in a suitable principal value sense, and one avoids those times where the zero  $z_k(t)$  fails to be simple; see [8, Lemma 2.4] for a verification of this in the regime  $t > \Lambda$ . In view of the Riemann-von Mangoldt formula (as well as variants such Corollary 10.3, it is expected that the number of zeroes in any region of the form  $\{x + iy : x + iy = x_* + O(1)\}$  for large  $x_*$  should be of the order of  $\log x_*$ . As a consequence, we expect a typical zero  $z_k(t)$  to move with speed  $O(\log |z_k(t)|)$ , although one may occasionally move much faster than this if two zeroes are exceptionally close together, or less than this if the zeroes are close to being evenly

spaced. As a consequence, if the Riemann hypothesis fails and there is a zero  $x + iy$  of  $H_0$  with  $y$  comparable to 1, it should take time comparable to  $\frac{1}{\log x}$  for this zero to move towards the real axis, leading to the heuristic lower bound  $\Lambda \gg \frac{1}{\log x}$ . Thus, in order to obtain an upper bound  $\Lambda \leq \Lambda_0$ , it will probably be necessary to verify that there are no zeroes  $x + iy$  of  $H_0$  with  $y$  comparable to 1 and  $|x| \leq c \log \frac{1}{\Lambda}$  for some small absolute constant  $c > 0$ . This suggests that the time complexity bound in Remark 10.4 is likely to be best possible (unless one is able to prove the Riemann hypothesis, of course).

In [8, Lemma 2.1] it is also shown that the velocity of a given zero  $z(t)$  is given by the formula

$$\partial_t z(t) = \frac{H_t''(z(t))}{H_t'(z(t))}$$

assuming that the zero is simple. By using the asymptotics in Proposition 10.1 and Corollary 10.3 together with the generalized Cauchy integral formula to then obtain asymptotics for  $H_t'$  and  $H_t''$ , it is possible to show that for the zeroes  $x(t)$  that are real and larger than  $\exp(C/t)$ , and move leftwards with velocity

$$\partial_t x(t) = -\frac{\pi}{4} + O(x^{-ct});$$

we leave the details to the interested reader. **maybe supply some numerically computed graphics here?**

#### REFERENCES

- [1] J. Arias de Reyna, *High-precision computation of Riemann's zeta function by the Riemann-Siegel asymptotic formula, I*, Mathematics of Computation, Volume 80, Number 274, April 2011, Pages 9951009.
- [2] J. Arias de Reyna, J. Van de Lune, *On the exact location of the non-trivial zeroes of Riemann's zeta function*, Acta Arith. **163** (2014), 215–245.
- [3] W. G. C. Boyd, *Gamma Function Asymptotics by an Extension of the Method of Steepest Descents*, Proceedings: Mathematical and Physical Sciences, Vol. 447, No. 1931 (Dec. 8, 1994), pp. 609–630.
- [4] N. C. de Bruijn, *The roots of trigonometric integrals*, Duke J. Math. **17** (1950), 197–226.
- [5] G. Csordas, T. S. Norfolk, R. S. Varga, *A lower bound for the de Bruijn-Newman constant  $\Lambda$* , Numer. Math. **52** (1988), 483–497.
- [6] G. Csordas, A. M. Odlyzko, W. Smith, R. S. Varga, *A new Lehmer pair of zeros and a new lower bound for the de Bruijn-Newman constant  $\Lambda$* , Electronic Transactions on Numerical Analysis. **1** (1993), 104–111.
- [7] G. Csordas, A. Ruttan, R. S. Varga, *The Laguerre inequalities with applications to a problem associated with the Riemann hypothesis*, Numer. Algorithms, **1** (1991), 305–329.
- [8] G. Csordas, W. Smith, R. S. Varga, *Lehmer pairs of zeros, the de Bruijn-Newman constant  $\Lambda$ , and the Riemann hypothesis*, Constr. Approx. **10** (1994), no. 1, 107–129.
- [9] H. Ki, Y. O. Kim, and J. Lee, *On the de Bruijn-Newman constant*, Advances in Mathematics, **22** (2009), 281–306.
- [10] D. H. Lehmer, *On the roots of the Riemann zeta-function*, Acta Math. **95** (1956) 291–298.
- [11] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, Analytic number theory (Proc. Sympos. Pure Math., Vol. XXIV, St. Louis Univ., St. Louis, Mo., 1972), pp. 181–193. Amer. Math. Soc., Providence, R.I., 1973.
- [12] H. L. Montgomery, R. C. Vaughan, *Multiplicative number theory. I. Classical theory*. Cambridge Studies in Advanced Mathematics, 97. Cambridge University Press, Cambridge, 2007.
- [13] C. M. Newman, *Fourier transforms with only real zeroes*, Proc. Amer. Math. Soc. **61** (1976), 246–251.
- [14] T. S. Norfolk, A. Ruttan, R. S. Varga, *A lower bound for the de Bruijn-Newman constant  $\Lambda$  II.*, in A. A. Gonchar and E. B. Saff, editors, Progress in Approximation Theory, 403–418. Springer-Verlag, 1992.
- [15] A. M. Odlyzko, *An improved bound for the de Bruijn-Newman constant*, Numerical Algorithms **25** (2000), 293–303.

- [16] T. K. Petersen, Eulerian numbers. With a foreword by Richard Stanley. Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser/Springer, New York, 2015.
- [17] D. J. Platt, *Isolating some non-trivial zeros of zeta*, Math. Comp. **86** (2017), 2449–2467.
- [18] B. Rodgers, T. Tao, *The De Bruijn-Newman constant is nonnegative*, preprint.
- [19] Y. Saouter, X. Gourdon, P. Demichel, *An improved lower bound for the de Bruijn-Newman constant*, Mathematics of Computation. **80** (2011), 2281–2287.
- [20] J. Stopple, *Notes on Low discriminants and the generalized Newman conjecture*, Funct. Approx. Comment. Math., vol. 51, no. 1 (2014), pp. 23–41.
- [21] J. Stopple, *Lehmer pairs revisited*, Exp. Math. **26** (2017), no. 1, 45–53.
- [22] H. J. J. te Riele, *A new lower bound for the de Bruijn-Newman constant*, Numer. Math., **58** (1991), 661–667.
- [23] E. C. Titchmarsh, *The Theory of the Riemann Zeta-function*, Second ed. (revised by D. R. Heath-Brown), Oxford University Press, Oxford, 1986.

<http://michaelnielsen.org/polymath1/index.php>