

The Well-ordering Theorem

Kevin Ma

January 22, 2020

1 Introduction

Given a set S , an order relation \prec is a well-ordering if all subsets of S contain a least element under \prec . The Axiom of Choice was introduced by Ernst Zermelo in 1904^[1] in order to prove his formulation of the well-ordering theorem, also known as Zermelo's theorem^[2]:

Every set can be well-ordered.

Given Zermelo's formulation, one has to assume the axiom of choice since sets are not implicitly ordered. However, if a strict total ordering on a set, one can construct a well-ordering. This implies that the following statement can be deduced in first order logic independent of the axiom of choice:

Every set that can be totally ordered can be well-ordered.

This has implications for the mathematical representation of physical systems, since one assumes upon construction of a model that there exists a total ordering on the system's components. Since defining a well-ordering given an existing order relation is valid independent of axioms, this provides further evidence that mathematics is an accurate representation of the natural world by removing the need to assume the axiom of choice when defining well-orderings on sets.

2 Nested Sets

We begin with the well-defined order properties of nested sets.

Definition 2.1. *Let S be a set. Then S is a collection of nested sets if*

$$\forall x, y \in S, x \neq y \iff (x \subset y) \vee (y \subset x)$$

Lemma 2.1. *Let S be a collection of nested sets, $A, B \in S$. Then*

$$A \subset B \iff (A \cap B = A) \wedge (A \cup B = B)$$

Lemma 2.2. *Let S be a collection of nested sets such that*

$$\exists z \in S, (z \neq \emptyset) \wedge (\forall x \in S, z \in x)$$

Then S contains a least element.

Proof. Since $\forall x \in S, z \in S$. the set

$$L = \bigcap_{x \in S} x \neq \emptyset$$

since $z \in L$. Furthermore, by the nested property of intersection,

$$\forall x \in S, L \subseteq x$$

which implies that $L \in S$ and L is the least element in S . □

3 The Well-ordering Principle

The order properties of nested sets can be generalized to any totally ordered set.

Definition 3.1 (Total order relation). *Let S be a set with identity relation $=$. Then a binary relation $<$ is a total order relation on S if for all $x, y \in S$:*

- $(x < y) \iff \neg(y < x) \wedge \neg(x = y)$
- $(y < x) \iff \neg(x < y) \wedge \neg(x = y)$
- $(x = y) \iff \neg(x < y) \wedge \neg(y < x)$

Define $(S, <)$ to be the set S ordered under $<$.

Definition 3.2 (Well-ordered). $(S, <)$ is well-ordered if every subset $X \subseteq S$ contains a least element under $<$.

Definition 3.3. Let $(A, <)$ and (B, \ll) be two totally ordered sets. Then a function $f : A \rightarrow B$ is order preserving if

$$\forall x, y \in A, x < y \iff f(x) \ll f(y)$$

One can construct an order-preserving bijection from any totally ordered set to a collection of nested sets. Define the partition of S induced by x to be the set $\{L_x, x, G_x\}$ such that

$$1. L_x = \{s \in S | s < x\}$$

$$2. G_x = \{s \in S | s > x\}$$

Clearly, $S = L_x \sqcup x \sqcup G_x$.

Define a closed map F on S to be,

$$F(x) = L_x \cup x$$

Define $\mathcal{L}[S] = \{F(s) | s \in S\}$.

As a totally ordered set, for each $x \in S$, $F(x)$ clearly contains a maximal element. However, when considering the set $F(x)$ as a member of $\mathcal{L}[S]$, for all $y \in S$,

$$x < y \implies (L_x \subset L_y) \wedge (x \in L_y) \implies F(x) \subseteq L_y \subset F(y)$$

Since F is an injectively defined function on S , it is trivially bijective. Hence, F is an order-preserving map such that

$$x < y \iff F(x) \subset F(y)$$

Theorem 3.1 (The well-ordering principle). Suppose $(S, <)$ is totally ordered and contains a least element. Then every subset $X \subseteq S$ has a least element under $<$.

Proof. Let z be the least element of S . For each $x \in S$, define

$$F(x) = L_x \sqcup x$$

By definition of least element, $F(z) = \emptyset \sqcup z = z$. Since

$$\forall x, y \in S, x < y \implies F(x) \subset F(y)$$

F is an order-preserving bijection. In addition,

$$\forall x \in S, z = F(z) \subseteq F(x)$$

thus, $\mathcal{L}[S] = \{F(x) | x \in S\}$ is totally ordered under \subset with least element z .

Let $X \subseteq S$. Define

$$\mathcal{L}[X] = \{F(x) | x \in X\}$$

Then $\forall F(x) \in \mathcal{L}[X], z \subseteq F(x)$ which implies that

$$F_0 := \bigcap_{F(x) \in \mathcal{L}[X]} F(x) \neq \emptyset$$

Thus, $F_0 \in \mathcal{L}[X]$ by the properties of intersection and

$$(\forall x \in X)(F_0 \subseteq F(x))$$

Since $\mathcal{L}[X]$ is bijective and $F_0 \in \mathcal{L}[X]$,

$$\exists! x_0 \in X, F(x_0) = F_0$$

Thus,

$$(\forall x \in X)(F_0 \subseteq F(x)) \implies (\forall x \in X)(x_0 \leq x)$$

and $x_0 \in X$ is the least element of X . □

4 The Well-ordering Theorem

Lemma 4.1 (Existence of a well-ordering). *Let $(S, <)$ be a totally ordered set. Then there exists a strict total ordering \prec such that (S, \prec) has a least element.*

Proof. Let $z \in S$ such that

$$\begin{aligned} L_z &:= \{x \in S | x < z\} \neq \emptyset \\ G_z &:= \{x \in S | z < x\} \neq \emptyset \end{aligned} \tag{1}$$

such that $S = L_z \sqcup z \sqcup G_z$. Define the order relation \prec such that

$$z \prec \{x_1 | (\forall x_1 \in L_z)\} \prec \{x_2 | (\forall x_2 \in G_z)\}$$

and $\forall x_1, x_2 \in S$,

$$x_1 \prec x_2 \iff \neg(x_1 = x_2) \wedge \begin{cases} ((x_1 \in L_z) \wedge (x_2 \in G_z)) \\ \vee ((x_1 < x_2) \wedge (x_1 \in L_z) \wedge (x_2 \in L_z)) \\ \vee ((x_1 < x_2) \wedge (x_1 \in G_z) \wedge (x_2 \in G_z)) \\ \vee (x_1 = z) \end{cases}$$

Then z is the least element of (S, \prec) . □

Theorem 4.2 (Well-ordering theorem). *Suppose $(S, <)$ is totally ordered. Then S can be well-ordered.*

Proof. If S has 2 elements, then it is trivially well-ordered. Thus, suppose $(S, <)$ is a total ordering on more than 2 elements. By Lemma 4.1, there exists a strict total order relation \prec such that (S, \prec) contains a least element. By the well-ordering principle, for all subsets $X \subseteq S$, (X, \prec) has a least element. Thus, every set $X \subseteq S$ has a least element under \prec and S is well-ordered. □

References

- [1] Bell, John L. (2008), "The Axiom of Choice", The Stanford Encyclopedia of Philosophy, Metaphysics Research Lab, Stanford University, <https://plato.stanford.edu/archives/spr2019/entries/axiom-choice/>
- [2] Wikipedia contributors (2019). "Well-ordering theorem", Wikipedia, *The Free Encyclopedia* https://en.wikipedia.org/w/index.php?title=Well-ordering_theorem&oldid=931579814