

# Equivalency of Poincaré inequality and functional on the weighted Sobolev spaces

Raj Narayan Dhara

Joint work with Prof. Agnieszka Kałamańska

Faculty of Mathematics, Informatics and Mechanics, University  
of Warsaw  
Warszawa

May 14, 2014

## Definition

Let  $K$  be a compact set in  $\Omega \in \mathbb{R}^n$  and let  $\Phi(x, \xi) \geq 0$  be a continuous function in  $\Omega \times \mathbb{R}^n$  and positive homogeneous of the first degree with respect to  $\xi$ . Denote

$\mathcal{W}(K, \Omega) := \{u \in \mathcal{D}(\Omega) : u \geq 1 \text{ on } K\}$ . The number

$$(p, \Phi)\text{-cap}(K, \Omega) := \inf \left\{ \int_{\Omega} [\Phi(x, \nabla u)]^p dx : u \in \mathcal{W}(K, \Omega) \right\}, \quad (1)$$

where  $p \geq 1$ , is called the  $(p, \Phi)$ -capacity of  $K$  relative to  $\Omega$  and is denoted by  $(p, \Phi)\text{-cap}(K, \Omega)$ .

## Theorem

1. *If there exists a constant  $\beta$  such that for any compact set  $F \subset \Omega$*

$$\mu(F)^{\alpha p} \leq \beta \cdot (p, \Phi)\text{-cap}(F, \Omega), \quad (2)$$

*where  $p \geq 1$ ,  $\alpha > 0$ ,  $\alpha p \leq 1$ , then for all  $u \in \mathcal{D}(\Omega)$*

$$\|u\|_{L^q(\Omega, \mu)}^p \leq C \int_{\Omega} [\Phi(x, \nabla u)]^p \, dx, \quad (3)$$

*where  $q = \alpha^{-1}$  and  $C \leq p^p(p-1)^{1-p}\beta$ .*

2. *If (3) holds for any  $u \in \mathcal{D}(\Omega)$  and if the constant  $C$  does not depend on  $u$ , then (2) is valid for all compact set  $F \subset \Omega$  with  $\alpha = q^{-1}$  and  $\beta \leq C$ .*

## Corollary

Assume  $\Phi(x, \lambda) = \rho(x)^{1/p}|\lambda|$ ,  $p = q$  and for  $p$ -admissible weight we have  $d\mu(x) = \rho(x)dx$  (see Chapter 1.1 in [?]), we have

- i) if there exists a constant  $\beta$  such that for any compact set  $F \subset \Omega$

$$\mu(F) := \int_F \rho(x) dx \leq \beta \cdot (p, \Phi)\text{-cap}(F, \Omega), \quad (4)$$

then for all  $u \in \mathcal{D}(\Omega)$

$$\|u\|_{L^p_\rho(\Omega)} \leq C \|\nabla u\|_{L^p_\rho(\Omega)}, \quad (5)$$

where  $C \leq p(p-1)^{(1-p)/p}$ .

- ii) If (4) holds for  $u \in \mathcal{D}(\Omega)$  and if the constant  $C$  does not depend on  $u$ , then (5) is valid for all compact set  $F \subset \Omega$  and  $\beta \leq C$ .

## Definition

Let  $\Omega$  be any open subset of  $\mathbb{R}^n$  and  $\Psi$  be the locally integrable function defined on  $\Omega$  such that for every nonnegative compactly supported  $w \in W^{1,p}(\Omega)$ ,

$$\int_{\Omega} \Psi w \, dx > -\infty. \quad (6)$$

Let  $u \in W_{\text{loc}}^{1,p}(\Omega)$  and  $u \neq 0$  a.e. We say that

$$-\Delta_p u \geq \Psi, \quad (7)$$

if for every non-negative compactly supported  $w \in W^{1,p}(\Omega)$  we have

$$\langle -\Delta_p u, w \rangle := \int_{\Omega} |\nabla u|^{p-2} \langle \nabla u, \nabla w \rangle \, dx \geq \int_{\Omega} \Psi w \, dx. \quad (8)$$

## Definition

**$(\Psi, p)$ -condition:** Suppose  $u$  and  $\Psi$  are as in Definition 1.4 and moreover there exists

$$(\Psi, p)\sigma_0 := \inf \{ \sigma \in \mathbb{R} : \Psi \cdot u + \sigma |\nabla u|^p \geq 0 \text{ a.e. in } \Omega \cap \{u > 0\} \} \in \quad (9)$$

where we set  $\inf \emptyset = +\infty$ .

## Theorem (I. SKRZYPCZAK)

Suppose  $1 < p < \infty$  and there exists a non-negative solution  $v \in W_{\text{loc}}^{1,p}(\Omega)$ , to PDI in the sense of Definition 1.4

$$\begin{cases} \Delta_p v(x) \geq \Psi \\ v(x) > 0 \end{cases} \quad \text{in } \Omega, \quad (10)$$

where  $\Psi$  is locally integrable and satisfies  $(\Psi, p)$  with  $\sigma_0 \in \mathbb{R}$  given by (9). Assume further that  $\beta$  and  $\sigma$  are arbitrary numbers such that  $\beta > 0$  and  $\beta > \sigma \geq \sigma_0$ . Then for every Lipschitz function  $\xi$  with compact support in  $\Omega$  we have

$$\int_{\Omega} |\xi|^p \mu_1(dx) \leq \int_{\Omega} |\nabla \xi|^p \mu_2 dx. \quad (11)$$

where

## Theorem (I. SKRZYPCZAK)

$$\begin{aligned}\mu_1(dx) &:= \left(\frac{\beta - \sigma}{p - 1}\right)^{p-1} [\Psi \cdot v + \sigma |\nabla v|^p] \cdot v^{-\beta-1} \chi_{\{v>0\}} dx \\ \mu_2(dx) &:= v^{p-\beta-1} \chi_{\{|\nabla v| \neq 0\}} dx =\end{aligned}$$

Applying this theorem for  $0 \leq \Psi = \varphi(x)|v(x)|^{p-2}v(x)$ ,  $\sigma = 0$ , where  $\varphi$  is a non-negative function. Assume

$$\mu_1(dx) = \rho_1(x)dx \quad \text{and} \quad \mu_2(dx) = \rho_2(x)dx. \quad (12)$$



Then we have

$$\begin{aligned}
 \rho_1(x) &= \left( \frac{\beta}{p-1} \right)^{p-1} [\Psi \cdot v] \cdot v^{-\beta-1} \chi_{\{v>0\}}, \\
 &\geq \left( \frac{\beta}{p-1} \right)^{p-1} [\varphi |v|^{p-2} v \cdot v] \cdot v^{-\beta-1} \chi_{\{v>0\}}, \\
 &= \left( \frac{\beta}{p-1} \right)^{p-1} \varphi |v|^{p-\beta-1} \chi_{\{|\nabla v| \neq 0\}} = \left( \frac{\beta}{p-1} \right)^{p-1} \phi \rho_2(x)
 \end{aligned}$$

Then we have for  $\phi \geq C$ , some constant and from (11),

$$\int_{\Omega} |\xi|^p \rho_1(x) (dx) \leq C \left( \frac{\beta}{p-1} \right)^{p-1} \int_{\Omega} |\nabla \xi|^p \rho_1(x) dx. \quad (13)$$

and from (ii) of Corollary 1.3, then we have for (1.6), there exists a constant  $\beta'$  such that for any compact set  $F \subset \Omega$

$$\mu(F) := \int_F \rho_1(x) dx \leq \beta' \cdot (p, \Phi)\text{-cap}(\Omega, F). \quad (14)$$

Thank  
you