

1 **Sticky Dead Microbes: rapid abiotic retention of microbial necromass in soil**

2 Buckeridge, Kate M.^{a,1,*}, La Rosa, Alfio Fabio^{a,2}, Mason, Kelly E.^b, Whitaker, Jeanette^b,
3 McNamara, Niall P.^b, Grant, Helen K.^b, Ostle, Nick^a

4 a. Lancaster Environment Centre, University of Lancaster, Lancaster, UK

5 b. UK Centre for Ecology & Hydrology, Lancaster, UK

6 * Corresponding author: kmbuckeridge@gmail.com, +44 (0) 7482 929 296

7 1. Present address: Global Academy of Agriculture and Food Security, The Royal (Dick)

8 School of Veterinary Studies, University of Edinburgh, Edinburgh, UK

9 2. Present address: Department of Mathematics, University of Bonn, Bonn, DE

10 Article type: Short communication

11

Abstract

Microbial necromass dominates soil organic matter. Recent research on necromass and soil carbon storage has focused on necromass production and stabilization mechanisms but not on the mechanisms of necromass retention. We present evidence from soil incubations with stable-isotope labelled necromass that abiotic adsorption may be more important than biotic immobilization for short-term necromass retention. We demonstrate that necromass adsorbs not only to mineral surfaces, but may also interact with other necromass. Furthermore, necromass cell chemistry alters necromass-necromass interaction, with more bacterial tracer retained when there is yeast necromass present. These findings suggest that the adsorption and abiotic interaction of microbial necromass and its functional properties, beyond chemical stability, deserve further investigation in the context of soil carbon sequestration.

Keywords

Soil organic matter; functional properties; stable isotope; grassland pasture; carbon sequestration; nitrogen

28 **Main text**

29 Soil organic matter (SOM) is a key indicator for healthy soil and sustainable agriculture
30 (Paustian et al., 2016). Research into SOM stability has traditionally focused on the quality
31 and quantity of plant inputs to soil (Lehmann and Kleber, 2015), however, recent research
32 reveals that SOM is dominated by dead microbial products and residues (hereafter,
33 ‘necromass’; (Kallenbach et al., 2015; Liang et al., 2019, 2011)). The persistence of
34 necromass in soil may be promoted via necromass uptake and immobilization into microbial
35 biomass, but retention ultimately relies on microaggregate formation via adsorption to soil
36 mineral surfaces. In actuality, both biotic (microbial immobilization) and abiotic retention
37 (adsorption and molecular interaction) may co-occur but the relative importance of these
38 short-term processes has not been assessed (Fig.1).

39 Necromass has been visualized on mineral surfaces (Kleber et al., 2011), supporting the
40 paradigm that accumulation of stable SOM is dominated by organic-mineral adsorption, and
41 is limited by mineral surface area (McNally et al., 2017). However, necromass is not
42 detected as a smooth coverage on mineral surfaces, but clumpy (Dignac et al., 2017; Vogel
43 et al., 2014), suggesting that SOM stabilization may also involve organic-organic
44 interactions, or necromass adhering to other necromass and organic matter for example by
45 ionic interactions, hydrogen bridges, van der Waals forces, and (partial) entrapment
46 (Schweizer et al., 2018; Vogel et al., 2014) (Fig.1). Understanding the relative importance of
47 these two abiotic adsorption processes will be critical for predicting upper limits of SOM
48 persistence.

49 The chemistry of microbial necromass has been assumed unimportant as a regulator of SOM
50 storage, because its chemical composition is more similar than diverse plant inputs (Liang et

al., 2017). Previous research on necromass chemistry and persistence has focused on its stability and emphasized chitin retention (Fernandez et al., 2016; Schreiner et al., 2014). However, cell chemistry can alter rates of cell-cell adhesion (Dufrêne, 2015) and organic-mineral adsorption rates (Meissner et al., 2015), particularly for N-rich necromass (Kopittke et al., 2017). Gram-positive bacterial envelopes have a thick cross-linked peptidoglycan layer outside the lipid membrane, whereas a Gram-negative cell envelope has an inner and outer lipid membrane enclosing a thinner peptidoglycan layer in the periplasm. Fungal cell walls are highly heterogeneous; yeast cell walls, for example, are composed of layered mannan, β -glucans and chitin outside a lipid membrane. Therefore, cell-membrane functional groups with a high N-content, such as peptidoglycan-rich Gram-positive bacteria, may be favored for organic-organic interaction, relative to Gram-negative bacteria or fungi. Given the potential for broad shifts in microbial community composition as a result of global change, land use change or even seasonality (Buckeridge et al., 2013; Ramirez et al., 2012), necromass cell chemistry may influence SOM stabilization at ecosystem scales.

We investigated the importance of biotic and abiotic necromass retention in grassland soil and the influence of necromass chemistry on this retention in short-term laboratory incubations. We hypothesized: H1. both biotic and abiotic necromass retention occur, and that biotic retention is more important; H2. more necromass is retained in soil with higher background concentrations of necromass (implying organic-organic adhesion); and H3. abiotic retention would be higher for Gram-positive bacterial membranes (implying cell chemistry is important). We tested these hypotheses with short-term (3-d) laboratory incubations of clay-loam agricultural pasture soil; the duration was chosen to limit biotic processing of rapid abiotic interactions. Detailed methods are available in supplementary

methods. Soil was assigned to live and sterile treatments with enhanced background
necromass (~5.5 mg C and ~1.5 mg N g⁻¹ soil fwt) from three soil microbial functional groups
(Gram-negative (*Escherichia coli*) and Gram-positive bacteria (*Micrococcus luteus*) and yeast
(*Saccharomyces cerevisiae*)), and incubated with a no-addition control. We also added a
small amount of isotopic tracer necromass consisting of ¹³C¹⁵N labeled (10 atm%) *E.coli*
necromass (0.050 mg C and 0.016 mg N g⁻¹ soil fwt) to all treatments; the dual label was
used to aid assessment of biotic vs. abiotic retention. Headspace CO₂ and N₂O samples were
collected throughout the 3-day incubation. At the end of the incubation, we used the
chloroform-direct-extraction method to estimate microbial biomass immobilization of the
tracer (measured in all, detected only in live soils), and the balance between immobilization
and loss as CO₂ was used to calculate carbon use efficiency of the live microbes. The C, N,
δ¹³C and δ¹⁵N of the post-extraction soils (i.e. removing any free or loosely-bound C or N)
was used to assess total retention (assumed biotic + abiotic in the live soils and abiotic only
in the sterile soils), and mixed models were used to assess the statistical significance
(α=0.05) of our results.

Retention of C from the tracer necromass was lower in live vs. sterile soils (P=0.006),
rejecting H1 and indicating that short-term retention of necromass C was dominated (>70%)
by abiotic processes (Fig.2a). Our results, however, cannot confirm that biotic processes are
unimportant for persistent SOM-C accrual. Our controlled environment incubation may
have overestimated the importance of abiotic processes: in a more dynamic, natural system
with active plant-microbial interactions, live microbial immobilization of necromass-C and
plant and microbial uptake/immobilization of necromass-N is potentially critical to
retention. Furthermore, our lab-grown, single-culture additions only approximate the

retention of chemically and taxonomically-complex native necromass. Nonetheless, our results illustrate an important short-term effect of abiotic C retention in soil.

The lower retention of ^{13}C in live vs. sterile soils was not predicted and we suspected this was either a side-effect of sterilization, or CO_2 loss from microbial activity. We discounted the sterilization effect, because we did not see a parallel lower retention of necromass- ^{15}N in live compared to dead soils (Fig.2b). This, combined with no change in C:N ratio of tracer retention in sterile soils and a drop in C:N ratio of retention in live soils ($P < 0.001$, data not shown) equates with live microbial processing of C, associated with immobilization/loss of 4-10% of the total C added, as extractable biomass or CO_2 (Fig.3a-b). Losses of N_2O from the labeled necromass were comparable to CO_2 ($< 2\%$, Fig.3c) but we presume that N uptake to microbial biomass (not measured) would be lower than C, reflecting the lower N demand for growth in these N-rich pasture soils. C-starvation during incubation may have resulted in use of necromass-C for maintenance, reflected in a low substrate use efficiency (CUE) (Fig.3d).

Higher C and N was retained in live and sterile soils with enhanced background necromass compared to controls ($\sim 10\text{-}40\%$ higher, Fig. 2a-b), supporting the hypothesis (H2) that necromass may be adhering to necromass. Organic-organic adhesion has been suggested previously based on isotopic and visual evidence in laboratory and long-term field studies (Schweizer et al., 2018). We did not observe higher C and N retention in soils with enhanced peptidoglycan (*M. luteus*) necromass, rejecting H3, that abiotic retention would be higher for Gram-positive bacterial membranes. However, we provide novel evidence that the retention of C and N from the necromass tracer was higher in live and dead soils augmented with *S. cerevisiae* necromass (C: $P < 0.001$; N: $P = 0.03$). This higher microbial C and N retention in *S. cerevisiae*-amended soils does not appear to be a biotic process, because the CUE of

the tracer necromass in live soils did not differ between background necromass types (Fig. 3d) (despite all enhanced background necromass treatments being lower than the no-addition control, which was presumably a response to higher substrate concentration relative to the control (Geyer et al., 2019)). Instead, this enhanced retention of the tracer necromass in the presence of *S. cerevisiae* necromass may be indicative of faster or stronger interaction between the complex morphogenesis of the Gram-negative outer cell membrane and yeast cell walls, such as occur in live microbial communities (Dufrêne, 2015). Further, compound-specific research is required to understand if the properties of yeast necromass extend to other fungal necromass generally and to specific bacterial membrane and fungal cell wall compounds. Regardless, this result indicates that cell chemistry contributes to an adhesion mechanism that promotes necromass stability in soil.

We conclude that abiotic processes are important for short-term retention of necromass-C and N in soils and require greater emphasis in studies investigating SOM stability. Our results support research that indicates organic-organic interaction promotes retention of C and N and contributes novel evidence that this mechanism is regulated by cell chemistry. If this short-term abiotic retention occurs *in situ* and persists, then microbial community structure and possibly the fungal:bacterial ratio, may influence C and N stabilization through variations in community cell chemistry. Field additions of isotopically-labelled necromass from different taxa, in different soils, and followed by secondary ion mass spectrometry (nanoSIMS), would be a valuable start. These findings suggest that abiotic adsorption and interaction of microbial necromass and its functional properties beyond chemical stability (i.e. cell molecular property, aggregations, and morphology), deserve further investigation in the context of soil carbon sequestration.

143

144 **Acknowledgements**

145 At UKCEH, we thank Glenn Rhodes for his help with the necromass-making process and
146 Andrew Stott for his assistance with isotope analysis. This research was funded by the
147 Natural Environment Research Council (NERC U-Grass NE/M017125/1 and NE/M016757)
148 awarded to NO, JW and NPM for KMB and the Soils Training and Research Studentships
149 (STARS) awarded to KMB for AFL.

150

151 **References**

- 152 Buckeridge, K.M., Banerjee, S., Siciliano, S.D., Grogan, P., 2013. The seasonal pattern of soil
153 microbial community structure in mesic low arctic tundra. *Soil Biol. Biochem.* 65, 338–
154 347. doi:10.1016/j.soilbio.2013.06.012
- 155 Dignac, M., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet,
156 G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., 2017. Increasing soil carbon
157 storage : mechanisms, effects of agricultural practices and proxies. A review. *Agron.*
158 *Sustain. Dev.* 37, 14. doi:10.1007/s13593-017-0421-2
- 159 Dufrêne, Y.F., 2015. Sticky microbes: Forces in microbial cell adhesion. *Trends Microbiol.* 23,
160 376–382. doi:10.1016/j.tim.2015.01.011
- 161 Fernandez, C.W., Langley, J.A., Chapman, S., McCormack, M.L., Koide, R.T., 2016. The
162 decomposition of ectomycorrhizal fungal necromass. *Soil Biol. Biochem.* 93, 38–49.
163 doi:10.1016/j.soilbio.2015.10.017

164 Fierer, N., Allen, A.S., Schimel, J.P., Holden, P.A., 2003. Controls on microbial CO₂
 165 production: a comparison of surface and subsurface soil horizons. *Glob. Chang. Biol.* 9,
 166 1322–1332.

167 Geyer, K.M., Dijkstra, P., Sinsabaugh, R., Frey, S.D., 2019. Clarifying the interpretation of
 168 carbon use efficiency in soil through methods comparison. *Soil Biol. Biochem.* 128, 79–
 169 88. doi:10.1016/j.soilbio.2018.09.036

170 Kallenbach, C.M., Grandy, A.S., Frey, S.D., Diefendorf, A.F., 2015. Microbial physiology and
 171 necromass regulate agricultural soil carbon accumulation. *Soil Biol. Biochem.* 91, 279–
 172 290. doi:10.1016/j.soilbio.2015.09.005

173 Kleber, M., Nico, P.S., Plante, A., Filley, T., Kramer, M., Swanston, C., Sollins, P., 2011. Old
 174 and stable soil organic matter is not necessarily chemically recalcitrant: Implications for
 175 modeling concepts and temperature sensitivity. *Glob. Chang. Biol.* 17, 1097–1107.
 176 doi:10.1111/j.1365-2486.2010.02278.x

177 Kopittke, P.M., Hernandez-Soriano, M.C., Dalal, R.C., Finn, D., Menzies, N.W., Hoeschen, C.,
 178 Mueller, C.W., 2017. Nitrogen-rich microbial products provide new organo-mineral
 179 associations for the stabilization of soil organic matter. *Glob. Chang. Biol.* 12, 3218–
 180 3221. doi:10.1111/gcb.14009

181 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature* 528,
 182 60–68. doi:10.1038/nature16069

183 Liang, C., Amelung, W., Lehmann, J., Kästner, M., 2019. Quantitative assessment of
 184 microbial necromass contribution to soil organic matter. *Glob. Chang. Biol.* 25, 3578–
 185 3590. doi:10.1111/gcb.14781

186 Liang, C., Cheng, G., Wixon, D.L., Balser, T.C., 2011. An Absorbing Markov Chain approach to
 187 understanding the microbial role in soil carbon stabilization. *Biogeochemistry* 106,
 188 303–309. doi:10.1007/s10533-010-9525-3

189 Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial control
 190 over soil carbon storage. *Nat. Microbiol.* 2, 1–6. doi:10.1038/nmicrobiol.2017.105

191 McNally, S.R., Beare, M.H., Curtin, D., Meenken, E.D., Kelliher, F.M., Calvelo Pereira, R.,
 192 Shen, Q., Baldock, J., 2017. Soil carbon sequestration potential of permanent pasture
 193 and continuous cropping soils in New Zealand. *Glob. Chang. Biol.* 23, 4544–4555.
 194 doi:10.1111/gcb.13720

195 Meissner, J., Prause, A., Bharti, B., Findenegg, G.H., 2015. Characterization of protein
 196 adsorption onto silica nanoparticles: influence of pH and ionic strength. *Colloid Polym.*
 197 *Sci.* 293, 3381–3391. doi:10.1007/s00396-015-3754-x

198 Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P., Smith, P., 2016. Perspective
 199 Climate-smart soils. *Nature* 532, 49–57. doi:10.1038/nature17174

200 Ramirez, K.S., Craine, J.M., Fierer, N., 2012. Consistent effects of nitrogen amendments on
 201 soil microbial communities and processes across biomes. *Glob. Chang. Biol.* 18, 1918–
 202 1927. doi:10.1111/j.1365-2486.2012.02639.x

203 Schreiner, K.M., Blair, N.E., Levinson, W., Egerton-Warburton, L.M., 2014. Contribution of
 204 fungal macromolecules to soil carbon sequestration, in: Hartemink, A., McSweeney, K.
 205 (Eds.), *Soil Carbon*. Springer International Publishing, pp. 155–161. doi:DOI
 206 10.1007/978-3-319-04084-4_16

Schweizer, S.A., Hoeschen, C., Schlüter, S., Kögel-Knabner, I., Mueller, C.W., 2018. Rapid soil formation after glacial retreat shaped by spatial patterns of organic matter accrual in microaggregates. *Glob. Chang. Biol.* 24, 1637–1650. doi:10.1111/gcb.14014

Vogel, C., Mueller, C.W., Hoeschen, C., Buegger, F., Heister, K., Schulz, S., Schlöter, M., Kögel-Knabner, I., 2014. Submicron structures provide preferential spots for carbon and nitrogen sequestration in soils. *Nat. Commun.* 5, 1–7. doi:10.1038/ncomms3947

Figure captions

Figure 1. Schematic diagram of microbial necromass fate in mineral soils. In this diagram, necromass is presented as a substrate for microbes, similar to plant inputs (litter or exudates). Microbes can acquire unassociated or mineral-associated necromass for substrate. Immobilization includes recycling of necromass by microbes into new biomass, and eventually necromass, with potential for some loss as CO₂. Stabilization is assumed to be through adsorption to mineral surfaces ('organic-mineral'), especially on fine silts and clays. In this study we hypothesize that this process is not limited to mineral surface availability ('organic-mineral'), but that necromass also adheres to necromass ('organic-organic'), promoting retention, and that this organic-organic process may be influenced by necromass cell chemistry.

Figure 2. The retention of necromass (¹³C¹⁵N-*E.coli*) carbon and nitrogen in live and sterile soil with different background necromass. Data presented as the mean (± standard error, n=5) % retention of applied ¹³C¹⁵N necromass as **(a)** carbon (C) or **(b)** nitrogen (N) after 3 days incubation in live (microbial immobilization and adsorption) and sterile (adsorption

only) agricultural pasture soils. Tracer $^{13}\text{C}^{15}\text{N}$ -labeled *E.coli* necromass was added (0.050 mg C and 0.016 mg N g⁻¹ soil fwt) in combination with different background (i.e. unlabeled) necromass (control, *E.coli*, *M. luteus* or *S. cerevisiae* (~5.5 mg C and ~1.5 mg N g⁻¹ soil fwt)). Sterile soils were either autoclaved or gamma-irradiated (results pooled) and thus presumably have a slightly higher level of background necromass C and N from the recently killed natural soil microbial community (~ 0.38 mg C and 0.030 mg N g⁻¹ soil fwt). 'Retention' is calculated from the amount of label remaining in the water-extracted soil residue at the end of a short-term (3 d) incubation. Upper and lower-case letters within each plot indicate a significant difference between live and sterile treatments, and different letters indicate significant difference between necromass treatments ($\alpha=0.05$).

Figure 3. Tracing the gaseous loss, biotic immobilization and carbon use efficiency of necromass tracer ($^{13}\text{C}^{15}\text{N}$ -*E.coli*) in live soils with different background necromass. Data presented as the mean (\pm standard error, n=5) % of applied $^{13}\text{C}^{15}\text{N}$ necromass in **(a)** CO₂, **(b)** microbial biomass ('MBC'), and **(c)** N₂O, and **(d)** the carbon use efficiency ('CUE') of necromass mineralization after 3 days incubation. Tracer $^{13}\text{C}^{15}\text{N}$ -labeled *E.coli* necromass was added (0.050 mg C and 0.016 mg N g⁻¹ soil fwt) in combination with different background (i.e. unlabeled) necromass (control, *E.coli*, *M. luteus* or *S. cerevisiae* (~5.5 mg C and ~1.5 mg N g⁻¹ soil fwt)). Different letters within each plot indicate significant difference between necromass treatments ($\alpha=0.05$).




Figure 1. Schematic diagram of microbial necromass fate in mineral soils. In this diagram, necromass is presented as a substrate for microbes, similar to plant inputs (litter or exudates). Microbes can acquire unassociated or mineral-associated necromass for substrate. Immobilization includes recycling of necromass by microbes into new biomass, and eventually necromass, with potential for some loss as CO₂. Stabilization is assumed to be through adsorption to mineral surfaces ('organic-mineral'), especially on fine silts and clays. In this study we hypothesize that this process is not limited to mineral surface availability ('organic-mineral'), but that necromass also adheres to necromass ('organic-organic'), promoting retention, and that this organic-organic process may be influenced by necromass cell chemistry.




Figure 2. The retention of necromass ($^{13}\text{C}^{15}\text{N}$ -*E.coli*) carbon and nitrogen in live and sterile soil with different background necromass. Data presented as the mean (\pm standard error, $n=5$) % retention of applied $^{13}\text{C}^{15}\text{N}$ necromass as **(a)** carbon (C) or **(b)** nitrogen (N) after 3 days incubation in live (microbial immobilization and adsorption) and sterile (adsorption only) agricultural pasture soils. Tracer $^{13}\text{C}^{15}\text{N}$ -labeled *E.coli* necromass was added (0.050 mg C and 0.016 mg N/g) in combination with different background (i.e. unlabeled) necromass (control, *E.coli*, *M. luteus* or *S. cerevisiae* (~ 5.5 mg C and ~ 1.5 mg N/g)). Sterile soils were either autoclaved or gamma-irradiated (results pooled) and thus presumably have a slightly higher level of background necromass C and N from the recently killed natural soil microbial community (~ 0.38 mg C and 0.030 mg N/g). ‘Retention’ is calculated from the amount of label remaining in the water-extracted soil residue at the end of a short-term (3 d) incubation. Upper and lower-case letters within each plot indicate a significant difference between live and sterile treatments, and different letters indicate significant difference between necromass treatments ($\alpha=0.05$).




Figure 3. Tracing the gaseous loss, biotic immobilization and carbon use efficiency of necromass tracer ($^{13}\text{C}^{15}\text{N}$ -*E.coli*) in live soils with different background necromass. Data presented as the mean (\pm standard error, $n=5$) % of applied $^{13}\text{C}^{15}\text{N}$ necromass in **(a)** CO_2 , **(b)** microbial biomass ('MBC'), and **(c)** N_2O , and **(d)** the carbon use efficiency ('CUE') of necromass mineralization after 3 days incubation. Tracer $^{13}\text{C}^{15}\text{N}$ -labeled *E.coli* necromass was added (0.050 mg C and 0.016 mg N/g) in combination with different background (i.e. unlabeled) necromass (control, *E.coli*, *M. luteus* or *S. cerevisiae* (~5.5 mg C and ~1.5 mg N/g)). Different letters within each plot indicate significant difference between necromass treatments ($\alpha=0.05$).

Supplementary Methods (online only)

To test our hypotheses, we used short-term (3 day) live and sterile soil incubations of clay loam soil (order: Chromic Eutric Albic Stagnosol; pH: 6.8; texture: 53 % sand, 19 % silt, 28 % clay; bulk density: 0.96 g soil cm⁻³; organic matter: 4.8 % C, 0.48 % N, 0.19 mg MBC (g dw soil)⁻¹) managed as an agricultural pasture (247 kg N ha⁻¹ y⁻¹ (slurry and manure), 2-3 mows y⁻¹, 0.21 livestock units ha⁻¹ y⁻¹ (cattle equivalent)), from northwest England (53°50'57"N, 2°47'01"W). The soil was collected from the top 10 cm of 1 m² plots separated by 20 m (n=5), sieved to 2 mm and adjusted to 60% water holding capacity, then subsampled to live and sterile treatments in 500 mL laboratory bottles fitted with septum-modified lids (γ -irradiated soils were sterilized in triple-layered polyethylene bags and added to sterile laboratory bottles at the start of the incubation). Soil sterilization was either via autoclaving (2x 120°C for 30 min) or γ -irradiation (>75kGy); we used two different methods to isolate effects of sterilization itself on C and N retention. Specifically, we were concerned that autoclaving more than γ -irradiation would release protected pools of C, confounding our results by providing new sites for C and N retention. We found no difference (P<0.05) between autoclaving and γ -irradiation, so we pool those data ('sterile') in our results (by averaging pairs to not inflate degrees of freedom; n=5). Necromass was made by growing cultures from streak plates in media (including unlabeled or labeled glucose (¹³C) and ammonium chloride (¹⁵N)), spinning down biomass to a pellet, rinsing, autoclaving twice, freeze-drying, subsampling the powder to aliquots and autoclaving again. The incubation included unlabeled necromass to enhance the background concentration of necromass in the soil: we assumed that soil necromass C is at least 50% of soil organic C (Liang et al., 2011) and doubled that amount by adding lab grown necromass (~5.5 mg C and ~1.5 mg N

g⁻¹ soil fw) from three functional group chemistries (Gram-negative and Gram-positive bacteria (*Escherichia coli* (%C: 44.2, %N: 13.9, C/N: 2.7) and *Micrococcus luteus* (%C: 40.3, %N: 11.5, C/N: 3.0)) and a yeast (*Saccharomyces cerevisiae* (%C: 41.8, %N: 9.6, C/N: 3.7))), and maintained a no-addition control. To quantify retention, we added a very small amount of isotopic tracer necromass consisting of ¹³C/¹⁵N labeled (10 atom %) *E. coli* necromass (0.050 mg C and 0.016 mg N g⁻¹ soil fw) to all samples, included a natural abundance control, and assessed retention of the tracer as a proxy for necromass retention. To maintain sterile conditions, in a laminar flow hood we processed the jars (60 g fw soil each) over the course of one day in groups of 6 jars (necromass-types and controls), processed 3 times (live/sterile treatments), repeated for each soil replicate (n=5). Sterile background necromass was added as a powder to addition treatments and all soils were lightly shaken to distribute the powder through soil or to mimic this addition in the controls. By jar, 3 mls of sterile labelled necromass solution or water was added, lids were immediately and tightly closed, and initial headspace gas samples were collected through a filtered syringe. Headspace gas samples were collected a total of 4 times throughout the 3-day incubation, plus a final gas sample was collected for isotope analysis. At the end of the incubation, we aseptically subsampled 20 g fw soil to chloroform-added and non-chloroform-added extraction tubes, and used chloroform direct extraction (Fierer et al., 2003), with 35 ml water as an extractant. Soil and extractant were shaken for 1 hr on a rotary shaker, centrifuged, and filtered, then sparged to remove excess chloroform. Extractable organic C (EOC) and extractable nitrogen (EN) of the sparged and blank-corrected chloroform extracts and water extract controls were used to calculate microbial biomass uptake of the label (measured in all but only detected in the live soils), on a Shimadzu TOC-L CPN (Shimadzu UK Ltd., UK). To measure substrate uptake into biomass,

chloroform extracts were freeze-dried, ground and analyzed for $\delta^{13}\text{C}$ on a Costech CN
streamed to a Picarro G-2131i CRDS. We analyzed CO_2 and N_2O concentrations in the
headspace gas samples on a Perkin-Elmer Autosystem XL gas chromatograph (GC) fit with a
methanizer and flame ionization detector (CO_2) and an electron capture detector (N_2O). The
 $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures of the headspace gas samples were analyzed on Picarro G2131-*i*
(CO_2) and Picarro G5131-*i* (N_2O) Isotope and Gas Concentration Analyzers (cavity ringdown
spectrometers). We used CO_2 , N_2O and MBC concentrations and their delta values in mixing
models to calculate labelled necromass incorporation into live-treatment CO_2 , N_2O , and
MBC, and the carbon use efficiency of the live microbes. The C, N, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of the post-
extraction soils (i.e. the extraction soil remaining on the filter) were used to assess total
abiotic and biotic retention. Soils were oven-dried, ground and analyzed for $\delta^{13}\text{C}$ on a
Costech CN Analyzer streamed to a Picarro G2131-*i* Isotope and Gas Concentration Analyzer,
and for $\delta^{15}\text{N}$ on a Carlo Erba NA1500 Elemental Analyzer coupled to a Dennis Leigh
Technologies IRMS. We used mixed models to assess the statistical significance ($\alpha=0.05$) of
our results with two fixed factors (live or sterilized, and added necromass type) and one
random effect (plot) using the function lmer (package lme4), Anova (car) and glht
(multcomp) in R (3.2.4).