



Predicting Spinal Condition using various biomechanical attributes of the pelvic and lumbar spine

Xueyao Wang, Ally Wardell, Liam Pongracz, and Lan Luo



Introduction

- Machine learning applications in exploratory orthopedic medicine research
- Random forest applications in spine research
- Support Vector Machine model usage in an orthopedic context.



Introduction: Spondylolisthesis and disk herniation

- Spondylolisthesis is a unilateral or bilateral defect (fracture or separation) in the vertebral pars interarticularis, usually in the lower lumbar vertebrae
- Symptoms: pain that spreads across the patient's lumbar region and radiates into posterior legs
- Disk Herniation is when the soft center of a spinal disk pushes into a crack in the tougher exterior casing
- Symptoms: Lower back pain (chronic)
- Both are often misdiagnosed



Introduction: Purpose of this study

- Accurately categorize spinal osteoarthritis conditions into the categories: normal, disk herniation, and spondylolisthesis based on biomechanical attributes of the shape and orientation of the pelvis and lumbar spine.
- Compare the accuracy of categorizations between support vector machine models and a random forest model
- An accurate machine learning algorithm could serve as an assistance tool for physicians to consult while considering a diagnosis, with the hopes of reducing misdiagnosis



Methods: The Data

- These data were obtained from a selection of three-hundred sixty patients during the medical residence period of Dr. Henrique da Mota, in the Group of Applied Research in Orthopedics (GARO), at the Centre Medico-Chirurgical de Réadaptation des Massues in Lyon, France
- 360 Patients: volunteers that were asymptomatic for disk herniation and Spondylolisthesis
- Spinal condition categorizations: Normal, Disk Hernia, and Spondylolisthesis




Methods: The predictors

- Six Predictors
 - Pelvic Incidence
 - Pelvic Tilt
 - Lumbar Lordosis Angle
 - Sacral Slope
 - Pelvic Radius
 - Grade of Spondylolisthesis

Characteristic	N	Hernia, N = 60 ¹	Normal, N = 100 ¹	Spondylolisthesis, N = 150 ¹	p-value ²
pelvic_incidence	310	48 (11)	52 (12)	72 (15)	<0.001
pelvic_tilt	310	17 (7)	13 (7)	21 (12)	<0.001
lumbar_lordosis_angle	310	35 (10)	44 (12)	64 (16)	<0.001
sacral_slope	310	30 (8)	39 (10)	51 (12)	<0.001
pelvic_radius	310	116 (9)	124 (9)	115 (16)	<0.001
degree_spondylolisthesis	310	2 (6)	2 (6)	52 (40)	<0.001

¹Mean (SD)

²One-way ANOVA





Methods: Modeling and Algorithms

- Compare a Random Forest and Support Vector Machine Model
- Random Forest model
 - 5-fold cross-validation used to train / test the algorithm and obtain CV errors and standard errors
 - Grid search to select the best parameter values (trees: 50, 250, 500 / predictors: 2, 3, 6)
 - The best parameters for a given training set were selected using out-of-bag MSE
- Support Vector Machine models
 - 5-fold cross-validation used to train / test the algorithm and obtain CV errors and standard errors
 - One model trained with a linear kernel; one model trained with a radial basis kernel
 - Grid of parameters were tested (epsilon: 0, 0.25, 0.50, 0.075, 1.00 / cost: 1-5)
- We wanted to find the best predictive model between these two approaches



Results: Overall Performance

	Hernia	Normal	Spondylolisthesis
SVM (Linear)	0.333(0.167)	0.170(0.057)	0.047(0.018)
SVM (RBF)	0.283(0.095)	0.180(0.110)	0.060(0.037)
Random Forest	0.433(0.199)	0.230(0.104)	0.040(0.043)

(1) Error (Standard Error)




Results: AUC (Hand and Till)


	AUC
SVM (Linear)	0.952
SVM (RBF)	0.944
Random Forest	0.923




SVM (Linear)




SVM (RBF)



Random Forest



Results: Gini importance





Discussion

- Our results indicate SVM-linear, SVM-RBF, and RF, show very promising performance.
 - Models differ in precision
- All three algorithms performed the best in terms of prediction accuracy on the spondylolisthesis condition, second on the normal condition, and the lowest on the disk hernia condition.
 - 60% accuracy across all three algorithms, despite disk hernia being low.
- An explanation for low prediction accuracy for disk herniation
- Overall promising results for prediction



Discussion

- Limitations
 - Some collinearity in predictors
 - Ex) pelvic incidence and sacral slope
 - Could reduce precision of estimation/classification, and in turn, statistical power
 - Grade of spondylolisthesis had a high mean decrease in Gini importance
 - Could dominate classification algorithm and bias estimation.
- Recommendations for the future:
 - Develop more condition specific classification algorithms (as opposed to a one-for-all-spinal-conditions classification), and be more cautious on predictions selection.
- Conclusion: The results of this study indicate that both the SVM and RF models have achieved a relatively satisfying level of precision in classifying spinal conditions.



Thank you for your attention!



References

- Boulay C , Bollini G, Legaye J, Tardieu C, Prat-Pradal D, Chabrol B, Jouve J, Duval-Beaupère G, Pélissier J, (2014). Pelvic Incidence: A Predictive Factor for Three-Dimensional Acetabular Orientation—A Preliminary Study, *Anatomy Research International*, vol. 2014, Article ID 594650, 9 pages <https://doi.org/10.1155/2014/594650>
- Cabitz F, Locoro A, Banfi G. (2018, Jun). Machine Learning in Orthopedics: A Literature Review. *Front Bioeng Biotechnol.* 6:75. doi:10.3389/fbioe.2018.00075
- Day J, Smidt G, Lehmann T, (1984, April). Effect of Pelvic Tilt on Standing Posture, *Physical Therapy*, Volume 64, Issue 4, Pages 510–516, <https://doi.org/10.1093/ptj/64.4.510>
- D’Hemecourt P, Micheli L. (2021) Spondylolysis and spondylolisthesis in child and adolescent athletes: Clinical presentation, imaging, and diagnosis. In: UpToDate, Post, TW (Ed).
- Drazin D, Hussain M, Harris J, Hao J, Phillips M, Kim T, Johnson J, & Bucklen B, (2015). The role of sacral slope in lumbosacral fusion: a biomechanical study, *Journal of Neurosurgery: Spine SPI*, 23(6), 754-762. Retrieved Apr 30, 2021, from <https://theins.org/spine/view/journals/j-neurosurg-spine/23/6/article-p754.xml>
- Galbusera F, Casaroli G, Bassani T. (2019, Mar). Artificial intelligence and machine learning in spine research. *JOR Spine*. 2019;2(1):e1044. doi:10.1002/jsp2.1044
- Hendler N, (2016). Why Chronic Pain Patients are Misdiagnosed 40 to 80% of the Time?. *Journal on Recent Advances in Pain*. 2. 94-98. 10.5005/jp-journals-10046-0051.
- Kotti M, Duffell L, Faisal A, McGregor A, (2017). Detecting knee osteoarthritis and its discriminating parameters using random forests. *Med. Eng. Phys.* 43, 19–29. 10.1016/j.medengphy.2017.02.004
- Legaye J, Duval-Beaupère G, Hecquet J, Marty C. (1998). Pelvic incidence: a fundamental pelvic parameter for three-dimensional regulation of spinal sagittal curves. *Eur Spine J* ;7(2):99-103. doi: 10.1007/s005860050038. PMID: 9629932; PMCID: PMC3611230.
- Nijeweme -d’Hollosy W, van Velsen L, Soer R, Hermens H. (2016) Design of a web- based clinical decision support system for guiding patients with low back pain to the best next step in primary healthcare *Proceedings of the 9th International Joint Conference on Biomedical Engineering Systems and Technologies (BIOSTEC 2016)*. Cham: Springer International Publishing
- Oktay A, Akgul Y. (2013). Simultaneous localization of lumbar vertebrae and intervertebral discs with SVM- based MRF. *IEEE Trans Biomed Eng*;60:2375- 2383
- R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,
- Seoud L, Adankon M, Labelle H, Dansereau J, Cheriet F. (2010) Prediction of scoliosis curve type based on the analysis of trunk surface topography *Biomedical Imaging: From Nano to Macro, 2010 IEEE International Symposium*. Piscataway, NJ: Institute of Electrical and Electronics Engineers (IEEE); 10.1109/ISBI.2010.5490322.
- Symou E, Tsitsopoulos PP, Marinopoulos D, Tsonidis C, Anagnostopoulos I, Tsitsopoulos PD. (2010) Spondylolysis: a review and reappraisal. *Hippokratia* ;14(1):17-21.
- UCI Machine Learning. (2017). Biomechanical Features of Orthopedic Patients. Retrieved April 04, 2021. From <https://www.kaggle.com/uciml/biomechanical-features-of-orthopedic-patients>
- Zhou Z, Hou C, Li D, Si J, (2018). The pelvic radius technique in the assessment of spinopelvic sagittal alignment of degenerative spondylolisthesis and lumbar spinal stenosis, *Journal of Orthopaedic Science*, Volume 23, Issue 6, Pages 902-907, ISSN 0949-2658, <https://doi.org/10.1016/j.jos.2018.06.021>.