#### 1

## Discrete Assignment

# Karyampudi Meghana Sai EE23BTECH11031

### PROBLEM STATEMENT

The ratio of the A.M and G.M of two positive numbers a and b is m: n. Show that a:  $b = (m + \sqrt{m^2 - n^2}) : (m - \sqrt{m^2 - n^2})$ .

### SOLUTION

Expressing A.M and G.M in terms of a and b:

$$\frac{a+b}{2\sqrt{ab}} = \frac{m}{n} \tag{1}$$

Let's assume that  $x = \sqrt{\frac{a}{b}}$ . Then, we have:

$$\frac{a}{b} = x^2 \tag{2}$$

Substituting the value of x in equation (1):

$$\frac{1+x^2}{2x} = \frac{m}{n} \tag{3}$$

$$\frac{1}{x} + x = \frac{2m}{n} \tag{4}$$

$$x^2 - \frac{2m}{n}x + 1 = 0\tag{5}$$

$$\implies x = \frac{m}{n} \pm \frac{\sqrt{m^2 - n^2}}{n} \tag{6}$$

Since  $x = \sqrt{\frac{a}{b}}$ , x must be positive.

$$x = \frac{m + \sqrt{m^2 - n^2}}{n} \tag{7}$$

Referencing the value of x from equation(2).

$$\frac{a}{b} = \left(\frac{m + \sqrt{m^2 - n^2}}{n}\right)^2 \tag{8}$$

Multiplying both the numerator and denominator with  $(m - \sqrt{m^2 - n^2})$ :

$$\frac{a}{b} = \frac{1}{n^2} \frac{\left(m + \sqrt{m^2 - n^2}\right)^2 \left(m - \sqrt{m^2 - n^2}\right)}{\left(m - \sqrt{m^2 - n^2}\right)} \tag{9}$$

$$\implies a: b = (m + \sqrt{m^2 - n^2}): (m - \sqrt{m^2 - n^2})$$
 (10)

nth term of the AP:

$$t(n) = [a + n(b - a)] u(n)$$
(11)

$$n^{k}u(n) \xrightarrow{Z} (-1)^{k} z^{k} \frac{d^{k}}{dz^{k}} U(z)$$
(12)

$$u(n) \xrightarrow{Z} \frac{1}{(1-z^{-1})} \quad |z| > 1$$
 (13)

$$nu(n) \xrightarrow{Z} \frac{z^{-1}}{(1-z^{-1})^2} \quad |z| > 1$$
 (14)

Referencing the equations from (13),(14).

$$t(n) \xrightarrow{Z} \frac{a}{(1-z^{-1})} + \frac{(b-a)z^{-1}}{(1-z^{-1})^2} \quad |z| > 1$$
 (15)

nth term of the GP:

$$t(n) = a \left(\frac{b}{a}\right)^n u(n) \tag{16}$$

$$r^{k}u(n) \xrightarrow{Z} \frac{1}{(1-rz^{-1})} \quad |z| > |r|$$
 (17)

Referencing the equation from (17).

$$t(n) \xrightarrow{Z} \frac{a^2 z^{-1}}{(a - b z^{-1})} \quad |z| > \left| \frac{b}{a} \right| \tag{18}$$