



# 非コンパクト空間上の準古典解析とシュレディンガー作用素のスペクトル理論

日時: 2025 年 9 月 3 日–9 月 5 日

会場: 京都大学数理解析研究所 420 号室

代表者: 三上 溪太

## プログラム

### 9 月 3 日

- 13:00–14:00 **Serge Richard** (名古屋大学)  
*Spectral and scattering theory in a discrete setting 1*
- 14:10–15:10 **藤家 雪朗** (立命館大学)  
*Semiclassical analysis for matrix-valued operators and applications to spectral asymptotics 1*
- 15:40–16:40 **Serge Richard** (名古屋大学)  
*Spectral and scattering theory in a discrete setting 2*

### 9 月 4 日

- 9:30–10:30 **藤家 雪朗** (立命館大学)  
*Semiclassical analysis for matrix-valued operators and applications to spectral asymptotics 2*
- 10:40–11:40 **福嶋 翔太** (群馬大学)  
*Mathematical analysis of transmission problem across ideal and resistive interface 1*
- 13:00–14:00 **藤家 雪朗** (立命館大学)  
*Semiclassical analysis for matrix-valued operators and applications to spectral asymptotics 3*
- 14:40–15:40 **亀岡 健太郎** (立命館大学)  
*Semiclassical shape resonances for Stark and magnetic Stark Hamiltonians 1*
- 15:50–16:50 **藤家 雪朗** (立命館大学)  
*Semiclassical analysis for matrix-valued operators and applications to spectral asymptotics 4*

### 9 月 5 日

- 9:30–10:30 **福嶋 翔太** (群馬大学)  
*Mathematical analysis of transmission problem across ideal and resistive interface 2*
- 10:40–11:40 **亀岡 健太郎** (立命館大学)  
*Semiclassical shape resonances for Stark and magnetic Stark Hamiltonians 2*

## タイトルとアブストラクト

藤家 雪朗 (立命館大学)

**Title:** Semiclassical analysis for matrix-valued operators and applications to spectral asymptotics

**Abstracts:** In this course we consider Hermitian matrix-valued semiclassical pseudo-differential operators, mainly  $2 \times 2$  matrix-valued operator with two Schrödinger operators on the diagonal and small of order  $h$  interactions on the off-diagonal (called *coupled Schrödinger operator* in the following). Each eigenvalue of the matrix-valued symbol defines a Hamiltonian system. We focus on the problem arising from a *crossing* of two or more Hamiltonian flows.

The course consists of three parts.

The first part is devoted to the basic theory of semiclassical pseudodifferential operators. The goal of this part is a theorem of propagation of semiclassical wave front set. The real principal type condition in the well known scalar case is generalized here to the *microhyperbolic* condition in the sense of Ivrii. A crossing point of Hamiltonian flows is microhyperbolic if there exists a scalar function increasing along these flows near the crossing point.

From the second part, we restrict ourselves, for more detailed study, to a one-dimensional coupled Schrödinger operator. We study the semiclassical asymptotic behavior of the solutions microlocally near a crossing point. It is expressed as what we call *microlocal scattering matrix*, which relates the WKB solutions on the incoming Hamiltonian flows with the ones on the outgoing flows. We will see that the first term of the semiclassical asymptotic expansion of this matrix is an identity and that the second term is off-diagonal, corresponding to the change of flows, of order  $h^{\frac{1}{m+1}}$  where  $m$  is the contact order of the crossing.

In the last part, we apply the previous studies to the semiclassical asymptotics of eigenvalues or resonances of a coupled Schroedinger operator with crossings. The Hamiltonian flows make a directed graph when we regard the crossing points as vertices. The WKB solutions on each edge (Hamiltonian flow) together with the microlocal scattering matrices at each vertex enable us to define a monodromy matrix of the microlocal solutions on the graph. Eigenvalues or resonances are then characterized in terms of this matrix.

福寛 翔太 (群馬大学)

**Title:** Mathematical analysis of transmission problem across ideal and resistive interface

**Abstracts:** We consider electrostatic or thermal systems with conductors placed in Euclidean space. Conductors have three main characteristics to be

controlled: shape, conductivity and interface resistance. If the conductivity is locally constant, then the system is modelled by the Laplace equation, with these three characteristics incorporated into the transmission conditions. In this talk, we deal with two types of conductors. One is a conductor with negative conductivity (plus small imaginary part) and ideal (non-resistive) interface, which motivates the spectral analysis of Neumann-Poincaré operator. The other is a perfect conductor with constant interface resistance. In this case, the solution exhibits distinct behavior against the ideal interface case, although these two cases are connected by the limit as the interface resistance tends to zero. This talk is based on joint work with Yong-Gwan Ji (KIAS), Hyeonbae Kang (Inha University), Xiaofei Li (Zhejiang University of Technology) and Yoshihisa Miyanishi (Shinshu University).

### 亀岡 健太郎 (立命館大学)

**Title:** Semiclassical shape resonances for Stark and magnetic Stark Hamiltonians

**Abstracts:** We consider resonances of Stark and magnetic Stark Hamiltonians. We explain the definition of resonances through the complex distortion of Hamiltonians. Then we assume that there are potential wells and study the semiclassical behavior of resonances which are exponentially close to the real axis. The magnetic Stark Hamiltonian part is based on joint work with Naoya Yoshida.

### Serge Richard (名古屋大学)

**Title:** Spectral and scattering theory in a discrete setting

**Abstracts:** During these two presentations, we shall start by recalling the framework of topological crystals, which corresponds to the most general  $\mathbb{Z}^d$  periodic discrete setting. The spectral analysis of the Laplace operator or of the adjacency matrix will then be briefly sketched, and the scattering theory for perturbations of these operators will be considered. Perturbed systems are allowed to have an infinite number of new edges, as long as suitable decay conditions are imposed. In the final part of the presentation, discrete time evolution groups will also be introduced and various concepts of scattering theory will be further illustrated.