
Measurement of cross sections and couplings of the Higgs boson in the WW decay channel using the ATLAS detector

corrinne mills

(University of Edinburgh)



On behalf of the ATLAS collaboration

ICHEP


4 July 2014

Higgs boson production


- Higgs boson couplings measurement essential SM consistency test
→ *Search for rare production modes*
- Continue search for additional states at high mass in parallel



associated production ("VH")




$H \rightarrow WW \rightarrow l\nu l\nu$




WW channel well-suited to exploring rare production modes and searching at high-mass

Large branching ratio and good S/B from clean dilepton signature



channel	$\sigma(pp \rightarrow H)$ @ 8 TeV	BR ($H \rightarrow VV$)	BR($VV \rightarrow 4l$)	evt. yield in 20 fb^{-1}	S/B (approx.)
$H \rightarrow \gamma\gamma$		0.0023	-	1000	0.03 – 0.5
$H \rightarrow ZZ \rightarrow 4l$	22.3 pb	0.026	0.0011	130	2
$H \rightarrow WW \rightarrow l\nu l\nu$		0.22	0.047	4700	0.1 – 0.4

The WW signature




jets depend on production mode

spin zero resonance

Can't go bump hunting like $\gamma\gamma$ or $ZZ \rightarrow 4l$ channels

Transverse mass M_T : invariant mass without p_z of vs → signal discriminant with edge at m_H

Final state
2 charged leptons + 2 neutrinos
 $(l\nu jj)$ important as well for high m_H searches)




$$M_T^2 = (E_T^{\ell\ell} + E_T^{\text{miss}})^2 - (\vec{p}_T^{\ell\ell} + \vec{E}_T^{\text{miss}})^2$$


$$(E_T^{\ell\ell})^2 = (\vec{p}_T^{\ell\ell})^2 + (m_{\ell\ell})^2$$

Backgrounds


WW: quasi-irreducible




top: has b-jets




W+jets: fake leptons



$Z/\gamma^* \rightarrow \tau\tau \rightarrow l\nu\nu l\nu\nu$





Z/ γ^* : no ν s




Analysis strategy

Starting point: dilepton data after missing transverse energy cut



1. Categorize events by number of jets, number of leptons, lepton flavors
 → Separate by production mode and background composition
2. Cut away backgrounds and normalize to control regions enriched in a particular background but orthogonal to the signal region

WW – quasi-irreducible



WW is dominant ggF background
7.4% total uncertainty in 0-jet
 → 1.6% from theory

1) Normalize to data


2) subtract other contributions

3) extrapolate using simulation

$$N_{SR} = \left(\frac{N_{SR}^{MC}}{N_{CR}^{MC}} \right) (N_{CR}^{\text{data}} - N_{CR}^{\text{other}})$$


Dilepton invariant mass m_{ll} is a good signal discriminant because the spin-0 of the Higgs boson, combined with V-A structure of W decay correlates lepton directions

W+X backgrounds




- **W+jets: isolation / ID on 2nd lepton separates SR and CR**
- Transfer factor for extrapolation to SR from dijet data
- Stringent lepton ID / isolation → “fake” leptons, particularly muons, are often true leptons from heavy-flavor hadron decays
→ 40-45% uncertainty on fake factor from sample composition

- “**Same-sign validation region**”
- **WZ , $W\gamma^*$, $W\gamma$** estimated from theory + MC simulation
 - 14% total uncertainty in 0-jet category
 - Z , γ , and γ^* produce lepton with same or opposite charge as W with equal probability




The raw data





0 jet




1 jet



2 jet (VBF)



lνlν signal strength




Combined $WW \rightarrow l\nu l\nu$ signal strength $\mu = \sigma/\sigma_{\text{SM}}$ (2011+2012, all Njet):

$$\mu = 0.99 \pm 0.21 \text{ (stat.)} \pm 0.21 \text{ (sys)}$$

VBF vs. ggF: virtual contributions for ggF only

Sensitive to BSM particles in the loops



$$\mu_{\text{ggF}} = 0.8 \pm 0.2 \text{ (stat.)} \pm 0.3 \text{ (syst.)}$$

$$\mu_{\text{VBF}} = 1.7 \pm 0.7 \text{ (stat.)} \pm 0.4 \text{ (syst.)}$$

Cross section measurement

Uncertainties on the signal strength $\mu = \sigma/\sigma_{\text{SM}}$

Category	Source	Uncertainty, up (%)	Uncertainty, down (%)
Statistical	Observed data	+21	-21
Theoretical	Signal yield ($\sigma \cdot \mathcal{B}$)	+12 → +12 ← -9	-9 ← -12
Theoretical	WW normalisation	+12	-12
Experimental	Objects and DY estimation	+9	-8
Theoretical	Signal acceptance	+9	-7
Experimental	MC statistics	+7	-7
Experimental	W+jets fake factor	+5	-5
Theoretical	Backgrounds, excluding WW	+5	-4
Luminosity	Integrated luminosity	+4	-4
Total		+32	-29

- $\mu = \sigma/\sigma_{\text{SM}}$; measured cross section $\sigma = \mu \times \sigma_{\text{SM}}$
- Remove uncertainties on signal cross section but not acceptance

$$(\sigma \cdot \text{Br}(H \rightarrow WW))_{\text{obs, 8 TeV}} = 6.0 \pm 1.1 \text{ (stat.)} \pm 0.8 \text{ (theo.)} \pm 0.7 \text{ (expt.)} \pm 0.3 \text{ (lumi.) pb}$$

$$= 6.0 \pm 1.6 \text{ pb}$$


$$(\sigma \cdot \text{Br}(H \rightarrow WW))_{\text{exp, 8 TeV}} = 4.8 \pm 0.7 \text{ pb}$$

VH, H → WW Signal Regions


Clean 3-lepton (WH) and 4-lepton (ZH) signatures,
additional info on HWW vertex
(complement to VBF mode)

4-lepton: 2 OS pairs, one with
 $|m_{||} - m_Z| < 10 \text{ GeV}$
1 event expected, S/B $\sim 1/5$
→ zero observed

WH: Z-enriched category (OS same-flavor pair)



WH: Z-depleted category (no OS same-flavor pair)




minimum ΔR between any two leptons: same physics as $\Delta\varphi(\text{ll})$ and $m_{||}$

VH, H → WW Results

cross section limits (@95% confidence) and significance of excess over BG for $m_H=125$ GeV

	expected	observed	significance
WH (8 TeV)	$5.2 \times \sigma_{SM}$	$10 \times \sigma_{SM}$	2.3σ ($p_0 = 1.2\%$)
ZH (8 TeV)	$9.6 \times \sigma_{SM}$	$14 \times \sigma_{SM}$	1.5σ ($p_0 = 7.2\%$)
Combined (7+8 TeV)	$3.6 \times \sigma_{SM}$	$7.2 \times \sigma_{SM}$	2.0σ ($p_0 = 2.1\%$)

- No SM sensitivity yet but observed limit higher than expected one





The Road Ahead

- WW was essential to discovery, combined rate measurement, and connecting the new particle to electroweak symmetry breaking
- Now: Is this the **SM Higgs boson or something more interesting?**
 - *WW channel can address this from multiple angles*
 - + Good S/B and large branching ratio
 - - Challenging backgrounds in ggF and VBF
 - *ggF and VBF modes key inputs to coupling measurement combinations*
 - *Search for rarer production modes (WH, VH, ttH...)*
 - *Properties measurements (starting from spin)*
 - *Search at high mass for additional states*
- 13 TeV run (2015-2017) will bring much-needed data and stronger statements in response to all of these questions

backup



Spin measurement

- Test spin hypothesis with kinematic variables through a BDT
 - *No direct measurement of angles (cannot reconstruct rest frame)*
- Use ggF $e\mu$ zero-jet category → highest sensitivity
 - *Then loosen requirements on spin-sensitive variables: $m_{||} < 80$ GeV, $p_T^{||} > 20$, $\Delta\phi(II) < 2.8$*



Spin measurement


- Train two BDTs
 - One for SM signal (0^+) against background
 - Second for alternate spin/parity hypothesis ($2^+, 1^+, 1^-$) against background
- Fit data to discriminate hypotheses
- 2^+ excluded at $\geq 95\%$ CL by WW



Drell Yan (Z/γ^*)

Measure missing transverse energy
only with tracks: more robust against
extra interactions (pileup)

$$\vec{E}_T^{\text{miss}} = - \sum_{\text{objects } i} \vec{p}_T^{\text{object}}$$



- Left: additional rejection after initial E_T^{miss} requirement
→ *First inclusion of ee+μμ channels in 2012 WW analysis*


The ATLAS Detector

Muon Detectors: large radii for precise momentum measurement

Precision: Drift tubes (MDT) and Cathode Strip Chambers (CSC)

Trigger: Resistive Plate Chamber (RPC) and Thin Gap Chamber

(TGC)



Toroids

Inner Detector (tracking):

Pixels (silicon)

SCT (silicon)


TRT (straw)

Hadronic calorimeter (jets, hadrons)

Steel absorber + scintillator

LAr with copper/tungsten absorber


Top background to VBF



≥ 2 -jet data after a b-veto: still top-dominated

Reject further background with unique VBF signature: **energetic well-separated jets**

These cuts reject additional **95% of total background but cost 70% of VBF signal**




Estimating top in VBF phase space

Uncertainties on top normalization

stat.	theory	exp.	total
10%	15%	29%	39%

- Normalize to control region (like WW)
 - Apply all VBF jet cuts, extrapolate from b -tagged → b -vetoed



Normalize to data

subtract other processes

$N_{SR} = \left(\frac{N_{SR}^{MC}}{N_{CR}^{MC}} \right) \left(N_{CR}^{\text{data}} - N_{CR}^{\text{other}} \right)$


extrapolate using simulation

Full systematic uncertainties

Source	Signal processes (%)			Background processes (%)		
	$N_{\text{jet}} = 0$	$N_{\text{jet}} = 1$	$N_{\text{jet}} \geq 2$	$N_{\text{jet}} = 0$	$N_{\text{jet}} = 1$	$N_{\text{jet}} \geq 2$
Theoretical uncertainties						
QCD scale for ggF signal for $N_{\text{jet}} \geq 0$	13	-	-	-	-	-
QCD scale for ggF signal for $N_{\text{jet}} \geq 1$	10	27	-	-	-	-
QCD scale for ggF signal for $N_{\text{jet}} \geq 2$	-	15	4	-	-	-
QCD scale for ggF signal for $N_{\text{jet}} \geq 3$	-	-	4	-	-	-
Parton shower and UE model (signal only)	3	10	5	-	-	-
PDF model	8	7	3	1	1	1
$H \rightarrow WW$ branching ratio	4	4	4	-	-	-
QCD scale (acceptance)	4	4	3	-	-	-
WW normalisation	-	-	-	1	2	4
Experimental uncertainties						
Jet energy scale and resolution	5	2	6	2	3	7
b -tagging efficiency	-	-	-	-	7	2
f_{recoil} efficiency	1	1	-	4	2	-

$Z/\gamma^* \rightarrow \tau\tau \rightarrow |\nu\nu| |\nu\nu|$

Reduce uncertainties by normalizing dominant background to data in signal-depleted “control regions” (CR)



Normalize to data

subtract other processes


$N_{SR} = \left(\frac{N_{SR}^{MC}}{N_{CR}^{MC}} \right) (N_{CR}^{\text{data}} - N_{CR}^{\text{other}})$

extrapolate using simulation


Azimuthal lepton opening angle
 $\Delta\varphi(\text{ll})$ works well for the same reasons

1-jet analysis: top and the b-veto

control region: b-jet tag



signal region: b-jet veto




Extrapolate from tag to veto

Experimental systematics dominate (b -tag efficiency calibration)

High-mass / BSM WW searches

- High-mass searches now explicitly BSM
- Include SM-like, narrow-width, 2HDM signal models

Combine ggF and VBF channels for SM-like hypothesis (assume production cross section ratio unchanged from SM)



2HDM: general BSM Higgs model, neural-net based search

heavy H masses up to ~200 GeV excluded
(model + mixing angle dependent)

