ONE Clinical Bladder Health Protocol

POSSIBLE F-TAG/QMS

Resident Rights

- Self-Determination

Quality of Life

- ADL Maintenance

Pharmacy

- Drug Regimen is Free from Unnecessary Drugs

Urinary Incontinence

Quality Measures

- Falls with Major Injury
- UTI
- Catheter
- Claims Based:
  - ER Visits
  - Hospitalizations

THERAPIES

- **PT**
- **OT**

PURPOSE OF THERAPY

- Help our residents with UI to be as safe and independent as possible with utilization of strategies, exercises, and modalities

GOAL EXAMPLES

PT/OT:

- Patient will manage incontinence products independently 5/5 sessions.
- Patient will demonstrate toilet transfers from WC using grab bar with modified independence using grab bar.
- Patient will stand for two minutes to complete clothing management before and after toileting with SBA 5/5 sessions.
- Patient will demonstrate ability to complete hygiene after toileting with minimal assist for 5/5 sessions to decrease risk for compromised skin integrity.
- Patient will perform sit to stand transfers without episodes of UI for 5/5 sessions.
- Patient will decrease nighttime trips to toilet from six to four to improve safety.
- Patient will decrease episodes of UI from 5 to 3 per day for 5/5 days.
- Patient will not have UI with simulated coughing/sneezing 5/5 sessions.
- Patient will decrease bladder irritant intake from four to two servings per day.

TOOLS & ASSESSMENTS

TOOLS & ASSESSMENTS

- Intake Diary
- Voiding Diary
- Pelvic and Low Back Pain Rating
- Posture Assessment
- Sitting & Standing Tolerance Time
- Diastasis Recti Test
- Berg Balance
- Tinetti Balance
- TUG
- OAB Assessment Tool
- OABSS

POSSIBLE F-TAG/QMS	THERAPIES	PURPOSE OF THERAPY	GOAL EXAMPLES	TOOLS & ASSESSMENTS
			Patient will increase water intake from 20 ounces to 60 ounces per day. Patient will demonstrate belly breath and initial upright transverse activation with 50% tactile cues to improve sitting tolerance. Patient will report 4/10 pain after sitting with in WC with back support for 30 minutes. Pt will demonstrate increased standing tolerance with less pain and improved balance after 10 min of active sitting with feet on the vibration plate. Pt will demonstrate increased standing tolerance with less pain and improved balance after 10 min of active sitting with feet on the vibration plate. Pt will be able to exhale and engage core while independently changing positions in bed and moving from supine to sit with SBA reporting low back pain <3/10. Patient's severity level rating of urinary incontinence will decrease to 8/10 to increase participation in functional activities and decrease fear of going places. Patient will decrease the number of incontinence episodes at night to 2 in order to improve sleep patterns, facilitate safe functional mobility and decrease risk for falls. Patient will exhibit a decrease in the number of incontinent urge episodes per day to 9 to decrease fear related to incontinence, facilitate safe functional mobility, and improve functional task performance.	**Implementation:** Core and pelvic floor muscle strengthening Bladder irritant avoidance training Fluid intake/hydration training Bladder emptying strategy training PTNM (Percutaneous Tibial Neuromodulation) PENS Utilization MFAC Utilization Clothing Management Toileting Hygiene Toilet Transfers Toileting Schedule