Comparing World Quality of Life Measures: Nonparametric vs. Parametric Approaches

Katherine M. Prioli December 20, 2018

Abstract

BACKGROUND Increasing globalization has generated interest in comparing countries by key quality of life (QoL) measures such as life expectancy, gender equality, and happiness, among others. When considering how countries compare by QoL, it is vital to understand how the measures are related, if at all, and in what ways the statistical tests chosen affect the results. This study aimed to explore the distributions of and relationships between QoL measures using both nonparametric and parametric approaches, and to assess which methods were appropriate. METHODS A dataset containing country names and selected QoL measures for 2016 was explored through descriptive statistics and sensible univariate and bivariate visualizations. Five sets of hypotheses pertaining to relationships within the data were generated and tested with both nonparametric methods and their parametric equivalents, and results of these parallel analyses and the assumptions they require were compared. RESULTS Most variables in the study were not normally distributed. Nonparametric methods dominated, performing acceptably for all five cases, whereas parametric methods were acceptable in only two of the five cases. CONCLUSION Nonparametric methods represent a powerful, important toolset for use in analyzing data that fails the assumptions needed for traditional parametric methods.

Background

With increasing economic globalization, a natural topic of interest is how the world's nations compare with respect to quality of life (QoL). Several organizations monitor global QoL indicators and report single-dimension or aggregate values for indicators of interest. For example, the World Bank reports Gross Domestic Product (GDP), which is a single-dimension indicator often strongly predictive of QoL in a given country (The World Bank (2018)). Additionally, the World Health Organization reports infant mortality rate, life expectancy at birth, and life expectancy at 60 years of age (World Health Organization (2018b), World Health Organization (2018a)).

Other quality of life measures represent compound scores or indices based on several inputs. For example, the United Nations calculates an annual Human Development Index (HDI), representing the developmental level of each country on a scale of zero to one based on several factors, including life expectancy at birth, years of schooling, and per-capita income (The United Nations Development Programme (2018b)). The HDI also categorizes countries into four levels of development (low, medium, high, and very high). Similarly, the Social Progress Imperative publishes the Social Progress Index (SPI), ranging from 0 to 100, and comprising over 50 dimensions in three broad categories: basic human needs (e.g., nutrition, safety), foundations of wellbeing (e.g., basic knowledge, environmental quality), and opportunity (e.g., personal rights, freedoms) (Social Progress Imperative (2018b)). The World Economic Forum's Global Gender Gap Index reports a gender equality index, scaled from 0-1, based on measurements of gender-related gaps in such dimensions as economic participation, level of education, health and survival, and political offices held (World Economic Forum (2016b)). Finally, the World Happiness Report calculates a score from 0-10 by considering per-capita GDP, healthy life expectancy, social support, freedoms, and perception of corruption, among others (Helliwell, Layard, and Sachs (2018)).

When studying QoL by country, it is important to understand how each QoL measure is distributed and how they may be interrelated. The objective of this analysis was to explore the distributions of and relationships between key QoL indicators using both nonparametric and parametric methods, and to assess the appropriateness of each method used.

Methods

$Data\ Exploration$

The dataset used in this analysis, titled alldata, was generated for the MAT 8790 course (Prioli (2018b)). It consists of country-level variables for calendar year 2016 as described in Table 1.

Table 1. alldata dataframe contents.

| Source | Variable Name | Description |
|---------------------|---------------|---------------|
| countrycode package | country | Country names |

| Source | Variable Name | Description |
|------------------------------|---------------|-------------------------------------------------------------|
| Social Progress Imperative | SPI | Social Progress Index value (scale of 0:100) |
| (2018a) | | |
| The World Bank (2018) | logGDP | Gross Domestic Product, log transform (valued in \$US 2018) |
| The United Nations | HDIrank | Human Development Index ranking |
| Development Programme | | |
| (2018a) | | |
| The United Nations | HDIindex | HDI index value (scale of 0:1) |
| Development Programme | | |
| (2018a) | | |
| The United Nations | HDI_cat | HDI index category (4 levels) |
| Development Programme | | |
| (2018a) | | |
| Helliwell, Layard, and Sachs | happiness | World Happiness Score (scale of 0:10) |
| (2018) | | |
| World Economic Forum (2016a) | gendereq | Gender Equality Index (scale of 0:1) |
| World Health Organization | infantmort | Infant mortality rate |
| (2018b) | | |
| World Health Organization | birth_MF | Life expectancy at birth, males & females |
| (2018a) | | |
| World Health Organization | sixty_MF | Life expectancy at 60 years, males & females |
| (2018a) | | |

All variables represent data from calendar year 2016. Countries having missing data were omitted from the dataset to ensure that the tests of interest could be performed.

For each variable except country, descriptive statistics were run and a sensible visualization was generated, following which a correlation matrix was produced to examine pairwise relationships between continuous variables.

Nonparametric and Parametric Analyses

Based on the data exploration results, five sets of formal hypotheses were generated about the data (Table 2), and sensible nonparametric tests and their parametric equivalents were chosen to test these hypotheses.

Table 2. Analyses Performed

| Analysis | Variable(s) | Null Hypothesis | Alternative Hypothesis | Nonparametric Test | Parametric Test |
|----------|--------------------------|----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|----------------------------------------------|-------------------------------|
| 1 | HDIindex | (No formal hypothesis test) Nonparametric 95% confidence interval for the population mean | (No formal hypothesis test) Parametric 95% confidence interval for the population mean | Bootstrapping | One-Sample t-Test |
| 2 | HDIindex, SPI | Human development and social progress are not associated | Human development and social progress are correlated | Kendall's Tau | Pearson's Correlation Test |
| 3 | logGDP, gendereq | There is no relationship between log(GDP) and gender equality index | There is a relationship between log(GDP) and gender equality index | Hoeffding's Test | Pearson's Correlation Test |
| 4 | happiness | Happiness is normally distributed | Happiness is not normally distributed | One-Sample Kolmogorov- Smirnov Test | Shapiro-Wilk Test |
| 5 | HDI_cat, log(infantmort) | Log infant mortality rate is the same across levels of human development | Log infant mortality rate differs by level of human development | Permutation F-Test | ANOVA |

In Analysis #1, bootstrapping and the one-sample t-test were employed to find nonparametric and parametric 95% confidence intervals (CIs) for the population mean. The widths of these intervals were compared, and assumptions required for the t-test were assessed. The bootstrapping CI for the mean was found using 1,000 sampling iterations with replacement via $\mathtt{sample}()$, and the CI for the mean was found via $\mathtt{t.test}()$.

Analysis #2 was motivated by the correlation matrix analysis, which indicates a linear relationship is likely between SPI and HDIindex. Kendall's Tau was chosen as the nonparametric test, with the usual Pearson's test as the parametric alternative. Both tests assume continuous data and are equipped to detect linear dependence, and employed two-sided alternatives. Additionally, because simulation is the preferred method when ties are present for Kendall's Tau, the variables were first assessed for ties before selecting a method for Kendall's Tau. The competing hypotheses for this analysis were $H_0: \tau = 0$ vs. $H_A: \tau \neq 0$ (for Kendall's Tau) and $H_0: \rho = 0$ vs. $H_A: \rho \neq 0$ (for Pearson's Test). Both tests were carried out via cor.test() using method = "kendall" or method = "pearson" as appropriate.

Correlation tests were chosen for Analysis #3 to determine whether log GDP and gender equality were associated. No obvious relationship was seen between these two variables on the correlation plot. Hoeffding's Test was selected because it is sensitive to any departure from independence, and Pearson's test was chosen as the parametric comparator. Both tests employed two-sided alternatives and assume continuous data, and Pearson's test additionally assumes a linear relationship. For Hoeffding's test, the formal hypotheses tested were $H_0: \Delta = 0$ vs. $H_A: \Delta \neq 0$ where

$$\Delta = \int_{X \in x} \int_{Y \in y} [F_{XY}(x, y) - F_X(x)F_Y(y)]^2 dF_{XY}(x, y).$$

For Pearson's test, the competing hypotheses were $H_0: \rho = 0$ versus $H_A: \rho \neq 0$. These tests were carried out using testforDEP::testforDEP() with the options test = "HOEFFD" and test = "PEARSON" respectively.

Analysis #4 was chosen because happiness appears to have a symmetric, possibly normal distribution on univariate analysis. To test for normality, the Lilliefors test (a normality-specific version of the one-sample Kolmogorov-Smirnov test) was chosen as the nonparametric test, and the Shapiro-Wilk test was chosen as the parametric alternative. Both assume a continuous sample distribution. The formal hypotheses for this analysis were H_0 : happiness is normally distributed vs. for H_A : happiness is not normally distributed. Lilliefors testing was performed via nortest::lillie.test(), and Shapiro-Wilk testing was carried out using shapiro.test().

To test for a difference in infant mortality rate by human development category, Analysis #5 employed the permutation F-test and ANOVA as the nonparametric and parametric tests respectively. Both tests assume samples are independent with constant variance, and ANOVA additionally assumes a normal distribution. The log transform of infant mortality rate was used to ensure that the assumption of constant variance would be met. The competing hypotheses were H_0 : the cumulative distribution functions (CDFs) for log infant mortality were equal across all four human development levels vs. H_A : at least one of these CDFs differs. The permutation F-test was performed via jmuOutlier::perm.f.test() with 1,000 simulations because the sample size was prohibitively large for calculating an exact p-value. The aov() function was used for ANOVA testing.

Results of these five paired analyses were compared in the context of the data and assumptions needed. All tests were performed at level $\alpha = 0.05$ in R. The dataset, full code, and this report are available in an online repository (Prioli (2018a)).

Results

$Data\ Exploration$


The original dataset represented 205 countries, and the analytic dataset contained 111 countries after omitting those with missing values. Of the 94 countries omitted, 26 were highly developed by HDI category, 20 were in the low developmental category, and 18 had unknown HDI category (Table 3).

Table 3. Omitted Countries by HDI Category

| HDI Category | Frequency |
|--------------|-----------|
| Low | 20 |
| Medium | 18 |
| High | 26 |
| Very High | 12 |
| NA | 18 |


Descriptive statistics and univariate visualizations are presented in Figs. 1-8. Figure 1 depicts the distribution and descriptive statistics of the HDIindex variable, along with counts by country for the HDI_cat variable. HDIindex has a very nonnormal appearance, yet has similar mean and median, and a small standard deviation as compared to the mean. The highest category of human development is the most represented in the data.

Figure 1. Human Development Index




SPI, described in Fig. 2, appears to have a multimodal distribution, with mean and median quite similar in value.

Figure 2. Social Progress Index Distribution



The distribution of logGDP (Fig. 3) appears possibly bimodal with mean and median reasonably close in value and standard deviation smaller than both.

Figure 3. Gross Domestic Product Distribution, Log Transform




Per Fig. 4, the World Happiness Score appears roughly normally distributed.

Figure 4. Happiness Score Distribution

Min 2.903
Q1 4.624
Median 5.578
Mean 5.553
Q3 6.339
Max 7.66
SD 1.139

Happiness Score

Figure 5 shows a roughly symmetric, possibly trimodal distribution for gendereq, with mean and median very close in value, and standard deviation approximately a tenth of either measure of location.



The probability of infant mortality is heavily right-skewed per Fig. 6.




Figure 7 explores total life expectancy at birth vs. age 60. Life expectancy at birth appears left-skewed, but total life expectancy at sixty looks reasonably symmetric. These data indicate that, across all countries, a person who survives until at least 60 years has a longer total life expectancy than does the general world population at birth.





Figure 8 depoits pairwise relationships between these variables. Strong positive linear relationships are seen between HDIindex and SPI, HDIindex and birth_MF, and SPI and birth_MF, indicating that higher levels of social progress and human development generally correlate with a longer life expectancy at birth. Additionally, positive linear relationships are seen between happiness and birth_MF, between happiness and HDIindex, and between happiness and SPI, indicating that increasing levels of human development and social progress generally lead to increased happiness (though there is considerable spread in the data), and that happiness and longevity may also be correlated.

Strong negative relationships are seen between infantmort and birth_MF, between HDIindex and infantmort, and between SPI and infantmort, though the latter two of these may not necessarily be linear. These indicate that increasing human development and social progress levels correlate with lower levels of infant mortality, and that as infant mortality decreases, life expectancy at birth increases.

Figure 8. Correlation Matrix



Nonparametric and Parametric Analyses


In Analysis #1, the bootstrapping approach yielded a 95% CI for the mean of (0.7157, 0.7698), having width 0.0541. The parametric approach via t.test() yielded a 95% CI given by (0.7163, 0.7728), having width 0.0565. The nonparametric method gave the smaller confidence interval, having 95.75% the width of the CI obtained through the parametric method. The one-sample t-test assumes that the sample is random with independent draws, and that the data are continuous and normally distributed. Independence and continuous data are met, but per Fig. 1, normality is violated, so the parametric approach is not appropriate. The nonparametric approach is acceptable.

For Analysis #2, ties were not present in either SPI or HDIindex; thus, Kendall's Tau was tested using the usual cor.test(x, y, method = "kendall") approach. When testing for independence in both the nonparametric and parametrics cases, the p-values were less than 0.0001, with correlation coefficients $r_{\tau} = 0.843$ and r = 0.966 respectively. Both are statistically significant and represent strong evidence of a linear relationship between social progress and human developmental level, thus H_0 was rejected with the conclusion that, at the $\alpha = 0.05$ level, social progress and human development are correlated. Both tests assume continuous data, which is satisfied, and both yield the same p-value. Both the nonparametric and parametric methods are appropriate.

When testing for independence in Analysis #3, Hoeffding's test returned $p_{Hoeffding} = 0.0384$, whereas Pearson's test yielded $p_{Pearson} = 0.093$. Hoeffding's test led to the rejection of H_0 (i.e., independence) and the conclusion that, at the $\alpha = 0.05$ level, there is sufficient evidence of an association between log(GDP) and gender equality. Pearson's test, however, resulted in retaining H_0 and concluding that, at the $\alpha = 0.05$ level, there is insufficient evidence of an association between log(GDP) and gender equality. The assumptions of continuous data were met for both tests, but the assumption of linearity failed for Pearson's test, thus the parametric method is inappropriate.

In Analysis #4, the Lilliefors test yielded p-value $p_{Lilliefors} = 0.624$, whereas the p-value for the Shapiro-Wilk test was $p_{SW} = 0.129$. Both exceed alpha; thus, H_0 was retained for both tests with the conclusion that there is insufficient evidence to assert that happiness is not normally distributed. Both tests assume a continuous distribution, which is satisfied by the data, and the tests agree, so either the nonparametric or parametric approach is acceptable in this case. Notably, although neither p-value is significant, p_{SW} was much smaller than $p_{Lilliefors}$. Since the objective for this analysis was to test for normality, the parametric test is more appropriate.

Both the permutation F-test and ANOVA yielded p < 0.0001 for Analysis #5. Since $p < \alpha$, H_0 was rejected for both the nonparametric and parametric cases, leading to the conclusion that there exists a difference in infant mortality by HDI category. The assumptions of independence and constant variance are satisfied by the data, the latter having p-value 0.4327 on Levene's test for homogeneity of variances. However, per Fig. 9, the ANOVA assumption of normality may be violated for the Medium category. Therefore, the parametric method is inappropriate in this case, and the nonparametric method is retained.



Discussion

Nonparametric methods are appropriate when analyzing data that violates the assumption of normality, and they can also be applied to normal data, but may be less powerful than the traditional parametric tests. The dataset used in this study consists of real-world QoL data, many variables of which fail normality. Across all five analyses, the nonparametric methods performed acceptably, whereas the parametric equivalents were only acceptable for two of the analyses.

THINK ABOUT WHAT ELSE TO ADD TO THE DISCUSSION - ADDRESS HOW UNDERSTANDING THE RELATIONSHIPS BETWEEN QOL MEASURES IS USEFUL

Limitations

An important limitation to this analysis is that countries with missing data were excluded from the analytic dataset. This may introduce bias, as a nontrivial proportion of countries omitted belong to the low and medium HDI developmental categories, and these countries are likely to perform poorly on the other QoL measures considered.

Another limitation is the inherent interrelatedness of some of these variables, particularly the compound indices. For example, one factor considered in the Human Development Index is life expectancy, so it is unsurprising that a strong association between HDIindex and birth_MF was seen in the correlation matrix. Similarly, some subitems of the Social Progress Index overlap with those included in the Gender Equality Index. Correlation tests are possible in such cases but may be inappropriate unless the underlying measures are fundamentally different.

Finally, this study is limited by the variables included in this dataset. There exist many other QoL measures that would have been worth including in this analysis, such as measures of educational attainment, civilian indebtedness, and population-level health utilities. Future work may entail expanding the study to include such variables.

Conclusion

Nonparametric methods represent an important toolset when working with data that violates normality. This analysis gives evidence that they can perform favorably as compared to their parametric equivalents, and should be considered when analyzing data that is unsuitable for parametric approaches.

References

