

recent posts | user list | about | contact

Search

research education mechanician opinion software industry conference job video

# **User login**

Username \*

Password \*

- Create new account
- Request new password

Log in

## **Navigation**

Search iMechanica

## **Recent blog posts**

- Postdoc/Research Fellow for computational mechanics of all solid state battery at the Department of Mechanical Engineering, National University of Singapore
- Two Postdoc positions for acoustics at the Department of Mechanical Engineering, National University of Singapore

#### Home » Blogs » ramdas chennamsetti's blog


## Thin plate theory...

Thu, 2007-05-24 14:13 - ramdas chennamsetti

Hi all!

I have a small doubt in the assumptions made in thin plate theory.

We make some of the following assumptions in thin plate theory (Kirchoff's classical plate theory) (KCPT).



[2] The vertical deflection 'w' is not a function of 'z' => dw/dz = 0

Now there are three stress components sigma(x), sigma(y) and sigma(xy). The other three stress components sigma(z), sigma(xz) and sigma(yz). This is like a plane stress.

But, from the second assumption ez=0 (strain in z-direction) and from the above exz=0 and eyz=0.

Then, this leads to plane strain.

From the constitutive equation for 'ez' => sigma(x) + sigma(y) = 0

But, this doesn't happen.....I am looking for explanations ...

Thanks in advance.

» ramdas chennamsetti's blog | Log in or register to post comments | 17566 reads

#### **Comments**

the constitutive equation is not used **Permalink** Submitted by **Ying Li** on Thu, 2007-05-24 14:17.



# Quick guide

- Ask iMechanica
- Having difficulty with posting comments?
- How to add an image
- How to post an entry
- Journal Club
- Lecture notes
- RSS Feeds
- FAQ

#### **Recent comments**

- A proposal to reform the national law for the ASN procedure 3 hours 16 min
- Cartesian vs Natural 8 hours 45 min ago
- my resignation has been revoked !!! 1 day 1 hour ago
- A facebook group of 14.500 members will help you in applying! 3 days 4 hours ago
- Thanks you Francisco for your 5 days 5 hours ago
- Thank you Jessica for your 5 days 5 hours ago
- correction to step b 5 days 6 hours ago
- global to natural 5 days 6 hours ago

- Postdoc position for a computational mechanician in the Computational Physics Group @ University of Michigan
- SES Symposium 8.9 Mechanics, Materials,
   Manufacture and Device
   Innovations of Soft
   Electronics Abstracts
   Due Monday May 16
- PhD Position in Mechanical Engineering at Temple University for Fall 2022
- EMI Announces the Winners of Prestigious 2022 Mechanics Awards
- SES Symposium 6.8 Mechanics of
   Electrochemical Systems
   - Abstracts Due Monday
   May 16
- PhD positions opening in contact mechanics at PoliBa
- Centenary of two pioneering theories in mechanics
- Postdoc openings in Mechanics at Central South University

More

# What we talked about

- ABAQUS UMAT
- ABAQUS tutorial
- Cauchy stress
- Geometry & Mechanics
- Large elastic deformation
- Logarithmic strain

In the Kirchoff's classical plate theory, such a constitutive equation is not considered. So, you needn't to use it.

Lee

» Log in or register to post comments



## R. Chennamsetti, Scientist,

Permalink Submitted by ramdas chennamsetti on Thu, 2007-05-24 14:39.

R. Chennamsetti, Scientist, India

Hi Lee,

Thank you. But, constituvie equatations are to be satisfied. In this case they relate the stresses and strains.

- R Chennamsetti
- » Log in or register to post comments

# Kirchhoff's assumptions.

Permalink Submitted by Erkan Oterkus on Tue, 2007-05-29 20:41.

This is just my idea and I cannot claim that it is exactly true. After these assumptions, we do not have any dependency on z (out-of plane direction) coordinate. So, we only end up with a problem on an in-plane surface. Therefore, we should only concentrate on x-y components of

stress and strain. What this means, we should not take into account the relations related with the z-coordinate. So, we do not need to take into account the constituve relation of <code>ezz</code> and corresponding stress components.

Erkan Oterkus.

» Log in or register to post comments

## R. Chennamsetti, R&DE(E),

Permalink Submitted by ramdas chennamsetti on Wed, 2007-05-30 00:53.

R. Chennamsetti, R&DE(E), INDIA

Hi Erkan,



- Thanks Emma for the very 5 days 8 hours ago
- Thanks Emma for such an insightful discussion 5 days 9 hours ago

#### More comments

comments at a glance

## **Popular content**

#### Today's:

- I share the vision of iMechanica, but am not ready to post anything, should I register?
- How to apply to Italian professorship positions -a little guide
- thermodynamics of nanoscale small systems
- Generate master curve for viscoelasticity
- InternationInternational Journal of Applied Mechanics (IJAM) Vol.2 No.3
- Abaqus step definition
- Rolling Moment
  Resistance of Particles on
  Surfaces
- Postdoc position for a computational mechanician in the Computational Physics Group @ University of Michigan
- Three-day intensive course "Invitation to nondestructive



- Mechanics of growth
- Mesh-free methods
- Objective rates of stress
- Plastic potential
- Poroelasticity
- Stress and strain
- Temperature
- Timoshenko lectures
- Viscoelasticity
- Why is rubber incompressible?
- Writing a paper
- Writing a proposal
- in situ Mechanics
- tensor
- tensor and its invariants

#### **Sites of interest**

- AAM
- AMD
- Basic energy science
- Buckled Shells
- CFD online
- EMI
- Electroactive polymers
- IUTAM
- PolymerFEM
- PoroNet
- SES
- Soft Matter World
- USNC/TAM
- eFluids

Thank you. Here relations means 'Constitutive relation?' But, we can't violate the constitutive relation in any direction (I think).

- Ramdas
- » Log in or register to post comments

### We estimated Kirchoff

Permalink Submitted by TarasVasiliev on Wed, 2007-05-30 01:05.

We estimated Kirchoff theory by solving spatial problem for small hight h=H/R (<0.1).

It can be concluded that all normal stresses and deflection give asimptoticaly right values.

But shear stress orz must be approximated only by theories wich includes shear.

For example Timoshenko-Reissner theory.

More detailed information about useful boundaries in Kirchoff theory see

http://www.springerlink.com/content/y725477x15885301/?p=3bd506ccac034d2f...

» Log in or register to post comments

# Kirchhoff's assumptions -2

Permalink Submitted by Erkan Oterkus on Wed, 2007-05-30 01:50.

Hi Ramdas,

Thank you for this interesting topic. What I mean with the term relations is in general. Maybe I should even correct my phrase. Under the assumptions that we are making, we should eliminate the terms related with the z-coordinate. As a crude example, like eliminating applied boundary conditions from our global governing equation. So,, we should only have the in-plane related terms in any relation suh as equilibrium equations, constitutive relations, etc. You are right, we should not violate constitutive relation, but here we are making assumptions, so we need to sacrifice something which is OK to ignore under some particular conditions.

Erkan.

» Log in or register to post comments

- evaluation and structural health monitoring" by Zahra Sharif Khodaei, 24--26 May 2022, Czech Republic, Prague
- Postdoc/Research Fellow for computational mechanics of all solid state battery at the Department of Mechanical Engineering, National University of Singapore

# **Syndicate**



# The Conflicting Kirchhoff Assumptions

Permalink Submitted by Wenbin Yu on Fri, 2008-09-19 15:10.

The original post made a point. The violation of the 3D constitutive relations is happened because the second assumption is not correct. The first assumption is asymptotically correct for the first approximation of the original 3D model using a 2D model. However, the second assumption, transverse normal remains rigid, is clearly violating the first assumption, plane stress. Because by assuming plane stress, we assume that the plate is free to move in the thickness direction, which means the transverse normal is not rigid. Both assumptions can be valid only if the Poisson's ratio is zero. Recall the well-known and readily observed Poission's effect. The reason the second assumption is used is because it is convient to derive a 2D version kinematics (strain-displacement relations). It is noted that same conflicting assumptions also used to derive beam models dealing with tension and bending: section remain rigid in its own plane and the beam is in uniaxial stress state.

As a final comment, both assumptions are not absolutely needed for one to derive a plate theory. One can use the variational asymptotic method to take advantage of the smallness of theory as the small parameter to reduce the 3D model to a 2D model with the first assumption comes out as a result of the first assumption and the transverse displacement will be a quadratic function of the thickness coordinate. Please refer to the following paper for more details.

Yu, W.: "Mathematical Construction of a Reissner-Mindlin Plate Theory for Composite Laminates," *International Journal of Solids and Structures*, vol.42, no. 26, 2005, pp. 6680-6699. (pdf)

» Log in or register to post comments



Each entry is © copyright 2006-2014 by the individual user and can be used in accordance with the **Creative Commons License**. iMechanica is powered by **Drupal**, and hosted at the **Harvard School of Engineering and Applied Sciences**.