


Honors Multivariable Calculus : : Homework 1x

April 4, 2024

Ahmed Saad Sabit, Rice University

Problem 01

Aid for my brain: Let the compact region be $R \in \mathbb{R}^{n-1}$ where $n = 3$. Common sense tells us R is basically a disk in \mathbb{R}^2 or for now $x - y$ plane if we want what is in the figure. The tip of the cone is \vec{a} . I think from the question our R doesn't really need to be necessarily a disk. Now the region is all the lines that join from \vec{a} to R . If $\vec{x} \in R$ then considering a linear map $\gamma_{\vec{x}} : [0, 1] \rightarrow \mathbb{R}^n$ such that $\gamma_{\vec{x}}(0) = \vec{a}$ and $\gamma_{\vec{x}}(1) = \vec{x} \in R$



The line segment is set of all points such that,

$$\Gamma = \{s \in \mathbb{R}^n : s = \vec{a}t + (1 - t)\vec{x} \text{ where } t \in [0, 1], \vec{x} \in R, \vec{a} \in \mathbb{R}^n\}$$

Here $x \in R$. Every point $\vec{p} \in \Gamma$ is a member of the cone.

The volume of the region Γ (which is the defined cone) is going to be,

$$\int_{\Gamma} 1 = \text{Volume}$$

Now the burden is to find a region Γ .

The trick we are going to apply is to translate a relatively more simple shape into our cone shape by the following definition.

Definition 1. Define the D region to be $R \times [0, 1]$. Volume of this shape is

$$\int_D 1 = \text{vol}(R)$$

We need to find a transformation $T : D \rightarrow \Gamma$. From definition of cone we can write,

$$T(x_1, x_2, \dots, x_{n-1}, x_n) = x_n(\vec{a} - (x_1, \dots, x_{n-1}, 0)) + (x_1, \dots, x_{n-1}, 0)$$

In vector form,

$$T(x_1, \dots, x_n) = \begin{pmatrix} x_n a_1 - x_n x_1 + x_1 \\ \vdots \\ x_n a_{n-1} - x_n x_{n-1} + x_{n-1} \\ x_n a_n \end{pmatrix}$$

T is injective (check appendix after this problem). So we can compute,

$$dT = \begin{bmatrix} 1 - x_n & 0 & \cdots & 0 & a_1 - x_1 \\ 0 & 1 - x_n & \cdots & 0 & a_2 - x_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 - x_n & a_{n-1} - x_{n-1} \\ 0 & 0 & \cdots & 0 & a_n \end{bmatrix}$$

dT being an upper triangular matrix,

$$\det(dT) = (1 - x_n)^{n-1} a_n$$

Now through change of coordinates we can find the integral,

$$\int_D 1 = \int_\Gamma 1 \cdot T |\det(dT)|$$

Because $1 \cdot T = 1$,

$$\begin{aligned} &= \int_\Gamma |\det(dT)| \\ &= \int_\Gamma |(1 - x_n)^{n-1} a_n| \\ &= \int_\Gamma (1 - x_n)^{n-1} a_n \quad (\text{integration over } x_n > 0) \\ &= a_n \int_\Gamma (1 - x_n)^{n-1} \\ &= a_n \int_R \int_0^1 (1 - x_n)^{n-1} dx_n \end{aligned}$$

We can carry out the integration using $u = 1 - x_n$ substitution method and we can get,

$$a_n \text{vol}(R) \left[\frac{u^n}{n} \right]_0^1$$

Hence,

$$\boxed{\frac{a_n}{n} \text{vol}(R)}$$

T is injective

For this transformation to be valid, T must be injective other than places that are content zero. We can see that T is injective everywhere on its domain except for the point where the $x_n = 1$.

T is only valid if $x_n = 1$ “place” or $R \times \{1\}$ is content zero in \mathbb{R}^n .

Consider $f : R \subset \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ given by $f(\vec{x}) = 1$. Note that the graph of f is precise $R \times \{1\}$. Since our region R is compact we know that the graph of a continuous function is content zero. Therefore $R \times \{1\}$ is content zero.

Problem 02

Let D denote the region described by the n dimensional solid ellipsoid

$$\frac{x_1^2}{a_1^2} + \cdots + \frac{x_n^2}{a_n^2} \leq 1$$

Here a_i are positive. The volume is,

$$\int_D 1$$

We need to do a transformation now, so define $T : D' \rightarrow D$ defined by

$$T(x_1, \dots, x_n) = (a_1 x_1, \dots, a_n x_n)$$

where D' is the unit ball in \mathbb{R}^n . Then note that T is surjective to D and injective on it's domain.

$$dT = \begin{bmatrix} a_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & a_n \end{bmatrix}$$

So easily,

$$\det(dT) = a_1 \cdots a_n$$

Now we perform the change of coordinates and compute,

$$\int_D 1 = \int_{D'} 1 \cdot |T| \det(dT)$$

Carrying out the computation we simply have,

$$\int_{D'} |a_1 \cdots a_n|$$

So from here,

$$= (a_1 \cdots a_n) (\text{volume of the unit ball in } \mathbb{R}^n)$$

$$(a_1 \cdots a_n) \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2} + 1)}$$

Problem 03

Denote the region enclosed by the blue curve and red line by D . The average value of f on this region is,

$$\bar{f} = \frac{\int_D f}{\text{vol}(D)}$$

We first need to compute $\int_D f$. So consider,

$$T : D' \rightarrow D \quad T(r, \theta) = (r \cos \theta, r \sin \theta)$$

From here we can see,

$$|\det(dT)| = r$$

We can compute the integral $\int_D f$ as,

$$\int_D f = \int_{D'} f \cdot T \cdot r = \int_{D'} (r^2 \cos^2 \theta + r^2 \sin^2 \theta) r$$

From here,

$$\int_{D'} r^3 = \int_0^{2\pi} \int_0^\theta r^3 dr d\theta$$

Computing the integral gives,

$$\frac{8}{5}\pi^5$$

Now we are required to evaluate the volume.

$$\int_D 1 = \int_{D'} (1 \cdot T) \cdot r = \int_0^{2\pi} \int_0^\theta r dr d\theta$$

Solving this gives us,

$$\frac{4}{3}\pi^3$$

So the division of the two results gives us the average value of f on D ,

$$\frac{6}{5}\pi^2$$

Problem 04

Similar to the first problem let's start by a simple volume first, define the region $R \times [0, 2\pi]$. From here we define

$$T : R \times [0, 2\pi] \rightarrow D$$

with

$$T(x, y, z) = (x, y \cos z, y \sin z)$$

Note how T is a surjective map onto D and is injective at every points except the content zero portion that is $z = 0$ and $z = 2\pi$.

We will now compute,

$$dT = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos z & -y \sin z \\ 0 & \sin z & y \cos z \end{pmatrix}$$

From here,

$$\det(dT) = 1 \cdot \det \begin{pmatrix} \cos z & -y \sin z \\ \sin z & y \cos z \end{pmatrix}$$

$$\det(dT) = y$$

Now we can carry out this integration,

$$\int_D 1 = \int_{R \times [0, 2\pi]} 1 \cdot |T| \det(dT) = \int y = \int_R \int_0^{2\pi} y dz$$

Solving this get's us,

$$2\pi \int_R y$$

Now from here we compute,

$$2\pi \cdot \text{avg of } y \text{ on } R \cdot \text{vol}(R)$$

We get,

$$2\pi \int_R y$$

From this we showed that,

$$\text{vol}(D) = \int_D 1 = 2\pi(\text{avg of } y \text{ on } R) \cdot \text{vol}(R)$$

1 Problem 05

The unit ball on \mathbb{R}^4 is defined by

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 \leq 1$$

Let denote D on this region the unit ball, and to find the unit ball volume we find $\int_D 1$.

The coordinates can be changed by the following $T : D' \rightarrow D$ through,

$$T(r_1, \theta, r_2, \theta_2) = (r_1 \cos \theta, r_1 \sin \theta, r_2 \cos \theta_2, r_2 \sin \theta_2)$$

where $D' = S_1 \times [0, 2\pi], S_2 \times [0, 2\pi]$ such that, $\forall r_1 \in S$ and $\forall r_2 \in S_2$ we have $r_1^2 + r_2^2 \leq 1$ and $r_1, r_2 \geq 0$.

$$dT = \begin{bmatrix} \cos \theta_1 & -r_1 \sin \theta_1 & 0 & 0 \\ \sin \theta_1 & r_1 \cos \theta_1 & 0 & 0 \\ 0 & 0 & \cos \theta_2 & -r_2 \sin \theta_2 \\ 0 & 0 & \sin \theta_2 & r_2 \cos \theta_2 \end{bmatrix}$$

Now that dT is a block matrix, therefore we calculate its determinant as,

$$\begin{aligned} \det(dT) &= \det \begin{pmatrix} \cos \theta_1 & -r_1 \sin \theta_1 \\ \sin \theta_1 & r_1 \cos \theta_1 \end{pmatrix} \cdot \det \begin{pmatrix} \cos \theta_2 & -r_2 \sin \theta_2 \\ \sin \theta_2 & r_2 \cos \theta_2 \end{pmatrix} \\ &= (r_1 \cos^2 \theta_1 + r_1 \sin^2 \theta_1) (r_2 \cos^2 \theta_2 + r_2 \sin^2 \theta_2) = r_1 r_2 \end{aligned}$$

Through change of variables,

$$\begin{aligned} \int_D 1 &= \int_{D'} 1 \cdot |T| |\det(dT)| \\ &= \int_{D'} r_1 r_2 \end{aligned}$$

This integral can be computed,

$$\begin{aligned} \int_{D'} r_1 r_2 &= \int_{S_1 \times S_2} \int_0^{2\pi} \int_0^{2\pi} r_1 r_2 d\theta_1 d\theta_2 \\ &= \int_{S_1 \times S_2} r_1 r_2 (4\pi^2) \end{aligned}$$

Inside this region we have $r_1^2 + r_2^2 \leq 1$. This is a unit circle and through change of ariables again,

$$G : S \rightarrow S_1 \times S_2$$

$$G(R, \alpha) = (R \cos \alpha, R \sin \alpha)$$

where $S = [0, 1] \times [0, \frac{\pi}{2}]$ we constrain the input $\alpha \in [0, \frac{\pi}{2}]$ upto content zero.

$$dG = \begin{bmatrix} \cos \alpha & -R \sin \alpha \\ \sin \alpha & R \cos \alpha \end{bmatrix}$$

$$\det(dG) = R \cos^2 \alpha + R \sin^2 \alpha = R$$

Then by the change of variables it follows,

$$\begin{aligned} \int_{S_1 \times S_2} r_1 r_2 &= \int_S (R \cos \alpha)(R \sin \alpha) \cdot R \\ &= \int_0^{\pi/2} \int_0^1 R^3 \cos \alpha \sin \alpha dR d\alpha \end{aligned}$$

Completing this integral we find,

$$\boxed{\frac{1}{8}}$$

Using this in our previous integral,

$$\int_{D'} r_1 r_2 = 4\pi^2 \left(\frac{1}{8} \right) = \frac{\pi^2}{2}$$

Hence the volume is $\pi^2/2$.