

# Pseudoscalar top-Higgs coupling: Exploration of CP-odd observables to resolve the sign ambiguity

Nicolas Mileo,<sup>1,\*</sup> Ken Kiers,<sup>2,†</sup> Alejandro Szynkman,<sup>1,‡</sup> Daniel Crane,<sup>2,§</sup> and Ethan Gegner<sup>2,¶</sup>

<sup>1</sup>*IFLP, CONICET – Dpto. de Física, Universidad Nacional de La Plata, C.C. 67, 1900 La Plata, Argentina*

<sup>2</sup>*Physics and Engineering Department, Taylor University, 236 West Reade Ave., Upland, IN 46989, USA*

(Dated: February 27, 2016)

## Abstract

We present a collection of CP-odd observables for the process  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$  that are linearly dependent on the scalar ( $\kappa_t$ ) and pseudoscalar ( $\tilde{\kappa}_t$ ) top-Higgs coupling and hence sensitive to the corresponding relative sign. The proposed observables are based on triple product (TP) structures that we extract from the expression of the differential cross section in terms of the spin vectors of the top and antitop quarks. In order to explore other possibilities, we progressively modify these TPs, first by combining them, and then by replacing the spin vectors by the lepton momenta or the  $t$  and  $\bar{t}$  momenta by their visible parts. Assuming an integrated luminosity that is consistent with that envisioned for the HL-LHC, we find that the most promising observable can disentangle the hypotheses  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$  by more than the  $\sim 20\sigma$  statistical level. In the case of observables that do not need the reconstruction of the  $t$  and  $\bar{t}$  momenta, the power of discrimination is up to the  $\sim 16\sigma$  statistical level for the same number of events. We also show that the capability of the most promising observables for separating the CP-mixed hypotheses prevails even when a number of events plausible within the short term LHC is considered.

---

\* mileo@fisica.unlp.edu.ar

† knkiers@taylor.edu

‡ szynkman@fisica.unlp.edu.ar

§ dkcrane@mtu.edu

¶ ethan\_gegner@taylor.edu

## I. INTRODUCTION

After the discovery of a new boson  $H$  by the ATLAS [?] and CMS [?] collaborations, it has become of crucial importance to determine its physical properties with the highest possible precision. The study of the new boson's couplings to fermions is of great relevance and will allow us to better understand this particle's CP-transformation properties, as well as the extent to which this particle is consistent with the Higgs boson predicted by the Standard Model (SM) of particle physics. It is of particular importance to test the coupling of the putative Higgs boson to the top quark. This coupling governs the main Higgs boson production mechanism (which proceeds via gluon fusion) and it contributes to the important Higgs boson decay mode to two photons. It is also involved in the scalar-field naturalness problem – giving rise to the leading dependence on the cut-off energy scale in the corrections to the Higgs mass – and it may play an important role in the mechanism for electroweak symmetry breaking.

Given that the main Higgs boson production process is dominated by a top quark loop and that the diphoton and digluon decay channels are also mediated by a top loop, these processes provide constraints on the scalar and pseudoscalar  $tH$  couplings,  $\kappa_t$  and  $\tilde{\kappa}_t$  [? ? ? ?]. However, these constraints assume that there are no other sources contributing to the corresponding effective couplings; furthermore, in the case of the diphoton decay channel (which also involves a  $W$  boson loop), it is also assumed that the coupling of the Higgs boson to the  $W$  is standard. In this sense, the constraints derived from measurements of Higgs boson production and decay rates are indirect constraints. Electric dipole moments can also impose stringent indirect constraints on  $\tilde{\kappa}_t$  by assuming that there are no new physics (NP) particles contributing to the loops of the relevant diagrams and in the case of the EDM of the electron that the electron-Higgs coupling is that predicted by the SM [? ?]. In order to probe the  $tH$  coupling directly, processes with smaller cross sections need to be taken into account.

In contrast to the  $\tau H$  coupling, which can be studied through the decay  $H \rightarrow \tau^+ \tau^-$  [?], the  $tH$  coupling can only be tested directly via production processes, since the Higgs boson is kinematically forbidden from decaying to a  $t\bar{t}$  pair. Two types of processes are of particular interest in this regard – the production of a Higgs boson in association with a  $t\bar{t}$  pair and in association with a single top or antitop. The cross section for associated Higgs production with a single top (antitop) is smaller than that for production with a  $t\bar{t}$  pair, and involves the interference between a diagram in which the Higgs is radiated from the top (antitop) leg and one with the Higgs emitted from the intermediate virtual  $W$  boson. Interestingly, this implies that the constraints on  $\kappa_t$  and  $\tilde{\kappa}_t$  derived from  $tH$  and  $\bar{t}H$  production are dependent on the assumption made regarding the coupling of the Higgs boson to the  $W$  gauge boson,  $\kappa_W$ . Nevertheless, it is important to note that the interference between the above mentioned diagrams can be exploited to determine the relative sign between  $\kappa_t$  and  $\kappa_W$  (see for example Ref. [?]). Associated Higgs production with a  $t\bar{t}$  pair has been studied by several authors, and various observables sensitive to the couplings  $\kappa_t$  and  $\tilde{\kappa}_t$  have been proposed. Examples of such observables (all of which are CP-even) are the cross section, invariant mass distributions, the transverse Higgs momentum distribution and the azimuthal angular separation between the  $t$  and  $\bar{t}$ , to name a few [? ? ? ? ?]. Also, an approach based on weighted moments and optimal observables has been developed in Ref. [? ? ? ?] to discriminate the hypothesis of a CP-even Higgs from that of a CP-mixed state within the context of an  $e^+e^-$  as well as a  $pp$  collider. Now, CP-even observables are not sensitive to the relative sign between the scalar and pseudoscalar couplings  $\kappa_t$  and  $\tilde{\kappa}_t$ . Such observables are quadratically dependent on these couplings and thus only provide an indirect measure of CP violation. In order to be sensitive

to the relative sign between  $\kappa_t$  and  $\tilde{\kappa}_t$ , CP-odd observables must be considered.

Since the top quark decays before it can hadronize, its spin information is passed on to the angular distributions of its decay products in such a way that these particles work as spin analyzers. As is well known, in the case of semileptonic top decay, the charged lepton is the most powerful in this regard. It is also known that the top quark and antiquark spins are highly correlated in  $t\bar{t}$  production, a feature that is manifested in the double angular distributions of the decay products of the  $t$  and  $\bar{t}$  systems [? ? ? ?]. In the case of  $t\bar{t}H$  associated production, the  $t\bar{t}$  spin correlations are also sensitive to the manner in which the top couples to the Higgs boson. In fact, observables that exploit the differences in the  $t\bar{t}$  spin configurations were used in Ref. [?] to improve the discrimination of the  $t\bar{t}H$  signal from the dominant irreducible background  $t\bar{t}b\bar{b}$ , which does not involve the Higgs boson.

In this paper, we define a set of observables that are linearly dependent on  $\kappa_t$  and  $\tilde{\kappa}_t$  and are thus sensitive to the relative sign of these couplings. The proposed observables are based on a particular set of triple product (TP) structures that we extract naturally from the expression for the differential cross section for  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$ , making use of the fact that the  $t$  and  $\bar{t}$  decay products contain spin information and are sensitive to the nature of the  $tH$  coupling, as noted above. By using spinor techniques we relate the top and antitop spin vectors to final state particle momenta and separate the production process from the decay. This allows to identify straightforwardly the contributions linearly sensitive to the couplings. Further, the TP correlations in these contributions incorporate the  $t$  and  $\bar{t}$  spin vectors. Starting with these TPs, we not only recover the observables given in [?] but also propose additional possibilities that increase the sensitivity. In order to establish a hierarchy in the sensitivity of the TPs under analysis we investigate three different types of observables by using simulated events: asymmetries, mean values and angular distributions. We note that TP correlations have been used in [?] in the context of top-quark production and decay and in [?] in the framework of anomalous color dipole operators.

The remainder of this paper is organized as follows. In Sec. II we study the theoretical framework for the process  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$  and derive a general expression for the differential cross section from which a first set of TP correlations is extracted. In Sec. III we probe the sensitivity of these TPs to the  $tH$  coupling by using various CP-odd observables. Subsequent sections are dedicated to explore other possibilities of CP-odd observables. In particular, observables based on TPs that incorporate the Higgs momentum are discussed in Sec. IV, whereas observables obtained without using the  $t$  and  $\bar{t}$  momenta are studied in Sec. V. Finally, Sec. VI is devoted to the discussion on the experimental feasibility of the most promising observables encountered here. The main conclusions are summarized in Sec. VII.

## II. THEORETICAL FRAMEWORK FOR $pp \rightarrow t(\rightarrow b\ell^+\nu_\ell) \bar{t}(\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$

Before turning to a discussion of CP-odd observables, it is useful to consider a few theoretical aspects of the process  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$ . In this section we derive a “factorized” expression for the gluon fusion contribution to this process. We then use this expression to isolate various mathematical quantities that will be useful as we construct CP-odd observables.

### A. Factorized expression for the scattering cross section

At the LHC  $t\bar{t}H$  production proceeds via  $q\bar{q}$  annihilation and  $gg$  fusion processes. The relevant leading-order Feynman diagrams are displayed in Fig. 1, where the first two rows show the  $q\bar{q}$  and  $gg$   $s$ -channel diagrams, and the last one depicts the  $gg$   $t$ -channel diagrams. Three more  $gg$ -initiated diagrams are obtained by exchanging the gluon lines in the third row. We describe the  $tH$  coupling with the effective Lagrangian

$$\mathcal{L}_{t\bar{t}H} = -\frac{m_t}{v}(\kappa_t \bar{t}t + i\tilde{\kappa}_t \bar{t}\gamma_5 t)H, \quad (1)$$

where  $v = 246$  GeV is the SM Higgs vacuum expectation value, and the coefficients  $\kappa_t$  and  $\tilde{\kappa}_t$  parameterize the scalar and pseudoscalar interaction, respectively. The SM case is obtained for  $\kappa_t = 1$  and  $\tilde{\kappa}_t = 0$ , while the values  $\kappa_t = 0$  and  $\tilde{\kappa}_t \neq 0$  parameterize a CP-odd Higgs **KK**: “boson”? I don’t think PRD allows us to just say “Higgs”.




FIG. 1: Tree-level Feynman diagrams contributing to  $t\bar{t}H$  production at the LHC. Three more diagrams are obtained by exchanging the gluon lines in the  $t$ -channel diagrams.

In this section we focus on the gluon fusion contributions to  $t\bar{t}H$  production, since these dominate over the the quark-antiquark annihilation contributions. Furthermore, we consider the case in which the top and antitop both decay semileptonically. As we shall show below, assuming the narrow width approximation for the top and antitop quarks, the unpolarized differential cross section for  $gg \rightarrow t(\rightarrow b\ell^+\nu_\ell)\bar{t}(\rightarrow \bar{b}\ell^-\bar{\nu}_\ell)H$  may be written in the following “factorized”

form,<sup>1</sup>

$$d\sigma = \sum_{\substack{b\ell^+\nu_\ell \\ \text{spins}}} \sum_{\substack{\bar{b}\ell^-\bar{\nu}_\ell \\ \text{spins}}} \left( \frac{2}{\Gamma_t} \right)^2 d\sigma(gg \rightarrow t(n_t)\bar{t}(n_{\bar{t}})H) d\Gamma(t \rightarrow b\ell^+\nu_\ell) d\Gamma(\bar{t} \rightarrow \bar{b}\ell^-\bar{\nu}_\ell), \quad (2)$$

where  $d\sigma(gg \rightarrow t(n_t)\bar{t}(n_{\bar{t}})H)$  is the differential cross section for the production of a top and antitop quark, with spin vectors  $n_t$  and  $n_{\bar{t}}$ , respectively, along with a Higgs boson. Also,  $d\Gamma(t \rightarrow b\ell^+\nu_\ell)$  and  $d\Gamma(\bar{t} \rightarrow \bar{b}\ell^-\bar{\nu}_\ell)$  are the partial differential decay widths for an unpolarized top and anti-top quark. The four-vectors  $n_t$  and  $n_{\bar{t}}$  are not arbitrary, but are given by particular combinations of the momenta of the  $t, \bar{t}, \ell^+$  and  $\ell^-$  [? ],

$$n_t = -\frac{p_t}{m_t} + \frac{m_t}{(p_t \cdot p_{\ell^+})} p_{\ell^+} \quad (3)$$

$$n_{\bar{t}} = \frac{p_{\bar{t}}}{m_t} - \frac{m_t}{(p_{\bar{t}} \cdot p_{\ell^-})} p_{\ell^-}. \quad (4)$$

Expressions similar to Eq. (2) have been derived previously for the production of short-lived particles in  $e^-e^+$  colliders [? ] and for  $t\bar{t}$  production both in  $e^-e^+$  colliders [? ] and  $pp$  colliders [? ? ? ? ].




FIG. 2: Schematic representation of the process  $g_a g_b \rightarrow t(\rightarrow b_i \ell^+ \nu_\ell) \bar{t}(\rightarrow \bar{b}_j \ell^- \bar{\nu}_\ell) H$ . The indices  $i, j$  denote the colour of the quarks while  $a, b$  are gluon indices.

To derive the above expressions, we begin by considering the schematic representation for the process  $g_a g_b \rightarrow t(\rightarrow b_i \ell^+ \nu_\ell) \bar{t}(\rightarrow \bar{b}_j \ell^- \bar{\nu}_\ell) H$  that is sketched in Fig. 2. Here  $a$  and  $b$  denote the initial-state gluons and  $i$  and  $j$  refer to the colours of the top and antitop quarks. The amplitude for this process may be written in the following compact form

$$\mathcal{M}^{ab,ij} = \bar{\psi}_t \mathcal{A}^{ab,ij} \psi_{\bar{t}}, \quad (5)$$

<sup>1</sup> The reader is referred to the discussion following Eq. (17) for some qualifying remarks regarding the “factorization” of this expression.

where the spinors  $\bar{\psi}_t$  and  $\psi_{\bar{t}}$  contain all of the information regarding the decay of the virtual top and anti-top, respectively, and where the quantity  $A^{ab,ij}$  is given by

$$\mathcal{A}^{ab,ij} \equiv A_{\mu\nu}^{ab,ij}(\epsilon_{\lambda_a})^\mu(\epsilon_{\lambda_b})^\nu = \sum_{k=1}^8 \mathcal{A}_k^{ab,ij} = \kappa_t \sum_{k=1}^8 \mathcal{S}_k^{ab,ij} + i\tilde{\kappa}_t \sum_{k=1}^8 \mathcal{P}_k^{ab,ij}. \quad (6)$$

The sum over  $k$  in the above expression corresponds to the eight gluon-initiated diagrams indicated in Fig. 1 (**KK: correct?**); also,  $\epsilon_{\lambda_a}$  and  $\epsilon_{\lambda_b}$  are the polarization vectors corresponding to  $g_a$  and  $g_b$ , respectively. In the last equality in Eq. (6) we have explicitly separated the amplitude into two sums, with one sum corresponding to the scalar contributions and the other to the pseudoscalar ones. Taking all of the final-state particles to be massless, we can use the spinor techniques developed in [?] to write  $\bar{\psi}_t$  and  $\psi_{\bar{t}}$  as follows<sup>2</sup>

$$\bar{\psi}_t = -g^2 \mathbb{P}_t(t) \mathbb{P}_W(t-b) \langle b - |\nu_\ell + \rangle \langle \ell^+ + |(\not{t} + m_t) \quad (7)$$

$$\psi_{\bar{t}} = -g^2 \mathbb{P}_t(\bar{t}) \mathbb{P}_W(\bar{t}-\bar{b}) \langle \bar{\nu}_\ell + | \bar{b} - \rangle (\not{\bar{t}} - m_t) | \ell^- + \rangle, \quad (8)$$

where  $|i+(-)\rangle \equiv (1/2)(1 \pm \gamma^5) \psi_i$  represents a right-handed (left-handed) chiral spinor for final-state particle  $i$  and  $\langle i+(-)|$  represents the corresponding adjoint spinor. (**KK: Do we mean the adjoint spinor, related by  $C\psi^T$ , or whatever?**) Also,  $\mathbb{P}_t(q) = (q^2 - m_t^2 + im_t\Gamma_t)^{-1}$  and  $\mathbb{P}_W(q) = (q^2 - M_W^2 + im_W\Gamma_W)^{-1}$ , and we have denoted the momenta of the various particles by the symbols that refer to the names of those particles [?].

Using the expressions defined above for  $\bar{\psi}_t$  and  $\psi_{\bar{t}}$ , we can write the amplitude  $\mathcal{M}^{ab,ij}$  in a form that is (in a sense) factorized. As a first step, we insert Eqs. (7) and (8) into Eq. (5), yielding

$$\mathcal{M}^{ab,ij} = -g^4 \mathbb{P}_t(t) \mathbb{P}_t(\bar{t}) \mathbb{P}_W(t-b) \mathbb{P}_W(\bar{t}-\bar{b}) \langle b - |\nu_\ell + \rangle \langle \bar{\nu}_\ell + | \bar{b} - \rangle \sqrt{2(t \cdot \ell^+)} \sqrt{2(\bar{t} \cdot \ell^-)} [\bar{\phi}_t \mathcal{A}^{ab,ij} \phi_{\bar{t}}], \quad (9)$$

where the spinors  $\phi_t$  and  $\phi_{\bar{t}}$  are defined as

$$\phi_t = \frac{(\not{t} + m_t)}{\sqrt{2(t \cdot \ell^+)}} |\ell^+ + \rangle \quad (10)$$

$$\phi_{\bar{t}} = \frac{(\not{\bar{t}} - m_t)}{\sqrt{2(\bar{t} \cdot \ell^-)}} |\ell^- + \rangle. \quad (11)$$

Note that in writing down the above expressions we have adopted the narrow-width approximation for the top and antitop quarks and for the  $W^\pm$  gauge bosons.<sup>3</sup> (**KK: Do I understand correctly that  $t$  is an on-shell four-vector in  $t \cdot \ell^+$ , but not in  $\mathbb{P}_t(t)$ ? This is similar to saying  $f(x)\delta(x-x_0) = f(x_0)\delta(x-x_0)\dots$ ?**) Working out the projection operators  $\phi_t \bar{\phi}_t$  and  $\phi_{\bar{t}} \bar{\phi}_{\bar{t}}$ , we have

$$\phi_t \bar{\phi}_t = \frac{1}{2}(1 + \not{t}\gamma^5)(\not{t} + m_t) \quad (12)$$

<sup>2</sup> These spinor techniques can also be used for massive final-state particles. Given the energy scale involved in the process in question, however, the assumption of massless final-state particles is sensible and greatly simplifies the derivation of Eq. (2).

<sup>3</sup> Since Eq. (9) contains the top quark propagator term  $\mathbb{P}_t(t)$ , for example,  $|\mathcal{M}^{ab,ij}|^2$  contains the factor  $((t^2 - m_t^2)^2 + m_t^2\Gamma_t^2)^{-1}$ , which is replaced by  $(\pi/m_t\Gamma_t)\delta(t^2 - m_t^2)$  in the narrow-width approximation. Thus we take the four-vector  $t$  appearing in the definition of  $\phi_t$  to be on shell, satisfying  $t^2 = m_t^2$ .

and

$$\phi_{\bar{t}} \bar{\phi}_{\bar{t}} = \frac{1}{2}(1 + \not{n}_{\bar{t}} \gamma^5)(\not{\bar{\ell}} - m_t), \quad (13)$$

with  $n_t$  and  $n_{\bar{t}}$  being the four-vectors defined in Eqs. (3) and (4). Thus,  $\phi_t$  and  $\phi_{\bar{t}}$  may be regarded as describing a top quark with spin vector  $n_t$  and an antitop quark with spin vector  $n_{\bar{t}}$ , respectively.

As a final step toward factorizing the amplitude  $\mathcal{M}^{ab,ij}$ , we note that the amplitude for a top quark with spin vector  $n_t$  to decay into  $b\ell^+\nu_\ell$  is given by

$$\mathcal{M}(t(n_t) \rightarrow b\ell^+\nu_\ell) = ig^2 \mathbb{P}_W(t-b) \langle b - |\nu_\ell + \rangle \sqrt{2(t \cdot \ell^+)} , \quad (14)$$

and likewise,

$$\mathcal{M}(\bar{t}(n_{\bar{t}}) \rightarrow \bar{b}\ell^-\bar{\nu}_\ell) = ig^2 \mathbb{P}_W(\bar{t}-\bar{b}) \langle \bar{\nu}_\ell + |\bar{b}- \rangle \sqrt{2(\bar{t} \cdot \ell^-)} . \quad (15)$$

Furthermore, the term inside the square brackets in Eq. (9) is the amplitude for producing a top quark with spin vector  $n_t$ , along with an anti-top with spin vector  $n_{\bar{t}}$  and a Higgs boson,

$$\mathcal{M}(g_a g_b \rightarrow t^i(n_t) \bar{t}^j(n_{\bar{t}}) H) = \bar{\phi}_t \mathcal{A}^{ab,ij} \phi_{\bar{t}}. \quad (16)$$

Combining Eqs. (14)-(16), we can write Eq. (9) in a form that appears to be factorized,

$$\mathcal{M}^{ab,ij} = \mathbb{P}_t(t) \mathbb{P}_{\bar{t}}(\bar{t}) \mathcal{M}(t(n_t) \rightarrow b\ell^+\nu_\ell) \mathcal{M}(\bar{t}(n_{\bar{t}}) \rightarrow \bar{b}\ell^-\bar{\nu}_\ell) \mathcal{M}(g_a g_b \rightarrow t^i(n_t) \bar{t}^j(n_{\bar{t}}) H). \quad (17)$$

It is important to note that, even though the above expression has the appearance of being factorized into production and decay parts, this apparent factorization is a bit misleading. In particular, the amplitude for  $t\bar{t}H$  production contains the top and antitop quark spin four-vectors  $n_t$  and  $n_{\bar{t}}$ , which depend on final-state kinematical quantities [see Eqs. (3) and (4)]. With this qualification in mind, we may now use the amplitude in Eq. (17) to determine the corresponding scattering cross section. After some manipulation of the phase space variables to take advantage of the presence of the propagator terms,  $\mathbb{P}_t(t)$  and  $\mathbb{P}_{\bar{t}}(\bar{t})$ , we arrive at the expression in Eq. (2). This expression also has the appearance of being factorized, but qualifying remarks, similar to those above, apply. **(KK: I'm still a bit confused as to why  $n_t$  appears in the amplitude expression for the decay, but not in the expression for the cross section. That is, the top spins are summed over in the width that appears in the factorized expression for the cross section. That part makes sense to me, but I'm not so clear on how to get there from Eq. (17). Probably need to look more closely at the notes!)**

~~Let us focus now in~~

## B. Triple products

(KK: maybe we should change this title for the subsection....)

The expression derived above for the scattering cross section [see Eq. (2), as well as Eq. (17)] provides significant insight into how one might analyze  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$  in order to determine the nature of the top-Higgs coupling. In particular, let us focus on the production amplitude,  $\mathcal{M}(g_a g_b \rightarrow t^i(n_t) \bar{t}^j(n_{\bar{t}}) H)$ , which forms part of the overall amplitude in Eq. (17) ~~needed for computing the production~~. The absolute value squared

of the production amplitude is used to determine  $d\sigma(gg \rightarrow t(n_t)\bar{t}(n_{\bar{t}})H)$ , which in turn forms part of the ~~unpolarized differential cross-section~~. By ~~summing~~ expression for the “factorized” cross section in Eq. (2). ~~Summing~~ over colour and gluon indices we have

$$\sum_{\substack{a,b \\ i,j}} |\mathcal{M}(g_a g_b \rightarrow t^i(n_t)\bar{t}^j(n_{\bar{t}})H)|^2 = \sum_{\substack{a,b \\ i,j}} \left| \sum_{k=1}^8 C_k^{ab,ij} \bar{\phi}_t(\kappa_t \mathcal{S}_k + i\tilde{\kappa}_t \mathcal{P}_k) \phi_{\bar{t}} \right|^2, \quad (18)$$

where we have separated the colour structure of each diagram by ~~using the definitions defining~~ (KK: do these need  $k$  subscripts?)  $\mathcal{S}^{ab,ij} = C_k^{ab,ij} \mathcal{S}$  and  $\mathcal{P}^{ab,ij} = C_k^{ab,ij} \mathcal{P}$  [see Eqs. (6) and (16)]. Also, the factors  $g_s^2 m_t/v$  and  $-ig_s^2 m_t/v$  arising from the vertices of the  ~~$t$ - and  $s$ -channel  $t$ - and  $s$ -channel~~ diagrams respectively have been included in the definition of  $C_k^{ab,ij}$  for convenience. The terms linear in  $\kappa_t$  and  $\tilde{\kappa}_t$  can be written as

$$\mathcal{O}(\kappa_t \tilde{\kappa}_t) \rightarrow \frac{1}{2} \kappa_t \tilde{\kappa}_t \sum_{k,r} \mathbb{C}_{kr} \text{Im} \left\{ \text{Tr} \left[ (1 + \not{n}_t \gamma^5) (\not{t} + m_t) \mathcal{S}_k (1 + \not{n}_{\bar{t}} \gamma^5) (\not{\bar{t}} - m_{\bar{t}}) \tilde{\mathcal{P}}_r \right] \right\}, \quad (19)$$

where the factor  $\mathbb{C}_{kr} = \sum_{ab,ij} C_k^{ab,ij} C_r^{ab,ij*}$  is real and ~~where~~  $\tilde{\mathcal{P}}_r = \gamma^0 \mathcal{P}_r \gamma^0$  (KK: wouldn't this be  $\mathcal{P}_r^\dagger$ ? I didn't look for this in the notes yet, so maybe I'm just missing something). The only ~~contributing terms~~ terms that yield non-zero contributions in the above sum are those with an odd number of  $\gamma^5$  matrices ~~and will lead to triple products~~; these lead to triple-product (TP) structures ~~, i.e. contractions between the levi-civita tensor  $\epsilon_{\alpha\beta\gamma\delta}$  with four momenta of the form  $\epsilon_{\alpha\beta\gamma\delta} p_a^\alpha p_b^\beta p_c^\gamma p_d^\delta$  where  $p_a$ - $p_d$  represent various four momenta associated with the process.~~ In contrast, it can be seen from Eq. (18) that the terms proportional to  $\kappa_t^2$  and  $\tilde{\kappa}_t^2$  ~~contain~~ descend from traces containing an even number of  $\gamma^5$  matrices and can be written in terms of scalar products of the available momenta.

~~In order to state this more clearly, we will consider the form of the differential cross-section~~ With the above considerations in mind, it is useful to write a general expression for the differential cross section  $d\sigma(gg \rightarrow t(n_t)\bar{t}(n_{\bar{t}})H)$  in terms of the momenta  $q = (q_1 - q_2)/2$ ,  $Q = (q_1 + q_2)/2$ ,  $t$ ,  $\bar{t}$ ,  $n_t$  and  $n_{\bar{t}}$ , where  $q_{1,2}$  denote the momenta of the ~~initial-state~~ gluons. Note that with this choice,  $q \cdot Q = 0$  and  $Q^2 = -q^2 = M_{t\bar{t}H}^2/4$ , where  $M_{t\bar{t}H}$  is the invariant mass of the ~~system  $t\bar{t}H$~~ . From these six vectors, ~~fifteen system~~ Fifteen TPs can be built<sup>4</sup>, ~~and the general form of the production differential cross-section will be given by constructed from these six four-vectors~~,<sup>4</sup> so that

$$d\sigma(gg \rightarrow t(n_t)\bar{t}(n_{\bar{t}})H) = \kappa_t^2 f_1(p_i \cdot p_j) + \tilde{\kappa}_t^2 f_2(p_i \cdot p_j) + \kappa_t \tilde{\kappa}_t \sum_{l=1}^{15} g_l(p_i \cdot p_j) \epsilon_l, \quad (20)$$

where  $\epsilon_l = \epsilon_{\alpha\beta\gamma\delta} p_a^\alpha p_b^\beta p_c^\gamma p_d^\delta$  denotes the  $l$ th TP ~~, the Roman indices (we adopt the convention  $\epsilon_{0123} = +1$ ) and where  $p_i$  and  $p_j$  refer to any of the involved momenta and we adopt the convention  $\epsilon_{0123} = +1$  six momenta~~. The functions  $f_{1,2}$  and  $g_k$  depend only on the possible scalar products and are therefore even under a parity transformation (P). However, the terms linear in  $\kappa_t \tilde{\kappa}_t$  are P-odd due to the presence of the P-odd TPs. Hence, only the functions  $f_{1,2}$  will contribute to the total cross-section, whereas the ~~TPs~~ TP terms will be sensitive to the

<sup>4</sup> We note that these fifteen TPs are not linearly independent (see the epsilon relations discussed in [?]).

<sup>4</sup> We note that these fifteen TPs are not linearly independent (see the epsilon relations discussed in Ref. [?]).



sign of the anomalous coupling  $\tilde{\kappa}_t$ . ~~From Of~~ the fifteen TPs mentioned above, we will focus on those that ~~do not include  $q$  and contain both~~ contain both of the spin vectors  $n_t$  and  $n_{\bar{t}}$ . ~~The former is motivated in~~ but do not include  $q$ . The decision not to consider  $q$ -dependent TPs is motivated by the fact that  $q$  cannot be ~~eliminated expressed~~ in terms of the momenta of final state particles (like as  $Q$  in virtue of momentum-energy can, by virtue of energy-momentum conservation). The ~~latter takes into account that for  $t\bar{t}$  production at hadron colliders the decision to focus on TPs that contain both  $n_t$  and  $n_{\bar{t}}$  is rooted in the fact that the spins of pair-produced top and antitop are not polarized but their spins are strongly correlated and observables combining the quarks are highly correlated at hadron colliders (even though the quarks themselves are unpolarized).~~ (KK: did I reword this correctly?) Observables that combine the decay products of the  $t$  and  $\bar{t}$  will be sensitive to this spin correlation [? ]. A similar behaviour is expected in  $t\bar{t}H$  production, where it can be shown that ~~the~~ single-spin asymmetries vanish [? ? ]. Hence, in order to construct observables sensitive to the structure of the  $tH$  coupling, we will ~~prefer restrict our attention to~~ those TPs that include information on the decay products of both the top and anti-top quarks. Only five of the fifteen TPs in Eq. (20) do not involve the ~~four~~ vector  $q$  and, among these, ~~we will focus on the three that only three~~ include both  $n_t$  and  $n_{\bar{t}}$ . ~~Thus, we will restrict our attention to the following TPs~~

$$\epsilon_1 \equiv \epsilon(t, \bar{t}, n_t, n_{\bar{t}}), \quad (21)$$

$$\epsilon_2 \equiv \epsilon(Q, \bar{t}, n_t, n_{\bar{t}}), \quad (22)$$

$$\epsilon_3 \equiv \epsilon(Q, t, n_t, n_{\bar{t}}). \quad (23)$$

~~Finally Before~~ turning to a consideration of various CP-odd observables, we remark that even though all of the above discussion ~~was given took place~~ within the context of  $gg$ -initiated production, similar conclusions are obtained for  $q\bar{q}$ -initiated production. In particular, the definitions of the spin vectors in Eqs. (3)-(4) and the general form of  $d\sigma$  introduced in Eq. (20) are valid in both cases.

### III. CP-ODD OBSERVABLES

We present below three types of observables based on the TPs discussed in Sec. II, mean values, asymmetries and angular distributions. These observables are sensitive not only to the magnitude of the pseudoscalar coupling  $\tilde{\kappa}_t$  but also to its sign. With the aim of testing them, we have used in all the cases a set of  $10^5$  events  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$  simulated at parton level with MadGraph5\_aMC@NLO [? ] at a center-of-mass energy of 14 TeV for different values of the couplings  $\kappa_t$  and  $\tilde{\kappa}_t$ <sup>5</sup>. We have used this somewhat large number of events with the aim of determining clearly the extent to which the proposed observables are sensitive to the NP coupling. However, we will discuss the corresponding experimental viability in Sec. VI. Finally, we have also imposed the following set of cuts:  $p_T$  of leptons  $> 10$  GeV,  $|\eta|$  of leptons  $< 2.5$ ,  $|\eta|$  of b jets  $< 2.5$  and  $\Delta R_{\ell\ell} > 0.4$ .

Under the assumption of no other new physics sources but the inclusion of the pseudoscalar coupling  $\tilde{\kappa}_t$ , indirect constraints based on the Higgs production from gluon fusion followed by its

<sup>5</sup> Although we use the same number of events in all the cases, the required luminosities are different since the cross section depends on the value of  $\tilde{\kappa}_t$ .

decay to two photons disfavour  $\kappa_t < 0$  without resolving the degeneracy in the sign of  $\tilde{\kappa}_t$  [? ]. On the other hand, by assuming the tensor structure of the Lagrangian to be the same as in the SM and considering a parameterization with one universal Higgs coupling to vector bosons,  $\kappa_V$ , and one universal Higgs coupling to fermions,  $\kappa_f$ , the measured signal strengths provided by ATLAS and CMS collaborations are compatible with the values predicted by the SM, ( $\kappa_f = 1, \kappa_V = 1$ ). Based on these facts, we will set the value of the scalar coupling to its SM value,  $\kappa_t = 1$ , and concentrate in modifications of the value as well as the sign of the pseudoscalar coupling. In particular, we have analyzed the cases  $\tilde{\kappa}_t = 0, \pm 0.25, \pm 0.5, \pm 0.75, \pm 1$ . However, we will give some results and comments regarding the pure CP-odd hypothesis ( $\kappa_t = 0, \tilde{\kappa}_t = 1$ ).

### A. Asymmetry

The first type of CP-odd observable we will consider is the asymmetry. For a certain TP, we define an asymmetry which gives the balance between the number of events with  $\epsilon > 0$  and those for which  $\epsilon < 0$ :

$$\mathcal{A}(\epsilon) = \frac{N(\epsilon > 0) - N(\epsilon < 0)}{N(\epsilon > 0) + N(\epsilon < 0)}. \quad (24)$$

Based on the general expression given in Eq. (20), we expect the following shape for the asymmetry,

$$\mathcal{A}(\epsilon) = \frac{A\kappa_t\tilde{\kappa}_t}{B\kappa_t^2 + C\tilde{\kappa}_t^2}, \quad (25)$$

which for  $\kappa_t = 1$  can be parameterized as

$$\mathcal{A}(\epsilon) = \frac{a\tilde{\kappa}_t}{1 + b\tilde{\kappa}_t^2}, \quad (26)$$

where the parameter  $a \equiv A/B$  determines the sensitivity to the pseudoscalar coupling, whereas  $b \equiv C/B$  quantifies the deviation from the linear behaviour. The results for  $\kappa_t = 1$  and  $\tilde{\kappa}_t = 0, \pm 1$  are shown in Table I, where the values in terms of the corresponding statistical uncertainty are also included. It can be seen that the considered asymmetries are useful to separate the CP-mixed cases from the SM case and also to discriminate the different sign of  $\tilde{\kappa}_t$ , giving deviations of order  $10\sigma$  in the former case and  $20\sigma$  in the latter.

The sensitivity of the asymmetry is quite similar for the three TPs, as can be seen by including other values of  $\tilde{\kappa}_t$  and using the expression in Eq. (26) as fitting function (see Fig. 3). In fact, we obtain the fit parameters ( $a = -0.057 \pm 0.006, b = 0.5 \pm 0.2$ ), ( $a = -0.056 \pm 0.006, b = 0.5 \pm 0.2$ ) and ( $a = 0.058 \pm 0.006, b = 0.6 \pm 0.2$ ) for  $\epsilon_1, \epsilon_2$  and  $\epsilon_3$ , respectively. In addition, we include in Fig. 4 the asymmetry for  $\epsilon_1$  isolating the  $gg$ -initiated,  $q\bar{q}$ -initiated and  $pp$  production. We see that the asymmetry is enhanced for  $gg$ -initiated production, while it is reduced and of opposite sign for the  $q\bar{q}$ -initiated case. The asymmetry for the  $pp$  case is then dominated by the  $gg$  contribution, albeit smaller due to the  $q\bar{q}$  contribution.

TABLE I: Asymmetries obtained by using  $10^5$  simulated events for the TPs  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$ ,  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  and  $\epsilon_3 = \epsilon(Q, t, n_t, n_{\bar{t}})$  in the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ .

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_1)$	$\mathcal{A}(\epsilon_1)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_2)$	$\mathcal{A}(\epsilon_2)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_3)$	$\mathcal{A}(\epsilon_3)/\sigma_{\mathcal{A}}$
1	-1	0.0315	10.0	0.0332	10.5	-0.0307	-9.7
1	0	-0.0021	-0.7	0.0009	0.3	-0.0011	-0.3
1	1	-0.0379	-12.0	-0.0411	-13.0	0.0378	12.0




FIG. 3: Asymmetries for the TPs  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$  (top-left),  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  (top-right) and  $\epsilon_3 = \epsilon(Q, t, n_t, n_{\bar{t}})$  (bottom). The points represent the values for  $\tilde{\kappa}_t = 0, \pm 0.25, \pm 0.5, \pm 0.75, \pm 1$  and the red solid line is the fitting curve.



FIG. 4: Asymmetry for the TP  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$ . The dashed line (red) corresponds to  $gg$ -initiated production, the dot-dashed line (grey) to  $q\bar{q}$ -initiated production and the solid line (blue) to  $pp$  production.

On the other hand, we have also tested various linear combinations of the TPs  $\epsilon_{1,2,3}$ , with the result that the asymmetry is enhanced in the following case:

$$\epsilon_4 = \epsilon_3 - \epsilon_2 = \epsilon(Q, t - \bar{t}, n_t, n_{\bar{t}}). \quad (27)$$

Note that in the  $Q$  rest frame  $\epsilon_4 = Q^0(\vec{t} - \vec{\bar{t}}) \cdot (\vec{n}_t \times \vec{n}_{\bar{t}})$  and the sign of this TP is determined by the quantity  $(\vec{t} - \vec{\bar{t}}) \cdot (\vec{n}_t \times \vec{n}_{\bar{t}})$ . The values obtained for the asymmetry of this TP are shown in II. We see by comparing the results in Tables I and II that the capability of the asymmetry to separate the two CP-mixed hypotheses is increased by at least  $3\sigma$ .

TABLE II: Asymmetry for the TP  $\epsilon_4$  for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . The values are obtained by using  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_4)$	$\mathcal{A}(\epsilon_4)/\sigma_{\mathcal{A}}$
1	-1	-0.0371	-12
1	0	0.0004	0.1
1	1	0.0461	14

Finally, it is worth noting that the asymmetry is not useful for discriminating between the SM hypothesis ( $\kappa_t = 1, \tilde{\kappa}_t = 0$ ) and the pure pseudoscalar hypothesis ( $\kappa_t = 0, \tilde{\kappa}_t = 1$ ) because, being linear in both  $\kappa_t$  and  $\tilde{\kappa}_t$ , it is expected to vanish in these cases. However, we will show in the next section that there exist angular distributions derived from the TPs that are actually suitable to distinguish these two hypothesis.

## B. Angular Distributions

Given a certain TP, it is possible to define associated angular distributions that are also sensitive to the pseudoscalar coupling  $\tilde{\kappa}_t$ . In order to clarify this, let us first consider the TP  $\epsilon(t, \bar{t}, n_t, n_{\bar{t}})$ . This TP can be written as  $\epsilon(t + \bar{t}, \bar{t}, n_t, n_{\bar{t}})$  so that in the reference frame defined by  $\vec{t} + \vec{\bar{t}} = 0$  and  $\vec{t} \parallel \hat{z}$  we have

$$\epsilon(t + \bar{t}, \bar{t}, n_t, n_{\bar{t}}) = M_{t\bar{t}} |\vec{t}| (\vec{n}_t \times \vec{n}_{\bar{t}})_z = M_{t\bar{t}} |\vec{t}| |\vec{n}_t| |\vec{n}_{\bar{t}}| \sin \theta_{n_t} \sin \theta_{n_{\bar{t}}} \sin \Delta\phi(n_t, n_{\bar{t}}), \quad (28)$$

where  $M_{t\bar{t}}$  is the invariant mass of the  $t\bar{t}$  pair, the angles  $\theta_{n_t}$  and  $\theta_{n_{\bar{t}}}$  denote the polar angles of  $\vec{n}_t$  and  $\vec{n}_{\bar{t}}$  respectively and finally  $\Delta\phi(n_t, n_{\bar{t}})$  is the angular difference between the projections of  $\vec{n}_t$  and  $\vec{n}_{\bar{t}}$  onto the plane perpendicular to  $\vec{t}$ . If we define this angle within the range  $[-\pi, \pi]$ , we see from Eq. (28) that its sign will determine the sign of the TP. In this sense, the distribution of the number of the events with respect to the angle  $\Delta\phi(n_t, n_{\bar{t}})$  is related to the asymmetry of the TP,

$$\mathcal{A}(\epsilon) = 1 - 2 \frac{N(\epsilon < 0)}{N_T} \quad \text{and} \quad \frac{N(\epsilon < 0)}{N_T} = \int_{-\pi}^0 \frac{1}{N_T} \frac{dN}{d\Delta\phi(n_t, n_{\bar{t}})} d\Delta\phi(n_t, n_{\bar{t}}), \quad (29)$$

where  $N_T$  is the total number of events. Moreover, from a certain TP one can derive different angular distributions by considering different reference frames, but all of these will satisfy Eq. (29). In particular, we will analyze the following angular distributions based on the TPs defined in Sec. II:

- Related to  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$  we consider  $d\sigma/d\Delta\phi_1(n_t, n_{\bar{t}})$  in the rest frame of  $t\bar{t}$  with  $\vec{t}$  in the  $z$ -axis. The angle  $\Delta\phi_1(n_t, n_{\bar{t}})$  is the angular difference between the projection of the spin vectors in the plane perpendicular to  $\vec{t}$ .
- Related to  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  we consider  $d\sigma/d\Delta\phi_2(n_t, n_{\bar{t}})$  in the rest frame of  $Q$  with  $\vec{t}$  in the  $z$ -axis. The angle  $\Delta\phi_2(n_t, n_{\bar{t}})$  is the angular difference between the projection of the spin vectors in the plane perpendicular to  $\vec{t}$ .
- Related to  $\epsilon_3 = \epsilon(Q, t, n_t, n_{\bar{t}})$  we consider  $d\sigma/d\Delta\phi_3(n_t, n_{\bar{t}})$  in the rest frame of  $Q$  with  $\vec{t}$  in the  $z$ -axis. The angle  $\Delta\phi_3(n_t, n_{\bar{t}})$  is the angular difference between the projection of the spin vectors in the plane perpendicular to  $\vec{t}$ .

In Fig. 5 we present the normalized distributions obtained for the first case listed above with  $\kappa_t = 1, \tilde{\kappa}_t = 0$  (CP-even Higgs boson),  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$  (CP-mixed Higgs boson) and  $\kappa_t = 0, \tilde{\kappa}_t = 1$  (CP-odd Higgs boson). In Fig. 6 we display in turn the analogous distributions for  $\epsilon_2$ . The distributions corresponding to  $\epsilon_3$  are similar to those of  $\epsilon_2$  but with the opposite shift for  $\kappa_t = \tilde{\kappa}_t = 1$  and  $\kappa_t = -\tilde{\kappa}_t = 1$  and then we do not include them here.




FIG. 5: Angular distributions associated with the TP  $\epsilon(t, \bar{t}, n_t, n_{\bar{t}})$  for various values of  $(\kappa_t, \tilde{\kappa}_t)$ . The error bars correspond to the statistical uncertainties.




FIG. 6: Angular distributions associated with the TP  $\epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  for various values of  $(\kappa_t, \tilde{\kappa}_t)$ . The error bars correspond to the statistical uncertainties.

As can be seen from Figs. 5 and 6, the peak of the distributions is shifted according to the values of the couplings  $\kappa_t, \tilde{\kappa}_t$ . This shift appears to be approximately the same but with opposite sign for  $\kappa_t = \tilde{\kappa}_t = 1$  and  $\kappa_t = -\tilde{\kappa}_t = 1$ , distinguishing then the sign of the pseudoscalar coupling. The last is the expected behaviour since the asymmetry is linear in  $\tilde{\kappa}_t$  (see Sec. III A) and the quantity  $N(\epsilon < 0)/N_T$  is related to the angular distribution according to Eq. (29). The distributions for the SM and the pure pseudoscalar hypotheses are clearly different. While in the former case the distributions exhibit a minimum at  $\Delta\phi_{1,2}(n_t, n_{\bar{t}}) = 0$ , in the latter the similar distributions display a peak, so that the two hypotheses can be discriminated, in contrast to the asymmetry that vanishes for both hypotheses. In order to quantify the mentioned shifts, we have fitted the simulated distributions with the following function proposed in [? ],

$$\frac{1}{\sigma} \frac{d\sigma}{d\Delta\phi_i(n_t, n_{\bar{t}})} = a_0 + a_1 \cos(\Delta\phi_i(n_t, n_{\bar{t}}) + \delta), \quad i = 1, 2, 3. \quad (30)$$

To the extent that this equation is exact we note that Eq. (32) gives  $\mathcal{A}(\epsilon_i) = 4a_1 \sin \delta$ . With this fitting function, we obtain phase shifts approximately between 0.9 and 1 (−1 and −0.9)

for  $\kappa_t = \tilde{\kappa}_t = 1$  ( $\kappa_t = -\tilde{\kappa}_t = 1$ ) both for  $\epsilon_1$  and  $\epsilon_2$ . However, the quality of the fits is not good enough, being worse in the case of the  $\epsilon_1$  distributions for which  $\chi^2/\text{d.o.f}$  is in the range 1.69-3.86, in contrast to the range 0.53-1.16 obtained for  $\epsilon_2$ . The main impact on the deviation from the functional form proposed in Eq. (30) appears to be caused by the  $\Delta R_{\ell\ell}$  cut we have imposed. In fact, when this cut is turned off the above ranges for  $\chi^2/\text{d.o.f}$  modify to 0.75-1.14 and 0.44-1.07 for the  $\epsilon_1$  and  $\epsilon_2$  distributions respectively. In Tables III and IV we list the results of the fits obtained when the  $\Delta R_{\ell\ell}$  cut is relaxed. The results for the TP  $\epsilon_3$  are pretty similar to those for  $\epsilon_2$  but with opposite sign in the phase shifts of the CP-mixed hypotheses and hence we do not include them here. In addition, we show in Fig. 7 the angular distributions along with the fit curves for both CP-mixed cases when no cut is imposed in  $\Delta R_{\ell\ell}$ .

TABLE III: Fit results for the angular distribution  $d\sigma/(\sigma d\Delta\phi_1(n_t, n_{\bar{t}}))$  related to the TP  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$  when the  $\Delta R_{\ell\ell}$  is turned off. Note that the parameter  $a_1$  change its sign for  $\kappa_t = 0, \kappa_t = 1$ .

$\kappa_t$	$\tilde{\kappa}_t$	$a_0$	$a_1$	$\delta$
1	-1	$0.1592 \pm 0.0006$	$-0.0139 \pm 0.0008$	$0.81 \pm 0.07$
1	0	$0.1595 \pm 0.0006$	$-0.0181 \pm 0.0008$	$0.002 \pm 0.06$
1	1	$0.1591 \pm 0.0006$	$-0.0131 \pm 0.0008$	$-0.82 \pm 0.07$
0	1	$0.1591 \pm 0.0006$	$0.0102 \pm 0.0008$	$0.11 \pm 0.08$

TABLE IV: Fit results for the angular distribution  $d\sigma/(\sigma d\Delta\phi_2(n_t, n_{\bar{t}}))$  related to the TP  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  when the  $\Delta R_{\ell\ell}$  is turned off. Note that the parameter  $a_1$  change its sign for  $\kappa_t = 0, \kappa_t = 1$ .

$\kappa_t$	$\tilde{\kappa}_t$	$a_0$	$a_1$	$\delta$
1	-1	$0.1591 \pm 0.0006$	$-0.0146 \pm 0.0008$	$0.73 \pm 0.06$
1	0	$0.1594 \pm 0.0007$	$-0.0190 \pm 0.0008$	$0.005 \pm 0.06$
1	1	$0.1592 \pm 0.0006$	$-0.0136 \pm 0.0008$	$-0.77 \pm 0.07$
0	1	$0.1591 \pm 0.0006$	$0.0113 \pm 0.0008$	$0.09 \pm 0.08$

From Tables III and IV we see that the parameter  $\delta$  is sensitive not only to the modulus of  $\tilde{\kappa}_t$  but also to its sign, as expected if Eq. (29) is taken into account. The phase shift  $\delta$  for the distribution of the angle  $\Delta\phi_1$  appears to exhibit a slightly higher sensitivity than that obtained from the  $\Delta\phi_2$ -distribution, albeit both are still compatible within the statistical uncertainties. However, it is important to stress that the fits of the  $\Delta\phi_2$ -distributions give always smaller values for  $\chi^2/\text{d.o.f}$ .






FIG. 7: Angular distributions  $d\sigma/(\sigma d\Delta\phi_1(n_t, n_{\bar{t}}))$  (top) and  $d\sigma/(\sigma d\Delta\phi_2(n_t, n_{\bar{t}}))$  (bottom) associated with the TPs  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$  and  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  respectively for the CP-mixed cases  $\kappa_t = \tilde{\kappa}_t = 1$  (left) and  $\kappa_t = -\tilde{\kappa}_t = 1$  (right) when the  $\Delta R_{\ell\ell}$  cut is turned off. Also, the corresponding fit curves are displayed in red.

Concerning the angular distributions that can be related to the combination  $\epsilon_4$ , we have analyzed the  $\Delta\phi(n_t, n_{\bar{t}})$  distribution in the  $Q$  rest frame with  $H$  in the  $z$ -axis for various values of  $\kappa_t$  and  $\tilde{\kappa}_t$ . We have found that these distributions are not described by Eq. (30) and their range of variation is larger than that of the distributions displayed in Figs. 5 and 6. Unlike the distributions related to  $\epsilon_1$ - $\epsilon_3$ , those arising from  $\epsilon_4$  exhibit small changes in their shapes for all the considered hypotheses and for this reason we have not included the corresponding plots here. However, the larger range of variation of the  $\epsilon_4$  distributions leads to higher values for the asymmetry (as can be seen from Tables I and II) even when the changes in the respective shapes are smaller than in the case of the distributions described by Eq. (30).

### C. Mean value

We turn now to consider the last type of observables constructed from the TPs that are sensitive to  $\tilde{\kappa}_t$ , the mean value. Given a certain TP, we define its mean value in the following manner,

$$\langle \epsilon \rangle = \frac{\int \epsilon \{d\sigma(pp \rightarrow b \ell^+ \nu_\ell \bar{b} \ell^- \bar{\nu}_\ell H)/d\Phi\} d\Phi}{\int \{d\sigma(pp \rightarrow b \ell^+ \nu_\ell \bar{b} \ell^- \bar{\nu}_\ell H)/d\Phi\} d\Phi}, \quad (31)$$

where  $\Phi$  is the Lorentz invariant phase space corresponding to the final state  $b \ell^+ \nu_\ell \bar{b} \ell^- \bar{\nu}_\ell H$ . From Eq. (20) we see that only the terms linear in  $\kappa_t$  and  $\tilde{\kappa}_t$  will contribute to the mean value, so that we expect this observable to be sensitive not only to the value but also to the relative sign of the couplings.

The results obtained for the TPs  $\epsilon_1 = \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$ ,  $\epsilon_2 = \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  and  $\epsilon_3 = \epsilon(Q, t, n_t, n_{\bar{t}})$  introduced in Sec. II are displayed in Table V, where we list the deviation of the mean values from the SM case ( $\tilde{\kappa}_t = 0$ ) in terms of the corresponding statistical uncertainty of the estimator of  $\langle \epsilon \rangle$ ,  $\bar{\epsilon}$ .

TABLE V: Mean values obtained for the TPs  $\epsilon_{1,2,3}$  for the SM case and two CP-mixed cases with opposite sign in the pseudoscalar coupling. The values are obtained by using  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\langle \epsilon_1 \rangle / \sigma_{\bar{\epsilon}_1}$	$\langle \epsilon_2 \rangle / \sigma_{\bar{\epsilon}_2}$	$\langle \epsilon_3 \rangle / \sigma_{\bar{\epsilon}_3}$
1	-1	4.26	4.94	-5.81
1	0	-0.91	-0.22	1.25
1	1	-7.98	-8.83	8.75

We see that the three observables are capable of distinguishing the SM case from both CP-mixed cases. Further, the observables are sensitive to the sign of  $\tilde{\kappa}_t$  and the two CP-mixed cases are also clearly disentangled. The observables  $\langle \epsilon_2 \rangle$  and  $\langle \epsilon_3 \rangle$  appear to be slightly more sensitive than  $\langle \epsilon_1 \rangle$ . On the other hand, the mean value for the combination  $\epsilon_4$  introduced in Sec. III A gives slightly smaller values than those listed in Table V, with -4.32, 1.11 and 7.23 for the cases ( $\kappa_t = 1, \tilde{\kappa}_t = -1, 0, 1$ ) respectively. As with the asymmetry, the hypotheses CP-even and CP-odd cannot be distinguished by the mean value that is also linear in both  $\kappa_t$  and  $\tilde{\kappa}_t$  (see Eqs. (20) and (31)). By taking into account the results given in Sec. III A, we can conclude that the sensitivity to the NP contribution is smaller for the mean values than for the asymmetries of the TPs under consideration.

### IV. CP-ODD OBSERVABLES NOT DEPENDING ON $t$ AND $\bar{t}$ SPIN VECTORS

So far we have considered three TPs involving the momenta  $t, \bar{t}$  and  $Q$  and the spin vectors  $n_t$  and  $n_{\bar{t}}$  given in Eqs. (3)-(4). In fact, we have described the general form of the differential cross-section in terms of these vectors in Eq. (20). Here, we will take into account another possibilities

for the choice of the vectors from which the CP-odd observables can be constructed. From the definitions in Eqs. (3)-(4), we see that the TPs  $\epsilon_{1,2,3}$  can be written as follows,

$$\epsilon(t, \bar{t}, n_t, n_{\bar{t}}) = \frac{m_t^2}{(t \cdot \ell^+)(\bar{t} \cdot \ell^-)} \epsilon(t, \bar{t}, \ell^-, \ell^+), \quad (32)$$

$$\epsilon(Q, \bar{t}, n_t, n_{\bar{t}}) = \frac{m_t^2}{(t \cdot \ell^+)(\bar{t} \cdot \ell^-)} \left( \epsilon(t, \bar{t}, \ell^-, \ell^+) + \epsilon(H, \bar{t}, \ell^-, \ell^+) + \frac{(t \cdot \ell^+)}{m_t^2} \epsilon(H, \bar{t}, t, \ell^-) \right) \quad (33)$$

$$\epsilon(Q, t, n_t, n_{\bar{t}}) = \frac{m_t^2}{(t \cdot \ell^+)(\bar{t} \cdot \ell^-)} \left( -\epsilon(t, \bar{t}, \ell^-, \ell^+) + \epsilon(H, t, \ell^-, \ell^+) + \frac{(\bar{t} \cdot \ell^-)}{m_t^2} \epsilon(H, \bar{t}, t, \ell^+) \right). \quad (34)$$

The above equations express the TPs studied in the last sections as linear combination of TPs involving the momenta  $t, \bar{t}, H, \ell^+$  and  $\ell^-$ , with coefficients that are functions of phase space variables. These five momenta give rise to five TPs whose sensitivity can also be tested by means of the observables introduced in Secs. III A-III C. In the first place, we have found that TPs not including both lepton and anti-lepton momenta present a negligible sensitivity, and then we will concentrate here on the results obtained for the remaining TPs,  $\epsilon_5 \equiv \epsilon(t, \bar{t}, \ell^-, \ell^+)$ ,  $\epsilon_6 \equiv \epsilon(H, t, \ell^-, \ell^+)$  and  $\epsilon_7 \equiv \epsilon(H, \bar{t}, \ell^-, \ell^+)$ . The Tables VI and VII summarize the results for these TPs.

TABLE VI: Mean values obtained for the TPs  $\epsilon_{5,6,7}$  for the SM case and two CP-mixed cases with opposite sign in the pseudoscalar coupling. The values correspond to  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\langle \epsilon_5 \rangle / \sigma_{\bar{\epsilon}_5}$	$\langle \epsilon_6 \rangle / \sigma_{\bar{\epsilon}_6}$	$\langle \epsilon_7 \rangle / \sigma_{\bar{\epsilon}_7}$
1	-1	3.98	-1.96	1.69
1	0	-0.43	1.25	0.74
1	1	-6.76	3.46	-3.29

TABLE VII: Asymmetries for the TPs  $\epsilon_{5,6,7}$  for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . The values correspond to  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_5)$	$\mathcal{A}(\epsilon_5)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_6)$	$\mathcal{A}(\epsilon_6)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_7)$	$\mathcal{A}(\epsilon_7)/\sigma_{\mathcal{A}}$
1	-1	0.0315	10.0	-0.0134	-4.2	0.0111	3.5
1	0	-0.0021	-0.7	-0.0011	-0.3	0.0009	0.3
1	1	-0.0379	-12.0	0.0143	4.5	-0.0137	-4.3

We see that the TP  $\epsilon_5$  gives rise to asymmetries and mean values that are clearly higher than those obtained from  $\epsilon_6$  and  $\epsilon_7$ . This is in contrast to the TPs  $\epsilon_{1,2,3}$ , for which the asymmetries and mean values are comparable (see Tables I and V). We also note that the asymmetry for  $\epsilon_5$  is exactly the same as for  $\epsilon_1$  as was expected from Eq. (32) since the proportionality factor relating them is positive definite. Regarding the mean values, we see by comparing Tables V and VI that the TPs  $\epsilon_{1,2,3}$  appear to have a higher sensitivity to the pseudoscalar coupling than  $\epsilon_{5,6,7}$ .

It is important to mention that in the  $t\bar{t}$  rest frame the sign of the TP  $\epsilon_5$  is defined through the angle  $\Delta\phi_{\ell-\ell^+}$  (recall the discussion on Eq. (31)), which is the angular difference between the projections of the leptons momenta onto the plane perpendicular to  $\vec{t}$ . By a similar argument to that discussed at the beginning of Sec. III B, we can construct an associated angular distribution that, being constrained by the  $\mathcal{A}(\epsilon_5)$ , will be also sensitive to the sign of the pseudoscalar coupling. This angular variable is nothing but that proposed in [?] as a useful CP-odd observable. Moreover, it is shown in [?] that this angular distribution follows the functional form given in Eq. (30). Due to the fact that this distribution is constrained by  $\mathcal{A}(\epsilon_5)$  which is equal to  $\mathcal{A}(\epsilon_1)$  and smaller than  $\mathcal{A}(\epsilon_{2,3})$ , the corresponding shifts ( $\delta$ ) obtained for different values of  $\tilde{\kappa}_t$  are expected to be of the same order than those exhibited by the  $\Delta\phi_1(n_t, n_{\bar{t}})$  distribution and somewhat smaller than those observed in the  $\Delta\phi_2(n_t, n_{\bar{t}})$  distribution.

In analogy to the combination of TPs considered in Sec. III, we have found a combination of the TPs  $\epsilon_{5,6,7}$  for which the asymmetry is enhanced with respect to  $\epsilon_5$ - $\epsilon_7$ ,

$$\epsilon_8 = 2\epsilon_5 - \epsilon_6 + \epsilon_7 = \epsilon(t + \bar{t} + H, t - \bar{t}, \ell^+, \ell^-). \quad (35)$$

We see from Eq. (35) that in the  $t\bar{t}H$  rest frame  $\epsilon_8 = M_{t\bar{t}H}(\vec{t} - \vec{\bar{t}}) \cdot (\vec{\ell}^+ \times \vec{\ell}^-)$ , where  $M_{t\bar{t}H}$  is the invariant mass of the system  $t\bar{t}H$ . Hence, in this reference frame the sign of the combination  $\epsilon_8$  is set by the quantity  $(\vec{t} - \vec{\bar{t}}) \cdot (\vec{\ell}^+ \times \vec{\ell}^-)$ . Taking into account Eqs. (27) and (35) along with the definition  $Q = (t + \bar{t} + H)/2$ , we see that the only relevant difference between  $\epsilon_4$  and  $\epsilon_8$  is that in the latter the spin vectors  $n_t$  and  $n_{\bar{t}}$  have been replaced by the momenta of the leptons  $\ell^+$  and  $\ell^-$  respectively. The values obtained for  $\mathcal{A}(\epsilon_8)$  are shown in Table VIII. Compared to the TPs  $\epsilon_5$ - $\epsilon_7$  and  $\epsilon_1, \epsilon_3$  (see Tables I and VII), the chosen combination exhibits a slightly higher sensitivity in the asymmetry for resolving the CP-mixed cases. This is not true in the case of the TPs  $\epsilon_2$  and  $\epsilon_4$ , for which the asymmetry is larger (see Tables I, II and VIII). In particular, the comparison between  $\epsilon_4$  and  $\epsilon_8$  indicates that the use of the momenta of the leptons instead of the spin vectors produces a decrease in the sensitivity of the asymmetry.

From the TPs  $\epsilon_5$ - $\epsilon_7$  one can derive various angular distributions in the same manner we have discussed in Sec. III B for  $\epsilon_1$ - $\epsilon_3$ . Of course, in this case the corresponding angle will be defined in terms of the momenta of the leptons instead of using the spin vectors. These distributions have the same behaviour than those derived from  $\epsilon_1$ - $\epsilon_3$ , but only the shift obtained in the case of  $\epsilon_5$  is compatible to the values given in Tables for  $\epsilon_1$ - $\epsilon_3$ . The shifts resulting from the fit of the distributions related to  $\epsilon_6$  and  $\epsilon_7$  are smaller. Taking into account these facts we do not give further details of the angular distributions for the TPs discussed in this section.

The mean value of  $\epsilon_8$  for the hypotheses under consideration is comparable with the values listed in Table VI for  $\epsilon_5$ . Concerning the associated angular distributions, their range of variation is larger than in the case of the distributions related to  $\epsilon_5$ - $\epsilon_7$  but exhibit smaller changes in their shapes for the different hypothesis and for this reason we do not include here the corresponding plots.

TABLE VIII: Asymmetry for the TP  $\epsilon_8$  for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . The values are obtained with  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_8)$	$\mathcal{A}(\epsilon_8)/\sigma_{\mathcal{A}}$
1	-1	0.0331	10.5
1	0	0.0023	0.7
1	1	-0.0403	-12.7

## V. CP-ODD OBSERVABLES NOT DEPENDING ON $t$ AND $\bar{t}$ MOMENTA

All the observables discussed in the above sections involve the momenta of the top and anti-top quarks thus requiring the full reconstruction of the kinematics of the individual  $t$  and  $\bar{t}$  systems in order to be measured. Although challenging due to the presence of two neutrinos in the final state, this can be in principle done by applying a kinematic reconstruction method such that the neutrino weighting technique [? ?]. Another possibility is to define observables that are not dependent on the  $t$  and  $\bar{t}$  momenta but make use of the quarks  $b$  and  $\bar{b}$ . In order to construct such observables we will modify here the most sensitive observables studied in Secs. III and IV, namely the combinations  $\epsilon_4$  and  $\epsilon_8$  respectively. Let us first consider the combination  $\epsilon_8$  defined in Eq. (35) and replace the top and antitop momenta by the bottom and antibottom momenta respectively. Thus, we define

$$\epsilon_9 = \epsilon(b + \bar{b} + H, b - \bar{b}, \ell^+, \ell^-). \quad (36)$$

Note that now the sign of the TP  $\epsilon_9$  is set by the sign of the quantity  $(\vec{b} - \vec{\bar{b}}) \cdot (\vec{\ell}^+ \times \vec{\ell}^-)$  in the  $b\bar{b}H$  rest frame. This quantity is in fact used in [?] but in the lab frame in order to define a CP-odd observable that only depends on lab frame variables. The values of the asymmetry for  $\epsilon_9$  are listed in Table IX. By comparing Tables VIII and IX we see that the use of the  $b, \bar{b}$  instead of  $t, \bar{t}$  leads to a decrease in the sensitivity of the asymmetry by  $\sim 5\sigma$  for  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . However, the observable is still capable of discriminating not only between the two CP-mixed hypotheses but also between these and the SM case. We proceed now in a similar way with the combination  $\epsilon_4$ . By using Eq. (27) along with the definitions of the spin vectors in Eqs. (3)-(4), we can write

$$\epsilon_4 = \frac{m_t^2}{(t \cdot \ell^+) \cdot (\bar{t} \cdot \ell^-)} \epsilon(Q, t - \bar{t}, \ell^-, \ell^+) + \frac{1}{(t \cdot \ell^+)} \epsilon(Q, t, \ell^+, \bar{t}) - \frac{1}{(\bar{t} \cdot \ell^-)} \epsilon(Q, \bar{t}, t, \ell^-). \quad (37)$$

Since the asymmetry is not changed by an overall positive definite factor, we will concentrate on the following combination arising from the expression in Eq. (37),

$$\epsilon(Q, t - \bar{t}, \ell^-, \ell^+) + \frac{(\bar{t} \cdot \ell^-)}{m_t^2} \epsilon(Q, t, \ell^+, \bar{t}) - \frac{(t \cdot \ell^+)}{m_t^2} \epsilon(Q, \bar{t}, t, \ell^-), \quad (38)$$

TABLE IX: Asymmetry for the TP  $\epsilon_9$  for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . The values are obtained with  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_9)$	$\mathcal{A}(\epsilon_9)/\sigma_{\mathcal{A}}$
1	-1	0.0171	5.4
1	0	0.0010	0.3
1	1	-0.0247	-7.8

and instead of replacing  $t$  and  $\bar{t}$  directly by  $b$  and  $\bar{b}$ , we use their visible parts, namely  $b + \ell^+$  and  $\bar{b} + \ell^-$  respectively. This results in the following definition

$$\epsilon_{10} = \epsilon(\tilde{Q}, c_{b\bar{b}}, \ell^-, \ell^+) - w_1 \epsilon(\tilde{Q}, b, \bar{b}, \ell^+) + w_2 \epsilon(\tilde{Q}, b, \bar{b}, \ell^-), \quad (39)$$

where  $\tilde{Q} \equiv (b + \ell^+ + \bar{b} + \ell^-)/2$  stands for the visible part of  $Q$ ,  $c_{b\bar{b}} = (1 - w_1)b - (1 - w_2)\bar{b}$  and the weights  $w_{1,2}$  are given by  $(\bar{b} \cdot \ell^-)/m_t^2$  and  $(b \cdot \ell^+)/m_t^2$  respectively. Also, the contribution  $m_\ell^2/m_t^2$  has been neglected both in  $w_1$  and in  $w_2$ . Note that if we set  $w_1 = w_2 = 0$ , the combination  $\epsilon_{10}$  reduces to  $\epsilon_9/2$  and  $\mathcal{A}(\epsilon_{10})$  becomes equal to  $\mathcal{A}(\epsilon_9)$ . The results obtained for the asymmetry of  $\epsilon_{10}$  are given in Table X. By comparing Tables II and X we see again that the sensitivity of the asymmetry decreases when  $t$  and  $\bar{t}$  are not included in the TP. Nevertheless, the combination  $\epsilon_{10}$  remains a useful observable for discriminating the CP nature of the Higgs boson, with the corresponding asymmetry having a sensitivity even higher than that of  $\epsilon_9$ , as can be checked by taking into account Tables IX and X. More precisely, the separation between the CP-mixed hypotheses is enhanced by about  $3\sigma$ . This improvement in the asymmetry of  $\epsilon_{10}$  with respect to  $\epsilon_9$  may be due to two facts. In the first place, as was pointed out in Sec. IV when comparing the TPs  $\epsilon_4$  and  $\epsilon_8$ , the asymmetry appears to be higher when the spin vectors are used instead of the lepton momenta and we see from Eqs. (36) and (39) that  $\epsilon_{10}$ , being obtained from  $\epsilon_4$ , contains the information on the spin vectors, as opposed to  $\epsilon_9$  that depends directly on the lepton momenta because it is derived from  $\epsilon_8$ . In the second place, in order to obtain  $\epsilon_{10}$  we have replaced in the expression for  $\epsilon_4$  the top and antitop momenta by their visible part, while in the case of  $\epsilon_9$  the bottom and antibottom momenta have been used.

TABLE X: Asymmetry for the TP  $\epsilon_{10}$  for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ . The values are obtained by using  $10^5$  simulated events.

$\kappa_t$	$\tilde{\kappa}_t$	$\mathcal{A}(\epsilon_{10})$	$\mathcal{A}(\epsilon_{10})/\sigma_{\mathcal{A}}$
1	-1	-0.0213	-6.7
1	0	0.0031	1.0
1	1	0.0300	9.5

By using our set of simulated events we have also tested, for comparison purposes, the asymmetry of the lab frame observable given in [? ]. We have obtained that this observable appears to be slightly less sensitive than the combination  $\epsilon_{10}$ , giving rise to a separation between the CP-mixed hypotheses smaller by about  $1.4\sigma$ .

## VI. EXPERIMENTAL FEASIBILITY

By considering the mild selection cuts introduced in Sec. III, the SM cross section for  $pp \rightarrow t (\rightarrow b\ell^+\nu_\ell) \bar{t} (\rightarrow \bar{b}\ell^-\bar{\nu}_\ell) H$ ,  $\ell = e, \mu$  at 14 TeV is  $\sim 15.3 \text{ fb}$  and hence the number of events expected within the context of the HL-LHC is  $\sim 15.3 \text{ fb} \times 3000 \text{ fb}^{-1} = 4.59 \times 10^4$ . This number is expected to be even larger in the case of  $\kappa_t = 1$ ,  $\tilde{\kappa}_t \neq 0$  since the corresponding cross section is then higher than the SM one. By taking into account NLO corrections to the production process and considering a  $K$ -factor around 1.2 [? ? ? ], the number of events can be raised up to  $\sim 5.49 \times 10^4$ . On the other hand, additional cuts as well as the efficiency related to the reconstruction of particles momenta will contribute to decrease this number. For instance, in order to obtain the asymmetry of  $\epsilon_4$ , the  $t$  and  $\bar{t}$  momenta need to be reconstructed. This is challenging not only due to the presence of two neutrinos in the final state which escape the detector undetected but also because final state objects need to be associated with the respective parent quark [? ]. As was already mentioned in Sec. V, a possibility is to use the neutrino weighting technique along with the kinematic equations arising from kinematic constraints related to the top and  $W$  masses as well as from the energy-momentum conservation at each of the decay vertices involved in the process. Within the context of  $t\bar{t}$  this procedure has been used, for instance, in order to obtain measurements of spin correlation [? ] and charge asymmetry [? ]. Also, events reconstructed with this technique has been used in [? ] for analyzing angular distributions that are useful for discriminating the signal from the backgrounds in  $t\bar{t}H(H \rightarrow b\bar{b})$ . In all these cases the corresponding efficiency in the reconstruction of the momenta is up to  $\sim 80\%$ .

Based on the discussion given in the previous paragraph, we have simulated sets of  $5 \times 10^4$ ,  $1 \times 10^4$  and  $5 \times 10^3$  events and recalculated the most sensitive observable, namely  $\mathcal{A}(\epsilon_4)$ , for each case. The results are displayed in Table XI, where it can be seen that for a number of events close to that roughly estimated above within the context of the HL-LHC, the observable is still sensitive to  $\tilde{\kappa}_t$  allowing to separate the CP-mixed cases by  $19\sigma$ . As expected, the sensitivity worsen as the number of events is reduced, but even with  $5 \times 10^3$  events the separation between the CP-mixed hypotheses under consideration is around  $6.5\sigma$ .

Although the combination  $\epsilon_{10}$  discussed in Sec. V avoids the problem of reconstructing the top and antitop momenta, we have also considered the respective asymmetry obtained for more conservative numbers of events. In Table XII we show the results for  $\mathcal{A}(\epsilon_{10})$  when the number of events is reduced from  $5 \times 10^4$  to  $1 \times 10^4$ . We see in this case that even with  $1 \times 10^4$  simulated events the observable is capable to discriminate the CP-mixed cases by  $5.6\sigma$ .

Finally, it is important to mention that a realistic analysis of the sensitivity of the observables discussed in this paper requires the study of the impact of the backgrounds as well as the hadronization of the quarks in the final state and the effects of the detector. If we consider the dominant decay mode of the higgs,  $H \rightarrow b\bar{b}$ , in order to maximize the cross section of the process, the signature is given by 4  $b$ -jets, two leptons and missing energy, while the main background arise from the production of  $t\bar{t}$  in association with additional jets, being the dominant source the production of  $t\bar{t} + b\bar{b}$ . By applying a small set of cuts it is shown in [? ] that the

TABLE XI: Asymmetry for the TP  $\epsilon_4$  obtained by using  $5 \times 10^4$ ,  $1 \times 10^4$  and  $5 \times 10^3$  simulated events for the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ .

$\kappa_t$	$\tilde{\kappa}_t$	$N_{\text{ev}} = 5 \times 10^4$		$N_{\text{ev}} = 1 \times 10^4$		$N_{\text{ev}} = 5 \times 10^3$	
		$\mathcal{A}(\epsilon_4)$	$\mathcal{A}(\epsilon_4)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_4)$	$\mathcal{A}(\epsilon_4)/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_4)$	$\mathcal{A}(\epsilon_4)/\sigma_{\mathcal{A}}$
1	-1	-0.0405	-9.1	-0.0426	-4.3	-0.0496	-3.5
1	0	0.0004	0.1	-0.0084	-0.8	-0.0004	-0.03
1	1	0.0443	9.9	0.0434	4.2	0.0420	3.0

TABLE XII: Asymmetry for the TP  $\epsilon_{10}$  in the SM case and the two CP-mixed cases defined by  $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$  when the number of simulated events is reduced from  $5 \times 10^4$  to  $1 \times 10^4$ .

$\kappa_t$	$\tilde{\kappa}_t$	$N_{\text{ev}} = 5 \times 10^4$		$N_{\text{ev}} = 1 \times 10^4$	
		$\mathcal{A}(\epsilon_{10})$	$\mathcal{A}(\epsilon_{10})/\sigma_{\mathcal{A}}$	$\mathcal{A}(\epsilon_{10})$	$\mathcal{A}(\epsilon_{10})/\sigma_{\mathcal{A}}$
1	-1	-0.0270	-6.0	-0.0184	-1.8
1	0	0.0022	0.5	-0.0086	-0.9
1	1	0.0313	7.0	0.0380	3.8

signal to background ratio is largely improved. On the other hand, a rigorous treatment of the backgrounds and their impact with respect to the signal is performed in [?] by using  $20.3 \text{ fb}^{-1}$  of data at  $\sqrt{s} = 8 \text{ TeV}$ .

The results shown in tables XI and XII reveal that with  $5 \times 10^3$  and  $1 \times 10^4$  events respectively the observables  $\mathcal{A}(\epsilon_4)$  and  $\mathcal{A}(\epsilon_{10})$  are still useful for testing  $\tilde{\kappa}_t$ . Without considering the lack of events related to the experimental analysis, these numbers of events correspond to a luminosity around  $\sim 300$ - $600 \text{ fb}^{-1}$  for the SM and even smaller for the CP-mixed cases due to the larger cross section. This range of luminosities is in principle achievable in the short term by the LHC. We note that in order to be fully conclusive about the required luminosity, it is important to include the effects of hadronization, detector resolution, reconstruction efficiencies and so forth. However, this kind of analysis is out of the scope of this paper.

## VII. CONCLUSIONS

In this paper we have presented a collection of CP-odd observables based on triple product correlations that are useful for disentangling the relative sign between the scalar ( $\kappa_t$ ) and a potential pseudoscalar ( $\tilde{\kappa}_t$ ) top-Higgs couplings in the  $t\bar{t}H$  production with both tops decaying



leptonically. We have tested the sensitivity of the proposed observables by considering three types of observables: asymmetries, mean values and angular distributions.

Through the use of spinor techniques we have written the expression for the differential cross section of the full process in such a manner that the production and the decay parts are separated, although connected by the spin vectors of the top and antitop which are given in terms of the momenta of the leptons in the final state. Moreover, we have indentified the terms linear in  $\kappa_t$  and  $\tilde{\kappa}_t$  as those involving TPs. Among these, we have explored the three that do not involve the momenta of the incoming quarks/gluons and at the same time incorporate both spin vectors:  $\epsilon_1 \equiv \epsilon(t, \bar{t}, n_t, n_{\bar{t}})$ ,  $\epsilon_2 \equiv \epsilon(Q, \bar{t}, n_t, n_{\bar{t}})$  and  $\epsilon_3 \equiv \epsilon(Q, t, n_t, n_{\bar{t}})$ .

We have found that  $\epsilon_{1,2,3}$  allow to distinguish between the CP-mixed hypotheses by more than  $\sim 20\sigma$  in the case of asymmetries and  $\sim 10\sigma$  in the case of mean values when  $1 \times 10^5$  simulated events are used. Furthermore, we have shown that the angular distributions associated with these TPs are also sensitive to the value of  $\kappa_t$  and  $\tilde{\kappa}_t$  exhibiting a phase shift that varies according to the values taken by these couplings. On the other hand, by exploring TPs that incorporate the momenta of the Higgs and the leptons instead of the spin vectors we can conclude that the observables studied here appear to be more sensitive when the spin vectors are used.

On the other hand, we have proposed a combination of the TPs considered in the first place,  $\epsilon_4 \equiv \epsilon_3 - \epsilon_2$ , that exhibits the highest sensitivity with the separation between the CP-mixed hypotheses being increased by at least  $3\sigma$  in the asymmetry for  $1 \times 10^5$  events with respect to  $\mathcal{A}(\epsilon_{1,2,3})$ . Again, when a similar combination is constructed by using the leptons momenta instead of the spin vectors ( $\epsilon_8$ ), the sensitivity in the asymmetry is decreased by  $\sim 3\sigma$  compared to  $\mathcal{A}(\epsilon_4)$  for the same number of events, giving values compatible with those obtained for the asymmetry of  $\epsilon_2$  and  $\epsilon_3$ .

Taking into account the challenge of reconstructing the top and antitop momenta due to the presence of two neutrinos in the final state, we have proposed and tested two TP correlations that avoid this difficulty. The first one is obtained by replacing the  $t$  and  $\bar{t}$  by the  $b$  and  $\bar{b}$  momenta ( $\epsilon_9$ ), whereas the second includes the visible part of the  $t$  and  $\bar{t}$  momenta ( $\epsilon_{10}$ ). We have encountered that the latter is the most sensitive leading to a discrimination of the CP-mixed cases under analysis by up to  $\sim 16\sigma$ .

Finally we have discussed the feasibility of the most sensitive observables proposed here. We have found that with  $5 \times 10^3$  and  $1 \times 10^4$  events respectively the observables  $\mathcal{A}(\epsilon_4)$  and  $\mathcal{A}(\epsilon_{10})$  are still useful for testing the hypotheses ( $\kappa_t = 1, \tilde{\kappa}_t = \pm 1$ ) giving rise to separations of order  $\sim 6\sigma$ . These required number of events would be reachable in principle in the short term by the LHC and hence the capability of the observables studied here for testing the sign of  $\tilde{\kappa}_t/\kappa_t$  could be probed in that context.

**Acknowledgments** This work has been partially supported by ANPCyT under grants No. PICT 2013-0433 and No. PICT 2013-2266, and by CONICET (NM, AS). The work of KK was supported by the U.S. National Science Foundation under Grant PHY-1215785. KK also acknowledges sabbatical support from Taylor University.

## REFERENCES