



Tips for General Practitioners in Managing Hypertension

Hypertension management requires a comprehensive approach that includes accurate diagnosis, risk assessment, lifestyle modifications, and appropriate medication. Here are key tips for general practitioners (GPs) to effectively manage hypertensive patients.

1. Accurate Diagnosis & BP Measurement

- ✓ Use proper technique:
 - Patient should be seated for 5 minutes, feet flat on the floor, and arm supported at heart level.
 - Avoid caffeine, smoking, or exercise 30 minutes before measurement.
 - Use an appropriately sized cuff (too small = falsely high readings).
 - Take at least two readings on different occasions to confirm hypertension.
- ✓ Consider Ambulatory Blood Pressure Monitoring (ABPM):
 - Helps diagnose white coat hypertension or masked hypertension.

2. Risk Stratification & Individualized Treatment

- ✓ Assess cardiovascular risk factors:
 - Check for diabetes, smoking, high cholesterol, kidney disease, family history of CVD.
 - Use risk calculators like ASCVD Risk Score to guide treatment intensity.
- ✓ Look for target organ damage:
 - ECG for LVH, arrhythmias.
 - Fundoscopy for hypertensive retinopathy.
 - Kidney function tests (serum creatinine, eGFR, urine albumin) for hypertensive nephropathy.
- ✓ Identify secondary hypertension causes if:
 - Sudden onset or severe hypertension (<40 years).
 - Resistant hypertension despite 3+ drugs.
 - Suspicion of hyperaldosteronism, renal artery stenosis, pheochromocytoma.

3. Lifestyle Modification Counseling (First-Line for All Patients)

- ✓ Dietary changes:



HyperTensia 2025



- DASH diet (Dietary Approaches to Stop Hypertension): High in fruits, vegetables, whole grains, low in sodium.
 - Reduce salt intake (<2.3g/day) and processed foods.
 - Increase potassium intake (bananas, oranges, spinach).
 - ✓ Weight management:
 - Aim for BMI <25 kg/m².
 - Even 5-10% weight loss can lower BP significantly.
 - ✓ Physical activity:
 - Encourage 30–45 minutes of moderate exercise (walking, cycling) 5 days a week.
 - ✓ Limit alcohol & smoking:
 - Men: ≤2 drinks/day, Women: ≤1 drink/day.
 - Strongly advise smoking cessation.
 - ✓ Stress management:
 - Teach relaxation techniques, meditation, and sleep hygiene.
-

4. Pharmacological Treatment: Choosing the Right Drug

- ✓ Follow guideline-based therapy: (e.g., AHA/ESC/ISH guidelines)
 - First-line choices:
 - ACE inhibitors (Lisinopril, Ramipril) or ARBs (Losartan, Telmisartan) → Best for diabetes, CKD, heart disease.
 - Calcium channel blockers (Amlodipine, Nifedipine) → Best for elderly, isolated systolic hypertension.
 - Diuretics (Hydrochlorothiazide, Chlorthalidone) → Good for salt-sensitive hypertension.
 - Beta-blockers (Metoprolol, Carvedilol) → Best for post-MI, heart failure.
- ✓ Combination therapy for better control:
 - Two-drug therapy (ACEI/ARB + CCB or Diuretic) if BP >150/90 mmHg.
- ✓ Adjust treatment for special populations:
 - Pregnancy: Use Labetalol, Methyldopa, or Nifedipine. Avoid ACEIs & ARBs.
 - Resistant hypertension: Add Spironolactone or consider Renal Denervation.



HyperTensia 2025



✓ Monitor for side effects:

- ACE inhibitors → Cough, hyperkalemia.
- CCBs → Edema, headache.
- Diuretics → Electrolyte imbalances.

5. Regular Monitoring & Patient Engagement

✓ Follow-up every 2–4 weeks after starting treatment, then every 3–6 months once BP is controlled.

✓ Encourage home BP monitoring for better compliance.

✓ Educate patients about medication adherence—common reason for treatment failure.

6. Recognizing Hypertensive Emergencies (Immediate Referral)

BP >180/120 mmHg with target organ damage (chest pain, breathlessness, confusion, kidney failure).

✓ Urgent hospital referral for IV antihypertensives (e.g., Labetalol, Nicardipine).


BIHS
British and Irish Hypertension Society
Committed to preventing cardiovascular disease

Adult Hypertension Pathway Therapeutic Management

Hypertension despite lifestyle measures^(a)
Confirmed using 24 hr ABPM or home average
IF Home Average BP $\geq 150/95$ mmHg
(clinic $\geq 160/100$ mmHg)
DIRECT TO STEP 2

Referral to a hypertension specialist may be necessary in some cases^(c). This pathway should not be used in pregnancy, malignant hypertension or hypertensive emergencies^(d).

Treat to Target
 $<135/85$ mmHg^(b)
(clinic $<140/90$ mmHg)





HyperTensia 2025



Conclusion:

- ✓ Accurate diagnosis, lifestyle modifications, and tailored drug therapy are key to effective hypertension management.
- ✓ Patient education & regular follow-ups improve long-term outcomes.
- ✓ Early recognition of complications (LVH, kidney disease, stroke risk) is crucial.