

Lagrange Interpolation for Equally-Spaced abscissa

Kunal Kishore

Theorem (Lagrange Interpolation for Equally-Spaced abscissa). *Suppose a, b are a pair of real number such that $a < b$, $m \geq 3$ be an integer and $h = \frac{b-a}{m-1}$. let $f \in C^m[a-mh, b+mh]$ and equispaced points x_k be defined as $x_k = \frac{b-a}{2} + kh$. Then, for any real number p there exists a real number ξ , $-mh < \xi < mh$, such that*

$$f(x_0 + ph) = \sum_k \mathcal{A}_k^m f(x_k) + \mathcal{R}_{m-1,p} \quad (1)$$

where k varies from

$$\begin{aligned} -\frac{1}{2}(m-2) \leq k \leq \frac{1}{2}m & \text{ for even } m \\ -\frac{1}{2}(m-1) \leq k \leq \frac{1}{2}(m-1) & \text{ for odd } m \end{aligned}$$

with

$$\mathcal{A}_k^m(p) = \frac{(-1)^{\frac{m-1}{2}+k}}{(\frac{m-1}{2}+k)!(\frac{m-1}{2}-k)!(p-k)} \sum_t (p-t)$$

and

$$\mathcal{R}_{m-1,p} = \frac{h^m}{m!} f^{(m)}(\xi) \sum_n (p-n)$$

here both t and n varies same as k .

Proof. Assuming m to be odd. For a function $f \in C^m[a-mh, b+mh]$, its Lagrange interpolation polynomial constructed from equispaced points

$$x_{-\frac{1}{2}(m-1)}, x_{-\frac{1}{2}(m-1)+1}, \dots, x_{\frac{1}{2}(m-1)-1}, x_{\frac{1}{2}(m-1)}$$

is given by

$$\mathcal{L}_{m-1}(x) = \sum_{j=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} l_j(x_j) f(x_j) \quad (2)$$

and,

$$f(x) = \sum_{j=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} l_j(x_j) f(x_j) + R_{m-1}(x) \quad (3)$$

with,

$$l_j(x) = \sum_{i=-\frac{1}{2}(m-1), i \neq j}^{\frac{1}{2}(m-1)} \frac{x - x_i}{x - x_j}$$

$$R_{m-1}(x) = \frac{f^{(m)}(\xi)}{m!} \left[\sum_{j=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} x - x_j \right]$$

We can write x as $x = x_0 + ph$ for some real number p .

$$\begin{aligned} R_{m-1}(x) &= R_{m-1}(x_0 + ph) = \frac{f^{(m)}(\xi)}{m!} \left[\sum_{j=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} x - x_j \right] \\ &= \frac{f^{(m)}(\xi)}{m!} \left[(x - x_{-\frac{1}{2}(m-1)}) (x - x_{-\frac{1}{2}(m-1)+1}) \dots (x - x_{\frac{1}{2}(m-1)}) \right] \\ &= \frac{f^{(m)}(\xi)}{m!} \left[(x_0 + ph - x_0 + \frac{1}{2}(m-1)h) \dots (x_0 + ph - x_0 - \frac{1}{2}(m-1)h) \right] \\ &= \frac{f^{(m)}(\xi)}{m!} \left[(ph + \frac{1}{2}(m-1)h) \dots (ph - \frac{1}{2}(m-1)h) \right] \\ &= \frac{f^{(m)}(\xi)}{m!} \left[h^m \left(p + \frac{1}{2}(m-1) \right) \dots \left(p - \frac{1}{2}(m-1) \right) \right] \\ &= \frac{f^{(m)}(\xi)}{m!} \left[h^m \left(p + \frac{1}{2}(m-1) \right) \dots \left(p - \frac{1}{2}(m-1) \right) \right] \\ &= \frac{f^{(m)}(\xi)}{m!} h^m \sum_{n=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} (p - n) \\ &= \mathcal{R}_{m-1,p} \end{aligned} \quad (4)$$

$$\begin{aligned} l_k(x) &= l_k(x_0 + ph) = \sum_{i=-\frac{1}{2}(m-1), i \neq k}^{\frac{1}{2}(m-1)} \frac{x - x_i}{x_k - x_i} \\ &= \left[\frac{(x - x_{-\frac{1}{2}(m-1)}) \dots (x - x_{k-1}) (x - x_{k+1}) \dots (x - x_{\frac{1}{2}(m-1)})}{(x_k - x_{-\frac{1}{2}(m-1)}) \dots (x_k - x_{k-1}) (x_k - x_{k+1}) \dots (x_k - x_{\frac{1}{2}(m-1)})} \right] \end{aligned}$$

$$\begin{aligned}
&= \left[\frac{\begin{aligned} &\left(x_0 + ph - x_0 + \frac{1}{2}(m-1)h\right) \dots \left(x_0 + ph - x_0 - (k-1)h\right) \left(x_0 + ph - x_0 - (k+1)h\right) \\ &\dots \left(x_0 + ph - x_0 - \frac{1}{2}(m-1)h\right) \end{aligned}}{\begin{aligned} &\left(x_0 + kh - x_0 + \frac{1}{2}(m-1)h\right) \dots \left(x_0 + kh - x_0 - (k-1)h\right) \left(x_0 + kh - x_0 - (k+1)h\right) \\ &\dots \left(x_0 + kh - x_0 - \frac{1}{2}(m-1)h\right) \end{aligned}} \right] \\
&= \left[\frac{\begin{aligned} &\left(p + \frac{1}{2}(m-1)\right) \dots \left(p - (k-1)\right) \left(p - (k+1)\right) \dots \left(p - \frac{1}{2}(m-1)\right) \end{aligned}}{\begin{aligned} &\left(k + \frac{1}{2}(m-1)\right) \dots \left(k - (k-1)\right) \left(k - (k+1)\right) \dots \left(k - \frac{1}{2}(m-1)\right) \end{aligned}} \right] \frac{(p-k)}{(p-k)} \\
&= \left[\frac{\begin{aligned} &\left(p + \frac{1}{2}(m-1)\right) \dots \left(p - k + 1\right) \left(p - k\right) \left(p - k - 1\right) \dots \left(p - \frac{1}{2}(m-1)\right) \end{aligned}}{\begin{aligned} &\left(k + \frac{1}{2}(m-1)\right) \dots \left(k - k + 1\right) \left(k - k - 1\right) \dots \left(k - \frac{1}{2}(m-1)\right) \end{aligned}} \frac{1}{(p-k)} \right] \\
&= \frac{\alpha}{\beta}
\end{aligned} \tag{5}$$

where $\alpha = (p + \frac{1}{2}(m-1)) \dots (p - k + 1) (p - k) (p - k - 1) \dots (p - \frac{1}{2}(m-1))$,
 $\beta = (k + \frac{1}{2}(m-1)) \dots (k - k + 1) (k - k - 1) \dots (k - \frac{1}{2}(m-1)) (p - k)$
Now,

$$\begin{aligned}
\beta &= \left(k + \frac{1}{2}(m-1)\right) \dots \left(k - k + 1\right) \left(k - k - 1\right) \dots \left(k - \frac{1}{2}(m-1)\right) (p - k) \\
&= \left(k + \frac{1}{2}(m-1)\right) \dots (2)(1)(-1)(-2) \dots \left(k - \frac{1}{2}(m-1)\right) (p - k) \\
&= \left(k + \frac{1}{2}(m-1)\right)! (-1)(-2) \dots \left(k - \frac{1}{2}(m-1)\right) (p - k) \\
&= \left(k + \frac{1}{2}(m-1)\right)! (-1)^{\frac{m-1}{2}-k} (1)(2) \dots \left(\frac{1}{2}(m-1) - k\right) (p - k) \\
&= \left(\frac{1}{2}(m-1) + k\right)! (-1)^{\frac{m-1}{2}-k} \left(\frac{1}{2}(m-1) - k\right)! (p - k) \\
\alpha &= \prod_{t=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} (p - t)
\end{aligned}$$

putting α, β back in equation 5

$$\begin{aligned}
l_k(x) &= \frac{\alpha}{\beta} \\
&= \frac{\prod_{t=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} (p - t)}{\left(\frac{1}{2}(m-1) + k\right)! (-1)^{\frac{m-1}{2}-k} \left(\frac{1}{2}(m-1) - k\right)! (p - k)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(-1)^{m-1}}{(-1)^{m-1}} \frac{(-1)^{k-\frac{m-1}{2}} \prod_{t=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} (p-t)}{\left(\frac{1}{2}(m-1)+k\right)! \left(\frac{1}{2}(m-1)-k\right)! (p-k)} \\
&= \frac{(-1)^{\frac{m-1}{2}+k}}{\left(\frac{m-1}{2}+k\right)! \left(\frac{m-1}{2}-k\right)! (p-k)} \prod_{t=-\frac{1}{2}(m-1)}^{\frac{1}{2}(m-1)} (p-t) \\
&= \mathcal{A}_k^m(p)
\end{aligned} \tag{6}$$

By using equations 6 , 3 , 4

$$f(x) = f(x_0 + ph) = \sum_k \mathcal{A}_k^m f(x_k) + \mathcal{R}_{m-1,p}$$

where k varies from $-\frac{1}{2}(m-1) \leq k \leq \frac{1}{2}(m-1)$, m is odd.

Similarly, for even m , we can get the result by constructing Lagrange interpolating polynomial from equispaced points

$$x_{-\frac{1}{2}(m-2)}, x_{-\frac{1}{2}(m-2)+1}, \dots, x_{\frac{1}{2}m-1}, x_{\frac{1}{2}m}$$

□

References

- [1] Sharad Kapur and Vladimir Rokhlin. High-order corrected trapezoidal quadrature rules for singular functions. *SIAM Journal on Numerical Analysis*, 34(4):1331–1356, 1997.
- [2] A Erdelyi, W Magnus, and F Oberhettinger. M. abramowitz and ia stegun, handbook of mathematical functions, 1972.