

## WORKED EXAMPLES 1

### TOTAL PROBABILITY AND BAYES' THEOREM

**EXAMPLE 1.** A biased coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed repeatedly and independently until the first head is observed. Compute the probability that the first head appears at an **even** numbered toss.

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible infinite binary sequences of coin tosses;
- event  $H_1$  - head on **first** toss;
- event  $E$  - first head on even numbered toss.

We want  $P(E)$  : using the Theorem of Total Probability, and the partition of  $\Omega$  given by  $\{H_1, H'_1\}$

$$P(E) = P(E|H_1)P(H_1) + P(E|H'_1)P(H'_1).$$

Now clearly,  $P(E|H_1) = 0$  (given  $H_1$ , that a head appears on the first toss,  $E$  cannot occur) and also  $P(E|H'_1)$  can be seen to be given by

$$P(E|H'_1) = P(E') = 1 - P(E),$$

(given that a head does **not** appear on the first toss, the required conditional probability is merely the probability that the sequence concludes after a further **odd** number of tosses, that is, the probability of  $E'$ ). Hence  $P(E)$  satisfies

$$P(E) = 0 \times p + (1 - P(E)) \times (1 - p) = (1 - p)(1 - P(E)),$$

so that

$$P(E) = \frac{1 - p}{2 - p}.$$

Alternatively, consider the partition of  $E$  into  $E_1, E_2, \dots$  where  $E_k$  is the event that the first head occurs on the  $2k$ th toss. Then  $E = \bigcup_{k=1}^{\infty} E_k$ , and

$$P(E) = P\left(\bigcup_{k=1}^{\infty} E_k\right) = \sum_{k=1}^{\infty} P(E_k).$$

Now  $P(E_k) = (1-p)^{2k-1}p$  (that is,  $2k-1$  tails, then a head), so

$$\begin{aligned} P(E) &= \sum_{k=1}^{\infty} (1-p)^{2k-1}p \\ &= \frac{p}{1-p} \sum_{k=1}^{\infty} (1-p)^{2k} = \frac{p}{1-p} \frac{(1-p)^2}{1-(1-p)^2} \\ &= \frac{1-p}{2-p}. \end{aligned}$$

**EXAMPLE 2.** Two players A and B are competing at a trivia quiz game involving a series of questions. On any individual question, the probabilities that A and B give the correct answer are  $\alpha$  and  $\beta$  respectively, for all questions, with outcomes for different questions being independent. The game finishes when a player wins by answering a question correctly.

Compute the probability that A wins if

- (a) A answers the first question,      (b) B answers the first question.

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible infinite sequences of answers;
- event  $A$  - A answers the first question;
- event  $F$  - game ends after the first question;
- event  $W$  - A wins.

We want

$$P(W|A) \quad \text{and} \quad P(W|A').$$

Using the Theorem of Total Probability, and the partition given by  $\{F, F'\}$

$$P(W|A) = P(W|A \cap F)P(F|A) + P(W|A \cap F')P(F'|A).$$

Now, clearly

$$P(F|A) = P[\text{A answers first question correctly}] = \alpha, \quad P(F'|A) = 1 - \alpha,$$

and  $P(W|A \cap F) = 1$ , but  $P(W|A \cap F') = P(W|A')$ , so that

$$P(W|A) = (1 \times \alpha) + (P(W|A') \times (1 - \alpha)) = \alpha + P(W|A')(1 - \alpha). \quad (1)$$

Similarly,

$$P(W|A') = P(W|A' \cap F)P(F|A') + P(W|A' \cap F')P(F'|A').$$

We have

$$P(F|A') = P[\text{B answers first question correctly}] = \beta, \quad P(F'|A) = 1 - \beta,$$

but  $P(W|A' \cap F) = 0$ . Finally  $P(W|A' \cap F') = P(W|A)$ , so that

$$P(W|A') = (0 \times \beta) + (P(W|A) \times (1 - \beta)) = P(W|A)(1 - \beta). \quad (2)$$

Solving (1) and (2) simultaneously gives, for (a) and (b)

$$P(W|A) = \frac{\alpha}{1 - (1 - \alpha)(1 - \beta)}, \quad P(W|A') = \frac{(1 - \beta)\alpha}{1 - (1 - \alpha)(1 - \beta)}.$$

*Note: recall, for any events  $E_1$  and  $E_2$  we have that*

$$P(E'_1|E_2) = 1 - P(E_1|E_2),$$

*but not necessarily that*

$$P(E_1|E'_2) = 1 - P(E_1|E_2).$$

**EXAMPLE 3.** Patients are recruited onto the two arms (0 - Control, 1-Treatment) of a clinical trial. The probability that an adverse outcome occurs on the control arm is  $p_0$ , and is  $p_1$  for the treatment arm. Patients are allocated alternately onto the two arms, and their outcomes are independent. What is the probability that the first adverse event occurs on the control arm?

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible infinite sequences of patient outcomes;
- event  $E_1$  - first patient (allocated onto the control arm) suffers an adverse outcome;
- event  $E_2$  - first patient (allocated onto the control arm) does not suffer an adverse outcome, but the second patient (allocated onto the treatment arm) does suffer an adverse outcome;
- event  $E_0$  - neither of the first two patients suffer adverse outcomes;
- event  $F$  - first adverse event occurs on the control arm.

We want  $P(F)$ . Now the events  $E_1, E_2$  and  $E_0$  partition  $\Omega$ , so, by the Theorem of Total Probability,

$$P(F) = P(F|E_1)P(E_1) + P(F|E_2)P(E_2) + P(F|E_0)P(E_0).$$

Now,  $P(E_1) = p_0$ ,  $P(E_2) = (1 - p_0)p_1$ ,  $P(E_0) = (1 - p_0)(1 - p_1)$  and  $P(F|E_1) = 1$ ,  $P(F|E_2) = 0$ . Finally, as after two non-adverse outcomes the allocation process effectively re-starts, we have  $P(F|E_0) = P(F)$ . Hence

$$\begin{aligned} P(F) &= (1 \times p_0) + (0 \times (1 - p_0)p_1) + (P(F) \times (1 - p_0)(1 - p_1)) \\ &= p_0 + (1 - p_0)(1 - p_1)P(F), \end{aligned}$$

which can be re-arranged to give

$$P(F) = \frac{p_0}{p_0 + p_1 - p_0p_1}.$$

**EXAMPLE 4.** In a tennis match, with the score at deuce, the game is won by the first player who gets a clear lead of two points.

If the probability that given player wins a particular point is  $\theta$ , and all points are played independently, what is the probability that player eventually wins the game?

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible infinite sequences of points;
- event  $W_i$  - nominated player wins the  $i$ th point;
- event  $V_i$  - nominated player wins the game on the  $i$ th point;
- event  $V$  - nominated player wins the game.

We want  $P(V)$ . The events  $\{W_1, W'_1\}$  partition  $\Omega$ , and thus, by the Theorem of Total Probability

$$P(V) = P(V|W_1)P(W_1) + P(V|W'_1)P(W'_1). \quad (3)$$

Now  $P(W_1) = \theta$  and  $P(W'_1) = 1 - \theta$ . To get  $P(V|W_1)$  and  $P(V|W'_1)$ , we need to further condition on the result of the second point, and again use the Theorem: for example

$$\begin{aligned} P(V|W_1) &= P(V|W_1 \cap W_2)P(W_2|W_1) + P(V|W_1 \cap W'_2)P(W'_2|W_1), \\ P(V|W'_1) &= P(V|W'_1 \cap W_2)P(W_2|W'_1) + P(V|W'_1 \cap W'_2)P(W'_2|W'_1), \end{aligned} \quad (4)$$

where

$$\begin{aligned} P(V|W_1 \cap W_2) &= 1, & P(W_2|W_1) &= P(W_2) = \theta, \\ P(V|W_1 \cap W'_2) &= P(V), & P(W'_2|W_1) &= P(W'_2) = 1 - \theta, \\ P(V|W'_1 \cap W_2) &= P(V), & P(W_2|W'_1) &= P(W_2) = \theta, \\ P(V|W'_1 \cap W'_2) &= 0, & P(W'_2|W'_1) &= P(W'_2) = 1 - \theta, \end{aligned}$$

as,

- given  $W_1 \cap W_2$  : the game is over, and the nominated player has won
- given  $W_1 \cap W'_2$  : the game is back at deuce
- given  $W'_1 \cap W_2$  : the game is back at deuce
- given  $W'_1 \cap W'_2$  : the game is over, and the player has lost

and the results of successive points are independent. Thus

$$\begin{aligned} P(V|W_1) &= (1 \times \theta) + (P(V) \times (1 - \theta)) = \theta + (1 - \theta) P(V) \\ P(V|W'_1) &= (P(V) \times \theta) + 0 \times (1 - \theta) = \theta P(V). \end{aligned}$$

Hence, combining (3) and (4) we have

$$\begin{aligned} P(V) &= (\theta + (1 - \theta) P(V)) \theta + \theta P(V) (1 - \theta) = \theta^2 + 2\theta(1 - \theta)P(V) \\ \implies P(V) &= \frac{\theta^2}{1 - 2\theta(1 - \theta)}. \end{aligned}$$

Alternatively,  $\{V_i, i = 1, 2, \dots\}$  partition  $V$ . Hence

$$P(V) = \sum_{i=1}^{\infty} P(V_i).$$

Now,  $P(V_i) = 0$  if  $i$  is odd, as the game can never be completed after an odd number of points. For  $i = 2$ ,  $P(V_2) = \theta^2$ , and for  $i = 2k + 2$  ( $k = 1, 2, 3, \dots$ ) we have

$$P(V_i) = P(V_{2k+2}) = 2^k \theta^k (1 - \theta)^k \times \theta^2$$

- the score must stand at deuce after  $2k$  points and the game must not have been completed prior to this, indicating that there must have been  $k$  successive drawn pairs of points, each of which could be arranged win/lose or lose/win for the nominated player. Then that player must win the final two points. Hence

$$P(V) = \sum_{k=0}^{\infty} P(V_{2k+2}) = \theta^2 \sum_{k=0}^{\infty} \{2\theta(1 - \theta)\}^k = \frac{\theta^2}{1 - 2\theta(1 - \theta)},$$

as the term in the geometric series satisfies  $|2\theta(1 - \theta)| < 1$ .

**EXAMPLE 5** A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained.

Find the probability that the experiment is completed on the  $n$ th toss.

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible infinite sequences of tosses;
- event  $E_1$  : first toss is  $H$ ;
- event  $E_2$  : first two tosses are  $TH$ ;
- event  $E_3$  : first two tosses are  $TT$ ;
- event  $F_n$  : experiment completed on the  $n$ th toss.

We want  $P(F_n)$  for  $n = 2, 3, \dots$ . The events  $\{E_1, E_2, E_3\}$  partition  $\Omega$ , and thus, by the Theorem of Total Probability

$$P(F_n) = P(F_n|E_1)P(E_1) + P(F_n|E_2)P(E_2) + P(F_n|E_3)P(E_3). \quad (5)$$

Now for  $n = 2$

$$P(F_2) = P(E_3) = (1 - p)^2$$

and for  $n > 2$ ,

$$P(F_n|E_1) = P(F_{n-1}), \quad P(F_n|E_2) = P(F_{n-2}), \quad P(F_n|E_3) = 0,$$

as, given  $E_1$ , we need  $n - 1$  further tosses that finish  $TT$  for  $F_n$  to occur, and given  $E_2$ , we need  $n - 2$  further tosses that finish  $TT$  for  $F_n$  to occur, with all tosses independent. Hence, if  $p_n = P(F_n)$  then  $p_2 = (1 - p)^2$ , otherwise, from (5),  $p_n$  satisfies

$$p_n = (p_{n-1} \times p) + (p_{n-2} \times (1 - p)) = pp_{n-1} + p(1 - p)p_{n-2}.$$

To find  $p_n$  explicitly, try a solution of the form  $p_n = A\lambda_1^n + B\lambda_2^n$  which gives

$$A\lambda_1^n + B\lambda_2^n = p(A\lambda_1^{n-1} + B\lambda_2^{n-1}) + p(1 - p)(A\lambda_1^{n-2} + B\lambda_2^{n-2}).$$



First, collecting terms in  $\lambda_1$  gives

$$\lambda_1^n = p\lambda_1^{n-1} + p(1-p)\lambda_1^{n-2} \implies \lambda_1^2 - p\lambda_1 - p(1-p) = 0,$$

indicating that  $\lambda_1$  and  $\lambda_2$  are given as the roots of this quadratic, that is

$$\lambda_1 = \frac{p - \sqrt{p^2 + 4p(1-p)}}{2}, \quad \lambda_2 = \frac{p + \sqrt{p^2 + 4p(1-p)}}{2}.$$

Furthermore,

$$\begin{aligned} n = 1 : p_1 = 0 & \implies A\lambda_1 + B\lambda_2 = 0 \\ n = 2 : p_2 = (1-p)^2 & \implies A\lambda_1^2 + B\lambda_2^2 = (1-p)^2 \end{aligned}$$

$$\implies A = \frac{(1-p)^2}{\lambda_1(\lambda_1 - \lambda_2)}, \quad B = -\frac{\lambda_1}{\lambda_2}A = -\frac{(1-p)^2}{\lambda_2(\lambda_1 - \lambda_2)}$$

$$\implies p_n = -\frac{(1-p)^2 \lambda_1^n}{\lambda_1(\lambda_2 - \lambda_1)} + \frac{(1-p)^2 \lambda_2^n}{\lambda_2(\lambda_2 - \lambda_1)} = \frac{(1-p)^2}{\sqrt{p(4-3p)}} (\lambda_2^{n-1} - \lambda_1^{n-1}), \quad n \geq 2.$$

**EXAMPLE 6** Information is transmitted digitally as a binary sequence known as “bits”. However, noise on the channel corrupts the signal, in that a digit transmitted as 0 is received as 1 with probability  $1 - \alpha$ , with a similar random corruption when the digit 1 is transmitted. It has been observed that, across a large number of transmitted signals, the 0s and 1s are transmitted in the ratio 3 : 4.

Given that the sequence 101 is received, what is the probability distribution over transmitted signals? Assume that the transmission and reception processes are independent

**SOLUTION:** Define:

- sample space  $\Omega$  to consist of all possible binary sequences of length three that is

$$\{000, 001, 010, 011, 100, 101, 110, 111\};$$

- a corresponding set of  
**signal** events  $\{S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7\}$  and  
**reception** events  $\{R_0, R_1, R_2, R_3, R_4, R_5, R_6, R_7\}$ .

We have observed event  $R_5$ , that 101 was received; we wish to compute  $P(S_i|R_5)$ , for  $i = 0, 1, \dots, 7$ . However, on the information given, we only have (or can compute)  $P(R_5|S_i)$ . First, using the Theorem of Total Probability, we compute  $P(R_5)$

$$P(R_5) = \sum_{i=0}^7 P(R_5|S_i)P(S_i). \quad (6)$$

Consider  $P(R_5|S_0)$ ; if 000 is transmitted, the probability that 101 is received is  $(1 - \alpha) \times \alpha \times (1 - \alpha) = \alpha(1 - \alpha)^2$  (corruption, no corruption, corruption) By complete evaluation we have

$$\begin{aligned} P(R_5|S_0) &= \alpha(1 - \alpha)^2, & P(R_5|S_1) &= \alpha^2(1 - \alpha), \\ P(R_5|S_2) &= (1 - \alpha)^3, & P(R_5|S_3) &= \alpha(1 - \alpha)^2, \\ P(R_5|S_4) &= \alpha^2(1 - \alpha), & P(R_5|S_5) &= \alpha^3, \\ P(R_5|S_6) &= \alpha(1 - \alpha)^2, & P(R_5|S_7) &= \alpha^2(1 - \alpha). \end{aligned}$$

Now, the prior information about digits transmitted is that the probability of transmitting a 1 is  $4/7$ , so

$$\begin{aligned} P(S_0) &= \left(\frac{3}{7}\right)^3, & P(S_1) &= \left(\frac{4}{7}\right) \left(\frac{3}{7}\right)^2, \\ P(S_2) &= \left(\frac{4}{7}\right) \left(\frac{3}{7}\right)^2, & P(S_3) &= \left(\frac{4}{7}\right)^2 \left(\frac{3}{7}\right), \\ P(S_4) &= \left(\frac{4}{7}\right) \left(\frac{3}{7}\right)^2, & P(S_5) &= \left(\frac{4}{7}\right)^2 \left(\frac{3}{7}\right), \\ P(S_6) &= \left(\frac{4}{7}\right)^2 \left(\frac{3}{7}\right), & P(S_7) &= \left(\frac{4}{7}\right)^3, \end{aligned}$$

and hence (6) can be computed as

$$P(R_5) = \frac{48\alpha^3 + 136\alpha^2(1-\alpha) + 123\alpha(1-\alpha)^2 + 36(1-\alpha)^3}{343}.$$

Finally, using Bayes' Theorem, we have the probability distribution over

$$\{S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7\}$$


given by

$$P(S_i|R_5) = \frac{P(R_5|S_i)P(S_i)}{P(R_5)}.$$

For example, the probability of correct reception is

$$P(S_5|R_5) = \frac{P(R_5|S_5)P(S_5)}{P(R_5)} = \frac{48\alpha^3}{48\alpha^3 + 136\alpha^2(1-\alpha) + 123\alpha(1-\alpha)^2 + 36(1-\alpha)^3}.$$

Posterior probability as a function of  $\alpha$ :



**EXAMPLE 7** While watching a game of Champions League football in a cafe, you observe someone who is clearly supporting Manchester United in the game.

What is the probability that they were actually born within 25 miles of Manchester ? Assume that:

- the probability that a randomly selected person in a typical local bar environment is born within 25 miles of Manchester is  $\frac{1}{20}$ , and;
- the chance that a person born within 25 miles of Manchester actually supports United is  $\frac{7}{10}$ ;
- the probability that a person not born within 25 miles of Manchester supports United with probability  $\frac{1}{10}$ .

**SOLUTION:** Define

- $B$  - event that the person is born within 25 miles of Manchester
- $U$  - event that the person supports United.

We want  $P(B|U)$ . By Bayes' Theorem,

$$\begin{aligned} P(B|U) &= \frac{P(U|B)P(B)}{P(U)} = \frac{P(U|B)P(B)}{P(U|B)P(B) + P(U|B')P(B')} \\ &= \frac{\frac{7}{10} \frac{1}{20}}{\frac{7}{10} \frac{1}{20} + \frac{1}{10} \frac{19}{20}} \\ &= \frac{7}{26} \approx 0.269. \end{aligned}$$