

THE DECIDABLE AND THE UNDECIDABLE. A SURVEY OF RECENT RESULTS

Søren Brinck Knudstorp
NihiL Workshop

February 5, 2024

University of Amsterdam

Plan for the talk

- (Un)decidability: what and why?
- Propositional team logics and their decidability
- Exploring boundaries between the decidable and the undecidable
 - Solving problems and obtaining insights along the way
 - Using insights to solve one last problem

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why?

(Un)decidability: what and why?

What?

A **decision problem** is a collection of inputs I , with a yes-or-no question for each $i \in I$.

A decision problem is **decidable** if there is an effective method that, given any $i \in I$, accurately answers the question. Otherwise, it is **undecidable**.

A logic \mathbf{L} , in a language \mathcal{L} , is decidable if there is an effective method that, given any $\varphi \in \mathcal{L}$, determines whether $\mathbf{L} \vdash \varphi$. Otherwise, it is undecidable.

Why? *Because it is a deep, profound and significant conceptual distinction.*

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ("teams") of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \vee \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ("teams") of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \vee \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ('teams') of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \vee \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ('teams') of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \cup \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ('teams') of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \vee \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Propositional team logics and their decidability

Traditionally (in, e.g., CPC), formulas φ are evaluated at **single valuations**

$v : \mathbf{Prop} \rightarrow \{0, 1\}$,

$$v \models \varphi.$$

In team semantics, formulas φ are evaluated at **sets ('teams') of valuations**

$s \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$,

$$s \models \varphi.$$

Definition (some team-semantic clauses)

Let $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$. For $s \in \mathcal{P}(X)$, we define

$s \models p$	iff	$\forall v \in s : v(p) = 1$,
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi$,
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi$,
$s \models \sim \varphi$	iff	$s \not\models \varphi$,
$s \models \varphi \cup \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

Observation. All propositional team logics are decidable: given φ , simply check whether $s \models \varphi$ for all $s \subseteq \{v \mid v : \mathbf{Prop}(\varphi) \rightarrow \{0, 1\}\}$.

Yet, this explanation is hardly satisfactory.
What is it that makes propositional team logics
decidable, *and others not?*

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \cup \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a powerset frame $\mathbb{F} = (\mathcal{P}(X), \cup)$, where 'o' is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$;

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi \text{ and } s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi \text{ or } s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi \text{ and } s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi \text{ or } s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi$; $s'' \models \psi$; and $s = s' \cup s''$.

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Team semantics as relational semantics

Recall our semantic clauses: For $X := \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ and $s \in \mathcal{P}(X)$, we had

$s \models p$	iff	$\forall v \in s : v(p) = 1,$
$s \models \varphi \wedge \psi$	iff	$s \models \varphi$ and $s \models \psi,$
$s \models \varphi \vee \psi$	iff	$s \models \varphi$ or $s \models \psi,$
$s \models \sim \varphi$	iff	$s \not\models \varphi,$
$s \models \varphi \circ \psi$	iff	there exist $s', s'' \in \mathcal{P}(X)$ such that $s' \models \varphi;$ $s'' \models \psi;$ and $s = s' \cup s''.$

This induces a **powerset frame** $\mathbb{F} = (\mathcal{P}(X), \cup)$, where ‘ \circ ’ is a binary modality referring to the ternary \cup -relation $s = s' \cup s''$; and a **model** $\mathbb{M} = (\mathcal{P}(X), \cup, V)$ with a ‘principal valuation’, i.e.,

$$V(p) := \{s \in \mathcal{P}(X) \mid \forall v \in s : v(p) = 1\} = \downarrow\{v \in X \mid v(p) = 1\}.$$

In fact, if we take all powerset frames $(\mathcal{P}(X), \cup)$, redefine the base clause

$$(\mathcal{P}(X), \cup, V), s \Vdash p \quad \text{iff} \quad s \in V(p),$$

and only allow principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$, we get **sound and complete relational semantics for team logics**.

Proof. A simple p-morphism argument.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: *Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for **arbitrary** valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?*

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for arbitrary valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: *Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for **arbitrary** valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?*

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: *Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for **arbitrary** valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?*

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: *Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for **arbitrary** valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?*

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Powerset frames and Boolean frames

Summarizing, (i) team logics are decidable, and (ii) relational semantics for team logics are given by powerset frames $(\mathcal{P}(X), \cup)$ with principal valuations $V : \mathbf{Prop} \rightarrow \{\downarrow s \mid s \in \mathcal{P}(X)\}$.

Question: *Sticking with the signature $\{\wedge, \vee, \sim, \circ\}$, what happens if we allow for **arbitrary** valuations $V : \mathbf{Prop} \rightarrow \mathcal{PP}(X)$? Does the logic remain decidable?*

In fact, this question is intimately related with an open problem: Goranko and Vakarelov (1999) consider the logic of Boolean frames – instead of a powerset $\mathcal{P}(X)$, the carrier is a Boolean algebra B – and raises the problem of its decidability.¹

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

¹Goranko and Vakarelov (1999) call their logic ‘hyperboolean modal logic’ and include modalities for all the Boolean operations, not just the join.

Proof method: tiling

- A (Wang) tile is a square with colors on each side.
- The tiling problem: given any finite set of tiles \mathcal{W} , determine whether each point in the quadrant \mathbb{N}^2 can be assigned a tile from \mathcal{W} such that neighboring tiles share matching colors on connecting sides.
- The tiling problem was introduced by Wang (1963) and proven undecidable by Berger (1966).




Figure 1: Wang tiles

Figures taken from: https://en.wikipedia.org/wiki/Wang_tile




Figure 2: A tiling of the plane

Proof method: tiling

- A (Wang) tile is a square with colors on each side.
- The tiling problem: given any finite set of tiles \mathcal{W} , determine whether each point in the quadrant \mathbb{N}^2 can be assigned a tile from \mathcal{W} such that neighboring tiles share matching colors on connecting sides.
- The tiling problem was introduced by Wang (1963) and proven undecidable by Berger (1966).




Figure 1: Wang tiles

Figures taken from: https://en.wikipedia.org/wiki/Wang_tile




Figure 2: A tiling of the plane

Proof method: tiling

- A (Wang) tile is a square with colors on each side.
- The tiling problem: given any finite set of tiles \mathcal{W} , determine whether each point in the quadrant \mathbb{N}^2 can be assigned a tile from \mathcal{W} such that neighboring tiles share matching colors on connecting sides.
- The tiling problem was introduced by Wang (1963) and proven undecidable by Berger (1966).




Figure 1: Wang tiles

Figures taken from: https://en.wikipedia.org/wiki/Wang_tile




Figure 2: A tiling of the plane

Proof method: tiling

- A (Wang) tile is a square with colors on each side.
- The tiling problem: given any finite set of tiles \mathcal{W} , determine whether each point in the quadrant \mathbb{N}^2 can be assigned a tile from \mathcal{W} such that neighboring tiles share matching colors on connecting sides.
- The tiling problem was introduced by Wang (1963) and proven **undecidable** by Berger (1966).




Figure 1: Wang tiles

Figures taken from: https://en.wikipedia.org/wiki/Wang_tile




Figure 2: A tiling of the plane

Proof method: tiling

- A (Wang) tile is a square with colors on each side.
- The tiling problem: given any finite set of tiles \mathcal{W} , determine whether each point in the quadrant \mathbb{N}^2 can be assigned a tile from \mathcal{W} such that neighboring tiles share matching colors on connecting sides.
- The tiling problem was introduced by Wang (1963) and proven **undecidable** by Berger (1966).




Figure 1: Wang tiles

Figures taken from: https://en.wikipedia.org/wiki/Wang_tile




Figure 2: A tiling of the plane

Proof method: tiling

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

Proof idea.

For each finite set of tiles \mathcal{W} , we construct a formula $\phi_{\mathcal{W}}$ such that \mathcal{W} tiles the quadrant if and only if $\phi_{\mathcal{W}}$ is satisfiable. \square

Dividing the proof into two lemmas, corresponding to a direction each, we can prove both results in one go:

Lemma

If $\phi_{\mathcal{W}}$ is satisfiable (in a Boolean frame), then \mathcal{W} tiles \mathbb{N}^2 .

Lemma

If \mathcal{W} tiles \mathbb{N}^2 , then $\phi_{\mathcal{W}}$ is satisfiable (in $(\mathcal{P}(\mathbb{N}), \cup)$).

Proof method: tiling

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

Proof idea.

For each finite set of tiles \mathcal{W} , we construct a formula $\phi_{\mathcal{W}}$ such that \mathcal{W} tiles the quadrant if and only if $\phi_{\mathcal{W}}$ is satisfiable. \square

Dividing the proof into two lemmas, corresponding to a direction each, we can prove both results in one go:

Lemma

If $\phi_{\mathcal{W}}$ is satisfiable (in a Boolean frame), then \mathcal{W} tiles \mathbb{N}^2 .

Lemma

If \mathcal{W} tiles \mathbb{N}^2 , then $\phi_{\mathcal{W}}$ is satisfiable (in $(\mathcal{P}(\mathbb{N}), \cup)$).

Proof method: tiling

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

Proof idea.

For each finite set of tiles \mathcal{W} , we construct a formula $\phi_{\mathcal{W}}$ such that \mathcal{W} tiles the quadrant if and only if $\phi_{\mathcal{W}}$ is satisfiable. \square

Dividing the proof into two lemmas, corresponding to a direction each, we can prove both results in one go:

Lemma

If $\phi_{\mathcal{W}}$ is satisfiable (in a Boolean frame), then \mathcal{W} tiles \mathbb{N}^2 .

Lemma

If \mathcal{W} tiles \mathbb{N}^2 , then $\phi_{\mathcal{W}}$ is satisfiable (in $(\mathcal{P}(\mathbb{N}), \cup)$).

Proof method: tiling

Theorem

The logic of powerset frames, in the signature $\{\wedge, \vee, \sim, \circ\}$, with arbitrary valuations is *undecidable*. And so is the hyperboolean modal logic of Goranko and Vakarelov (1999).

Proof idea.

For each finite set of tiles \mathcal{W} , we construct a formula $\phi_{\mathcal{W}}$ such that \mathcal{W} tiles the quadrant if and only if $\phi_{\mathcal{W}}$ is satisfiable. \square

Dividing the proof into two lemmas, corresponding to a direction each, we can prove both results in one go:

Lemma

If $\phi_{\mathcal{W}}$ is satisfiable (in a Boolean frame), then \mathcal{W} tiles \mathbb{N}^2 .

Lemma

If \mathcal{W} tiles \mathbb{N}^2 , then $\phi_{\mathcal{W}}$ is satisfiable (in $(\mathcal{P}(\mathbb{N}), \cup)$).

Insight 1: valuations matter

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Semilattice frames, associativity and negation

Question: *Since we can weaken from powersets to Boolean algebras and stay undecidable, how much further can we go while remaining undecidable?*

Weakening from powersets $(\mathcal{P}(X), \cup)$ to general (join-)semilattices (S, \sqcup) , we get a the problem posed by Bergman (2018) and Jipsen et al. (2021) (and SBK (2023a)).

Theorem

For any class \mathcal{S} of semilattices containing $(\mathcal{P}(\mathbb{N}), \cup)$, its logic in the signature $\{\wedge, \vee, \sim, \circ\}$, is undecidable.

Question: *What if we weaken even further than semilattices?*

(Partial) answer: As semilattices are partial orders ' \leq ' with all binary suprema, we could consider the logic of all *partial orders simpliciter*. This is modal information logic, which is proven **decidable** in SBK (2023b).

Question: *What if we, instead, reduce our signature $\{\wedge, \vee, \sim, \circ\}$?*

Answer: If we stick to semilattices but *omit negation*, so signature is $\{\wedge, \vee, \circ\}$, we obtain *Finean truthmaker semantics*, proven **decidable** in SBK (2023a).

Insight 2: associativity matters

Insight 3: negation matters

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqsubseteq, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting' **undecidability**]
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting' **undecidability**]
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**']
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**']

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting' **undecidability**]
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

(Un)decidability of relevant \mathbf{S} : using our insights

Problem of concern: *Is relevant logic \mathbf{S} decidable?*

\mathbf{S} is the logic of semilattice frames $(S, \sqcup, \mathbf{0})$ with a bottom element $\mathbf{0}$, with arbitrary valuations, in the signature $\{\wedge, \vee, \rightarrow\}$. ' \rightarrow ' is closely connected to ' \circ ' (it is its residual).

What we know about the problem:

- Omitting disjunction, the logic $\mathbf{S}_{\wedge, \rightarrow}$ is **decidable**.
- If we restrict to hereditary valuations, we obtain positive intuitionistic logic, which is **decidable**.
- \mathbf{S} is closely connected to positive relevant \mathbf{R}^+ , which is **undecidable**.
 - Und. of \mathbf{R}^+ was shown by Urquhart (1984), but \mathbf{S} eluded these techniques.
 - Eventually, this led Urquhart (2016) to conjecture that \mathbf{S} is **decidable**.

What we notice about the problem:

- *Valuations are arbitrary*, contra positive intuitionistic logic. ['suggesting **undecidability**']
- \mathbf{S} is positive, *no negation!* [suggesting **decidability**]
- Frames of \mathbf{S} are semilattices, *they are associative!* [suggesting **undecidability**]

Theorem: S is undecidable



Berger, R. (1966). *The undecidability of the domino problem*. English. Vol. 66. Mem. Am. Math. Soc. Providence, RI: American Mathematical Society (AMS). DOI: [10.1090/memo/0066](https://doi.org/10.1090/memo/0066) (cit. on pp. 33–37).







Bergman, C. (2018). “Introducing Boolean Semilattices”. In: *Don Pigozzi on Abstract Algebraic Logic, Universal Algebra, and Computer Science. Outstanding Contributions to Logic*. Ed. by J. Czelakowski. Springer, pp. 103–130 (cit. on pp. 43–54).



Goranko, V. and D. Vakarelov (1999). “Hyperboolean Algebras and Hyperboolean Modal Logic”. In: *Journal of Applied Non-Classical Logics* 9.2-3, pp. 345–368. DOI: [10.1080/11663081.1999.10510971](https://doi.org/10.1080/11663081.1999.10510971) (cit. on pp. 27–32, 38–41).

References II

-  Jipsen, P., M. Eyad Kurd-Misto, and J. Wimberley (2021). **“On the Representation of Boolean Magmas and Boolean Semilattices”**. In: *Hajnal Andr  ka and Istv  n N  meti on Unity of Science: From Computing to Relativity Theory Through Algebraic Logic*. Ed. by J. Madar  sz and G. Sz  kely. Cham: Springer International Publishing, pp. 289–312. DOI: [10.1007/978-3-030-64187-0_12](https://doi.org/10.1007/978-3-030-64187-0_12) (cit. on pp. 43–54).
-  Knudstorp, S. B. (2023a). **“Logics of Truthmaker Semantics: Comparison, Compactness and Decidability”**. In: *Synthese* (cit. on pp. 43–54).
-  — (2023b). **“Modal Information Logics: Axiomatizations and Decidability”**. In: *Journal of Philosophical Logic* (cit. on pp. 43–54).
-  Urquhart, A. (1984). **“The undecidability of entailment and relevant implication”**. In: *Journal of Symbolic Logic* 49, pp. 1059–1073 (cit. on pp. 57–74).



Urquhart, A. (2016). **“Relevance Logic: Problems Open and Closed”**. In: *The Australasian Journal of Logic* 13 (cit. on pp. 57–74).



Wang, H. (1963). **“Dominoes and the $\forall\exists\forall$ case of the decision problem”**. In: *Mathematical Theory of Automata*, pp. 23–55 (cit. on pp. 33–37).

Thank you!