


How (Not) to Probe The Axion-Electron Coupling

Kevin Zhou




UChicago Particle Theory Seminar — April 16, 2025
arXiv:2312.11601, with Asher Berlin, Alex Millar, Tanner Trickle



Ultralight dark matter is a subject of growing experimental interest:

- Explains apparent abundance of dark matter with simple production mechanisms
- Generic: weakly coupled ultralight fields appear in many models
- Low-hanging fruit: new small-scale experiments are needed, and very effective
- Minimal: requires introduction of only a single new field at low energies
- Bounded: only a few interactions are natural and leading in effective field theory

Couplings of a Field Theory Axion




arises from spontaneous breaking of chiral symmetry at high energies

axion is Goldstone boson, so recast most interactions in derivative form by **chiral field redefinitions**, inducing couplings to gauge bosons by anomaly

take **low-energy limit** to find effects on nucleons

Couplings of a Field Theory Axion



arises from spontaneous breaking of chiral symmetry at high energies

axion is Goldstone boson, so recast most interactions in derivative form by chiral field redefinitions, inducing couplings to gauge bosons by anomaly

take low-energy limit to find effects on nucleons

only a few dimension 5 operators, and all derivative couplings generically arise

$$(\partial_\mu a) K_{\text{EM}}^\mu$$


axion-photon

$$(\partial_\mu a) \bar{N} \gamma^\mu \gamma^5 N$$

axion-nucleon

$$(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$

axion-electron



$$(\partial_\mu a) K_{\text{EM}}^\mu$$



axion-photon

$$(\partial_\mu a) \bar{N} \gamma^\mu \gamma^5 N$$

axion-nucleon

$$(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$

axion-electron



Less explored, but several concepts could probe well beyond astrophysical bounds

Generally achieved by resonantly amplifying a nuclear spin torque $\tau = g_{aN} \hat{s} \times \nabla a$

Enhanced by the exceptional stability ($Q_{\text{eff}} \gtrsim 10^{10}$) of nuclear spin precession

$$(\partial_\mu a) K_{\text{EM}}^\mu$$


axion-photon

$$(\partial_\mu a) \bar{N} \gamma^\mu \gamma^5 N$$

axion-nucleon

$$(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$

axion-electron



No active experiments search for this coupling,
and even ambitious proposals have difficulty beating astrophysical bounds!

$$(\partial_\mu a) K_{\text{EM}}^\mu$$

axion-photon

$$(\partial_\mu a) \bar{N} \gamma^\mu \gamma^5 N$$

axion-nucleon

$$(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$

axion-electron

This talk explains:

- What this coupling does
- Why it's difficult to probe
- Why 8 previous claims of strong sensitivity were all wrong
- How it's still possible to use it to probe an meV QCD axion

The Axion-Electron Coupling

Lagrangian has axial vector current:

$$\mathcal{L} \supset g(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$

Which is a classical particle's spin 4-vector:


$$\int d^3x \bar{e} \gamma^\mu \gamma^5 e \rightarrow s^\mu \simeq (\mathbf{v} \cdot \hat{\mathbf{s}}, \hat{\mathbf{s}})^\mu$$

So for a nonrelativistic quantum particle:

$$H \supset -g(\nabla a) \cdot \sigma - \frac{g}{m_e} \dot{a} \sigma \cdot (\mathbf{p} - q\mathbf{A})$$

“axion wind” spin torque

$$\tau = g \hat{\mathbf{s}} \times \nabla a$$



“axioelectric” term

both terms oscillate with the axion field, at $f \sim (m_a/\mu\text{eV})$ GHz

$$|\nabla a| \sim v_{\text{DM}} \dot{a} \text{ where } v_{\text{DM}} \sim 10^{-3}$$

axioelectric term seems larger, but subject to traps!

Axioelectric Pitfalls

For simplicity, set $\nabla a = \mathbf{B} = 0$ and work in gauge $\mathbf{E} = -\dot{\mathbf{A}}$

$$H \simeq \frac{(\mathbf{p} - q\mathbf{A})^2}{2m_e} - \frac{g}{m_e} \dot{a} \sigma \cdot (\mathbf{p} - q\mathbf{A})$$

Idea #1: this implies $\mathcal{H} \supset -\mathbf{J}_{\text{eff}} \cdot \mathbf{A}$ where $\mathbf{J}_{\text{eff}} = (g\dot{a}q/m_e) \langle n_e \sigma \rangle$, a large current!

However, it would be more transparent to write: $H \simeq \frac{(\mathbf{p} - q\mathbf{A} - g\dot{a}\sigma)^2}{2m_e}$

No extra contribution to the current, but a new force: $\mathbf{A}_{\text{eff}} = (g/q) \dot{a}\sigma$ $\mathbf{F} = -g \frac{d}{dt} \langle \dot{a}\sigma \rangle$

Generally very small, because it is suppressed by two time derivatives

Idea #2: integrating the force over time yields $\mathbf{J} = (g\dot{a}q/m_e) \langle n_e \sigma \rangle$, a large current!

(For DFSZ axion, would be $\sim 10^3$ larger than \mathbf{J}_{eff} from axion-photon coupling)

Estimating the Axioelectric Current

How large of a current can we really get from $\mathbf{E}_{\text{eff}} = (g/q) \ddot{a} \sigma$?

In a spin-polarized dielectric, we have: $\mathbf{J}_a = (\epsilon - 1) \dot{\mathbf{E}}_{\text{eff}}$

But material also creates electric field: ~~$\nabla \times \nabla \times \mathbf{E} + \epsilon \ddot{\mathbf{E}} = -\dot{\mathbf{J}}_a$~~

Total polarization current:


$$\mathbf{J}_{\text{tot}} = (\epsilon - 1)(\dot{\mathbf{E}} + \dot{\mathbf{E}}_{\text{eff}}) = \frac{\epsilon - 1}{\epsilon} \dot{\mathbf{E}}_{\text{eff}}$$

Result suppressed by **three** time derivatives! But what if charges can move?


$$\epsilon(\omega) \sim \begin{cases} 1 + i\sigma/\omega & \text{ideal conductor} \\ 1 - \omega_p^2/\omega^2 & \text{ideal superconductor} \end{cases}$$

Above result holds, and implies $\mathbf{J}_{\text{tot}} \sim \dot{\mathbf{E}}_{\text{eff}}$ even when $|\epsilon| \gg 1$!


Physically, better conductors shield \mathbf{E}_{eff} more



Axioelectric Reach vs. Other Projections




Since $\mathbf{J} \propto d^3a/dt^3$, best probes are at high m_a , and still not very strong...



...but many recent papers claimed **vastly** stronger effects proportional to a !

The EDM Controversy

The axion-electron coupling has two equivalent forms



which are related by redefining states, $|\psi'\rangle = U|\psi\rangle$, so that all observables same

$$U \simeq \exp \left(-\frac{iga}{2m} (\mathbf{p} - q\mathbf{A}) \cdot \boldsymbol{\sigma} \right)$$

Already noted for pion-nucleon interactions in 1950s – but not obvious how!

Coefficient of the Axioelectric Term

For simplicity take $q = 0$, where only difference is axioelectric coefficient:

$$H_d \supset -\frac{g\dot{a}}{m} \boldsymbol{\sigma} \cdot \mathbf{p}$$

$$H_{nd} \supset -\frac{g\dot{a}}{2m} \boldsymbol{\sigma} \cdot \mathbf{p}$$

Puzzle: why can the coefficient of a force be freely adjusted?

Resolution: **any** force can be removed from a single particle Hamiltonian by working in the particle's reference frame; only relative accelerations physical

$$H_d \supset V(\mathbf{x}_1 - \mathbf{x}_2) - \frac{g\dot{a}}{m} (\boldsymbol{\sigma}_1 \cdot \mathbf{p}_1 + \boldsymbol{\sigma}_2 \cdot \mathbf{p}_2)$$

$$H_{nd} \supset V(\mathbf{x}_1 - \mathbf{x}_2) - \frac{g\dot{a}}{2m} (\boldsymbol{\sigma}_1 \cdot \mathbf{p}_1 + \boldsymbol{\sigma}_2 \cdot \mathbf{p}_2) - \frac{ga}{2m} (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \nabla_1 V(\mathbf{x}_1 - \mathbf{x}_2)$$


Shifting the axioelectric coefficient adds other terms with same physical effect

True and Spurious EDMs

The new term is the nonrelativistic limit of a genuine EDM:

$$H_{\text{nd}} \supset \frac{qga}{2m} \boldsymbol{\sigma} \cdot \mathbf{E} \quad H_{\text{EDM}} = \frac{d}{2} \Psi \gamma^5 \sigma^{\mu\nu} \Psi F_{\mu\nu} = -d \boldsymbol{\sigma} \cdot \mathbf{E} + (\text{relativistic corrections})$$

But for charged nonrelativistic particles, a constant EDM has no physical effects!



In this limit, just equivalent to unobservable shift in definition of position!
(key intuition behind “Schiff screening”)

True and Spurious EDMs

The new term is the nonrelativistic limit of a genuine EDM:

$$H_{\text{nd}} \supset \frac{qga}{2m} \boldsymbol{\sigma} \cdot \mathbf{E} \quad H_{\text{EDM}} = \frac{d}{2} \Psi \gamma^5 \sigma^{\mu\nu} \Psi F_{\mu\nu} = -d \boldsymbol{\sigma} \cdot \mathbf{E} + (\text{relativistic corrections})$$

But for charged nonrelativistic particles, a constant EDM has no physical effects!

The effects of a true EDM probed in experiments are from relativistic corrections:

$O(v/c)$ magnetic dipole, $O(v^2/c^2)$ length contraction

Commins, Jackson, DeMille (2007)


so H_{nd} contains **only** the part of H_{EDM} with no physical effect!

(time-varying $a(t)$ does have effect, but highly suppressed by $\sim (m_a/\text{eV})^2$)

Stadnik and Flambaum (2014)

Probing the Axion Wind Torque

The axion wind spin torque tilts spins, yielding oscillating transverse magnetic field



Naively seems easier for electrons, as $\mu_e/\mu_n \sim m_p/m_e \sim 10^3$
But typical quality factors much lower!

Solid magnetic material


Naturally probes $m_a \sim \text{meV}$ but interactions prevent very high Q

Signal is emitted meV radiation

"Separated" or "locked" electrons

Much higher Q and lower resonant frequencies possible

Signal is quasistatic magnetic field, detected by magnetometer




Separated electrons: comagnetometer vapor, storage ring bunch, NV centers
 (but much lower number density results in limited sensitivity)

Locked electrons: align spins to crystal, so spin torque becomes mechanical
 (but suppressed by inertia of nuclei, doesn't benefit from electron being light)

Probing the Axion Wind Torque

The axion wind spin torque tilts spins, yielding oscillating transverse magnetic field



Naively seems easier for electrons, as $\mu_e/\mu_n \sim m_p/m_e \sim 10^3$
But typical quality factors much lower!

Solid magnetic material



Naturally probes $m_a \sim \text{meV}$ but interactions prevent very high Q

Signal is emitted meV radiation

“Separated” or “locked” electrons


Much higher Q and lower resonant frequencies possible

Signal is quasistatic magnetic field, detected by magnetometer



Before our work, all existing and proposed experiments used the “ferromagnetic haloscope” concept

The Ferromagnetic Haloscope



effective magnetic field


$$\mathbf{B}_{\text{eff}} = \frac{g_{ae}}{\mu_B} \nabla a$$

magnetic quality factor

$$Q_m \sim 10^4$$

sample magnetization

$$\omega_m = 2\mu_B M_0 \sim 0.03 \text{ meV}$$



$$V \sim \begin{cases} 0.005 \text{ cm}^3 & \text{QUAX (2018)} \\ 0.01 \text{ cm}^3 & \text{Flower et al. (2019)} \\ 0.1 \text{ cm}^3 & \text{QUAX (2020)} \\ 100 \text{ cm}^3 & \text{QUAX target} \end{cases}$$

Main issue: scaling sample volume

Single crystal YIG made very slowly, in mm spheres, at cost of over \$10⁷/kg

The Ferromagnetic Haloscope

High Field Magnet

TEM102
Cavity

Alternative idea: replace strong resonant enhancement with overall scale

Ferromagnet spin precesses on resonance
with axion, driving a cavity mode

Polycrystalline spinel ferrites are cheap and mass produced



Ferrite Blocks Ceramic Magnets, 1 7/8" x 7/8" x 3/8" Rectangular Magnets, Ceramic Rectangular Square Magnets, Grade-8 Hard Ferrite Magnets for Crafts, Science and Hobbies (8 Pieces)

★★★★★ ~33

\$14⁹⁸

✓prime One-Day
FREE delivery Tomorrow, Nov 18


Lower carbon delivery ~

Add to Cart

M_0 double that of YIG

$$Q_m \sim 10^2$$

$$\omega_m = 2\mu_B M_0 \sim 0.03 \text{ meV}$$




$$V \sim \begin{cases} 0.005 \text{ cm}^3 & \text{QUAX (2018)} \\ 0.01 \text{ cm}^3 & \text{Flower et al. (2019)} \\ 0.1 \text{ cm}^3 & \text{QUAX (2020)} \\ 100 \text{ cm}^3 & \text{QUAX target} \end{cases}$$

Main issue: scaling sample volume

Single crystal YIG made very slowly, in mm spheres, at cost of over \$10⁷/kg

Radiation From a Slab



Solving spin equation of motion:


$$\mathbf{J}_a = \nabla \times ((1 - \mu^{-1}) \mathbf{B}_{\text{eff}})$$

Magnetized slab carries surface currents

$$1 - \mu^{-1} \simeq \frac{M_0}{B_0 - (m_a/2\mu_B) - iB_0/2Q_m}$$

(for clockwise polarization)

Radiation From a Slab



much of this power is converted to outgoing radiation

$$\mathbf{J}_a = \nabla \times ((1 - \mu^{-1}) \mathbf{B}_{\text{eff}})$$


Magnetized slab carries surface currents

$$1 - \mu^{-1} \simeq \frac{M_0}{B_0 - (m_a/2\mu_B) - iB_0/2Q_m}$$

Power deposited in slab maximized on resonance, at tunable frequency

$$(B_0 \leq 10 \text{ T} \text{ implies } m_a \leq 10^{-3} \text{ eV})$$

Radiation From a Slab



$$\mathbf{J}_a = \nabla \times ((1 - \mu^{-1}) \mathbf{B}_{\text{eff}})$$

Magnetized slab carries surface currents


$$1 - \mu^{-1} \simeq \frac{M_0}{B_0 - (m_a/2\mu_B) - iB_0/2Q_m}$$

Power deposited in slab maximized on resonance, at tunable frequency

$$(B_0 \leq 10 \text{ T} \text{ implies } m_a \leq 10^{-3} \text{ eV})$$

Place layers, with tunable separation, so emitted radiation interferes constructively

Magnetized Multilayers



Like the MADMAX experiment, radiation emitted from layers is focused on a detector

For simplicity, we consider the “transparent mode” setup:

slab spacing π/m_a , slab thickness $\pi/(\text{Re}(n) m_a)$

$$P_{\text{sig}} \sim \mathcal{F}^2 N^2 |B_{\text{eff}}|^2 A$$

slab form factor

$$\mathcal{F} \sim 1 - \mu^{-1}$$

(+ material loss effects)

slab area


$$A = 1 \text{ m}^2$$

number of layers

$$N \leq 80$$


$$N \leq \text{Re}(n)/(\pi \text{Im}(n))$$

Optimizing a Magnetized Multilayer



A few thick slabs tuned close to resonance; large volume needed, but only low magnetic field

Optimizing a Magnetized Multilayer



A few thick slabs tuned close to resonance; large volume needed, but only low magnetic field

Many thin slabs, which must be well off resonance to avoid absorption; high field but only in small volume

Benchmark Reach


$$\text{SNR} \simeq \frac{P_{\text{sig}}}{T_n} \sqrt{\frac{t_{\text{int}}}{\Delta\nu_a}}$$

Standard HEMT amplifier



Helium cryostat cooling

$$T_n = 4 \text{ K} + T_{\text{amp}}$$



Benchmark Reach


$$\text{SNR} \simeq \frac{P_{\text{sig}}}{T_n} \sqrt{\frac{t_{\text{int}}}{\Delta\nu_a}}$$

Standard quantum limited amplification

Dilution fridge cooling

$$T_n = 40 \text{ mK} + T_{\text{amp}}$$

More challenging, but realized in several expts




Benchmark Reach

$$\text{SNR} \simeq \frac{P_{\text{sig}} \sqrt{t_{\text{int}}}}{\text{NEP}}$$

Frequency insensitive
bolometer

$$\text{NEP} = 10^{-22} \text{ W}/\sqrt{\text{Hz}}$$

About 10x improvement
over state of the art;
target for BREAD



Benchmark Reach


$$\text{SNR} \simeq \frac{P_{\text{sig}}}{m_a} \sqrt{\frac{t_{\text{int}}}{\text{DCR}}}$$

Thermal limited
photon counting



$$\text{DCR} \sim \frac{\Delta\nu_s}{e^{m_a/T} - 1}$$

Challenging, but some
prototypes at meV

Ferromagnetic haloscopes
already assume noise-free
photon counting!



Open Questions



Conclusion

- The axion-electron coupling is minimal and generic
- Its physical effects are simple, but can be subtle to compute
- Recent attempts to probe it all used a single concept...
- ...but new ideas may allow scaling to stronger sensitivity

$$\mathcal{L} \supset g(\partial_\mu a) \bar{e} \gamma^\mu \gamma^5 e$$
$$H \supset -g(\nabla a) \cdot \sigma - \frac{g}{m_e} \dot{a} \sigma \cdot (\mathbf{p} - q\mathbf{A})$$

