

Math 341: Homework 4

Daniel Ko

Spring 2020

§1 A

Let V be a vector space having dimension n , and let S be a subset of V that generates V .

- a. Prove that there is a subset of S that is a basis for V . (Be careful not to assume that S is finite).

Proof.

Since V is finite dimensional, there exists a basis for V .

$$B = \{v_1, v_2, \dots, v_n\}$$

Any $v \in B$ can be expressed as a linear combination of S because $\text{span}(S) = V$.

Let the subset of S that generates v_i be S_i

$$v_i = \sum_{j=1}^{m_i} a_j^i s_j^i \text{ where } a \in F \text{ and } s \in S_i$$

The span of the union of the sets that generates v , $\text{span}(\bigcup_{i=1}^n S_i) = V$

Corollary 2(a) of Theorem 1.10 states that a generating set for V that contains exactly n vectors is a basis for V . The set above, which is a subset of S , contains exactly n vectors and generates V . Therefore, there is subset of S that is a basis for V . \square

- b. Prove that S contains at least n vectors.

Proof.

From (a) we know there is a subset of S that forms a basis. Since that subset contains n vectors, S must contain n or more vectors. \square

§2 B

Let $f(x)$ be a polynomial of degree n in $P_n(R)$. Prove that for any $g(x) \in P_n(R)$ there exists scalars c_0, c_1, \dots, c_n such that

$$g(x) = c_0 f(x) + c_1 f'(x) + c_2 f''(x) + \dots + c_n f^{(n)}(x)$$

Proof.

Let $B = \{f, f', f'', \dots, f^{(n)}\}$.

If B forms a basis we can express any $g(x) \in P_n(R)$ in the format above (a linear combination).

We can determine if B is basis by seeing if it is linearly independent by using a matrix.

$$\mu_0 f + \mu_1 f' + \mu_2 f'' + \dots + \mu_n f^{(n)} = 0$$

$$\begin{bmatrix} a_n & a_{n-1} & \cdots & \cdots & a_0 \\ & na_n & \cdots & \cdots & a_1 \\ & & \ddots & \cdots & \vdots \\ & & & \ddots & \vdots \\ & & & & n!a_n & (n-1)!a_{n-1} \\ & & & & & n!a_n \end{bmatrix} \begin{bmatrix} \mu_0 \\ \mu_1 \\ \vdots \\ \vdots \\ \mu_{n-1} \\ \mu_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ \vdots \\ 0 \\ 0 \end{bmatrix}$$

Solving this system of equations:

Looking at the bottom row, $n!a_n\mu_n = 0$

$\mu_n = \frac{0}{n!a_n} = 0$, a_n is non zero because f is a n th degree polynomial and a_n is its coefficient.

Looking at row $n-1$, $n!a_n\mu_{n-1} + (n-1)!a_{n-1}\mu_n = 0$

Because $\mu_n = 0$, $n!a_n\mu_{n-1} + 0 = 0$

$\mu_{n-1} = \frac{0}{n!a_n} = 0$

By back substitution, $\mu_n = \mu_{n-1} = \cdots = \mu_1 = \mu_0 = 0$

This means that B is linearly independent, which also means that B is a basis.

Therefore, any $g(x) \in P_n(R)$ can be a linear combination of B with the scalars c_0, c_1, \dots, c_n \square

§3 C

a. prove blah blah

Proof.

W_1 and W_2 are finite dimensional subspaces of $V \Rightarrow$ subspace $W_1 + W_2$ is finite dimensional and $\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2)$

Let $B_{1 \cap 2}$ be a basis for $W_1 \cap W_2$

$B_{1 \cap 2} = \{u_1, u_2, \dots, u_k\}$

By using the replacement theorem, we can extend $B_{1 \cap 2}$ to be a basis for W_1

So the basis for W_1 is B_1

$B_1 = \{u_1, u_2, \dots, u_k, v_1, v_2, \dots, v_m\}$

Likewise, we can extend $B_{1 \cap 2}$ to be a basis for W_2

$B_2 = \{u_1, u_2, \dots, u_k, w_1, w_2, \dots, w_p\}$

Basis for $W_1 + W_2$ will be $B_1 \cup B_2$, however they may contain the same vectors twice.

To prevent double counting, we must subtract $B_1 \cap B_2$ from $B_1 \cup B_2$

Thus the basis for $W_1 + W_2$ is

$B_{1+2} = \{u_1, u_2, \dots, u_k, v_1, v_2, \dots, v_m, w_1, w_2, \dots, w_p\}$

$W_1 + W_2$ is finite dimensional because its basis contains only a finite amount of vectors.

$\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2)$

$k + m + p = k + m + k + p - k$

$k + m + p = k + m + p$ \square

b. Let W_1 and W_2 be finite-dimensional subspaces of a vector space V , and let $V = W_1 + W_2$. Deduce that V is the direct sum of W_1 and W_2 if and only if $\dim(V) = \dim(W_1) + \dim(W_2)$

$V = W_1 \oplus W_2 \Leftrightarrow \dim(V) = \dim(W_1) + \dim(W_2)$

Proof.

$V = W_1 \oplus W_2 \Rightarrow \dim(V) = \dim(W_1) + \dim(W_2)$

From the definition of direct sum, $W_1 \cap W_2 = \{0\}$

This means $\dim(W_1 \cap W_2) = 0$

From (a), we proved that $\dim(V) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2)$
 $= \dim(W_1) + \dim(W_2) - 0$

$$= \dim(W_1) + \dim(W_2)$$

$$\dim(V) = \dim(W_1) + \dim(W_2) \Rightarrow V = W_1 \oplus W_2$$

$$V = W_1 \oplus W_2 \text{ if and only if } V = W_1 + W_2 \text{ and } W_1 \cap W_2 = \{0\}$$

$V = W_1 + W_2$ is true by the definition of the problem.

$$\text{From part (a), } \dim(V) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2)$$

$$\text{Our antecedent is } \dim(V) = \dim(W_1) + \dim(W_2)$$

Setting the two equations equal to each other:

$$\dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2) = \dim(W_1) + \dim(W_2)$$

$$\dim(W_1 \cap W_2) = 0$$

$$\text{This means } W_1 \cap W_2 = \{0\}$$

$$\text{Thus, } \dim(V) = \dim(W_1) + \dim(W_2)$$

$$\text{Therefore, } V = W_1 \oplus W_2 \Leftrightarrow \dim(V) = \dim(W_1) + \dim(W_2) \quad \square$$

§4 D

§5 E

Let V be the vector space of sequences. Define the functions $T, U : V \rightarrow V$ by $T(a_1, a_2, \dots) = (a_2, a_3, \dots)$ and $U(a_1, a_2, \dots) = (0, a_1, a_2, \dots)$. T and U are called the left shift and right shift operators on V respectively.

- a. Prove that T and U are linear.

Proof.

T is linear if and only if $T(x + y) = T(x) + T(y)$ and $T(cx) = cT(x)$

Let $x, y \in V$ $c \in F$

$$x = (x_1, x_2, \dots) \quad y = (y_1, y_2, \dots)$$

$$x + y = (x_1 + y_1, x_2 + y_2, \dots)$$

$$T(x + y) = (x_2 + y_2, x_3 + y_3, \dots)$$

$$T(x) = (x_2, x_3, \dots)$$

$$T(y) = (y_2, y_3, \dots)$$

$$T(x) + T(y) = (x_2 + y_2, x_3 + y_3, \dots)$$

$$\text{Thus, } T(x + y) = T(x) + T(y)$$

$$x = (x_1, x_2, \dots)$$

$$cx = (cx_1, cx_2, \dots)$$

$$T(cx) = (cx_2, cx_3, \dots)$$

$$T(x) = (x_2, x_3, \dots)$$

$$cT(x) = (cx_2, cx_3, \dots)$$

$$\text{Thus, } T(cx) = cT(x)$$

Therefore, T is linear. The proof for U being linear is similar. \square

- b. T is onto but not one to one

Let $v \in V$

- c. U is one to one but not onto.

§6 F

Let S be the subspace of $M_{n \times n}(R)$ generated by all matrices of the form $AB - BA$ with A and B in $M_{n \times n}(R)$. Prove that $\dim(S) = n^2 - 1$. (You may want to use the trace together with the rank-nullity theorem)

Proof.

Trace is a linear transformation.

$$\text{Tr} : M_{n \times n}(R) \rightarrow R$$

The subspace S is defined as $\{AB - BA : A, B \in M_{n \times n}(R)\}$

$$\text{Tr}(AB - BA) = \text{Tr}(AB) - \text{Tr}(BA)$$

$$= \text{Tr}(AB) - \text{Tr}(AB)$$

$$= 0$$

All matrices that can be expressed as $AB - BA$ is in the null space of Tr . This means that $N(\text{Tr}) = S$.

The rank-nullity theorem states:

$$\dim(N(\text{Tr})) + \dim(R(\text{Tr})) = \dim(M_{n \times n}(R))$$

$$N(\text{Tr}) = S, \text{ so } \dim(S) + \dim(R(\text{Tr})) = \dim(M_{n \times n}(R))$$

$$\dim(S) = \dim(M_{n \times n}(R)) - \dim(R(\text{Tr}))$$

$$= n^2 - \dim(R)$$

$$= n^2 - 1$$

□

§7 G

Let T be a linear transformation of a vector space V into itself. Suppose that $x \in V$ is such that $T^m(x) = 0$, and $T^{m-1}(x) \neq 0$ for some positive m . Show that $x, T(x), T^2(x), \dots, T^{m-1}(x)$ are linearly independent.

Proof.

The linear combination of the above set is

$$a_0x + a_1T(x) + a_2T^2(x) + \dots + a_{m-1}T^{m-1}(x) = 0$$

Notice that $T^n(x) = 0$ for all $n > m$.

$$T^{m+1}(x) = T(T^m(x)) = T(0) = 0$$

Let's take T^{m-1} on both sides of the linear combination.

$$\begin{aligned} T^{m-1}(a_0x + a_1T(x) + a_2T^2(x) + \dots + a_{m-1}T^{m-1}(x)) &= T^{m-1}(0) \\ T^{m-1}(a_0x) &= 0 \end{aligned}$$

So $z_1 = \sum_{i=1}^n a_i x_i$ where $a \in F$ and $x \in S_1$

If $S_1 \subseteq S_2$, then $x \in S_2$

So $z_1 \in \text{span}(S_2)$ because we can write z_1 as a linear combination of S_2

Therefore, if $S_1 \subseteq S_2$ then $\text{span}(S_1) \subseteq \text{span}(S_2)$ (*)

Defined in the problem, $\text{span}(S_1) = V$

By (*), $\text{span}(S_1) = V \subseteq \text{span}(S_2)$

Using theorem 1.5, $\text{span}(S_2) \subseteq V$

Therefore, $\text{span}(S_2) \subseteq V \subseteq \text{span}(S_2) \Leftrightarrow V = \text{span}(S_2)$

□

§8 H

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$

- a. If $T(a,b,c) = (a,b,0)$, show that T is the projection on the xy -plane along the z -axis.

Proof.

We want the projection to be on the xy -plane along the z -axis. Let the projection be $(x,y,0)$. To minimize the distance, we must choose x and y such that

$$(a-x)^2 + (b-y)^2 + (c-0)^2$$

is minimum. Since the equation above is a difference of squares, $x = a$ and $y = b$ will give us the minimum value. Therefore, the projection on the xy -plane will be $(a,b,0)$, which is T . \square

- b. Find a formula for $T(a,b,c)$, where T represents the projection on the z -axis along the xy -plane.

Proof.

We want the projection to be on the z -axis along the xy -plane. Let the projection be $(0,0,z)$. To minimize the distance, we must choose z such that

$$(a-0)^2 + (b-0)^2 + (c-z)^2$$

is minimum. $z = c$ will give us the minimum value. Therefore, the equation for T will be $T(a,b,c) = (0,0,c)$. \square

- c. If $T(a,b,c) = (a-c,b,0)$, show that T is the projection on the xy -plane along the line $L = \{(a, 0, a) : a \in \mathbb{R}\}$

Proof.

We want the projection to be on the xy -plane along the line L . Let the projection be $(x, y, 0)$. A vector that is on L is $(1, 0, 1)$. To minimize the distance, we must choose λ such that

$$(a, b, c) + \lambda(1, 0, 1) = (x, y, 0)$$

is minimum. Writing the equation above as a system:

$$a + \lambda = x$$

$$b = y$$

$$c + \lambda = 0$$

Solving this system gives us, $x = a - c, y = b$

Therefore, the projection on the xy -plane along the line L will be $(a - c, b, 0)$. \square

§9 I

In $M_{m \times n}(F)$, let E^{ij} denote the matrix whose only nonzero entry is 1 in the i th row and j th column. Prove that $\{E^{ij} : 1 \leq i \leq m, 1 \leq j \leq n\}$ is linearly independent.

Proof.

If E^{ij} is linearly independent then $a_{1,1}E^{1,1} + \dots + a_{m,n}E^{m,n} \neq 0$

This sum can only equal the 0 matrix if all a are 0.

Therefore, E^{ij} is linearly independent. \square

§10 J

Let u and v be distinct vectors in a vector space V . Show that $\{u, v\}$ is linearly dependent if and only if u or v is a multiple of the other.

Proof.

Let's first show that if u or v is a multiple of the other then $\{u, v\}$ is linearly dependent.

Being a multiple means $u = nv$ or $v = nu$ where $n \in F$

If $\{u, v\}$ is linearly dependent then $a_1u + a_2v = 0$ where $a \in F$

Using definition of multiple $a_1u + a_2nu = 0$

Factoring, $u(a_1 + a_2n) = 0$

This means $(a_1 + a_2n) = 0$

So, $n = \frac{-a_1}{a_2}$ which is a solution for linearly dependency.

Without loss of generality, we can prove the case where $v = nu$

Therefore, $\{u, v\}$ is linearly dependent.

Now let's show that if $\{u, v\}$ is linearly dependent then u or v is a multiple of the other.

If $\{u, v\}$ is linearly dependent then $a_1u + a_2v = 0$ where $a \in F$

We can rewrite the equation above as $a_1u = -a_2v$

$$u = \frac{-a_2}{a_1}v$$

Thus, u is a multiple of v .

Without loss of generality, we can prove v is a multiple of u .

Therefore, u or v is a multiple of the other. □