

Linear Algebra

Dan

Spring 2020

§1

Let

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ and } \mathbf{b} = \begin{pmatrix} e \\ f \end{pmatrix} \quad (1)$$

- a. Suppose that $a \neq 0$, compute the solution of $\mathbf{Ax} = \mathbf{b}$ using row reduction and provide the conditions on a, b, c, d such that your computations are valid. Express the result as a simplified expression. (**Hint:** recall that you can not divide by zero)

Proof. We perform reduced row echelon form (rref) on the augmented matrix

$$\begin{aligned} (A|b) &= \left[\begin{array}{cc|c} a & b & e \\ c & d & f \end{array} \right] \\ R_2 &\leftarrow R_2 - \frac{c}{a}R_1 \left[\begin{array}{cc|c} a & b & e \\ 0 & d - \frac{cb}{a} & f - \frac{ce}{a} \end{array} \right] \\ &\left[\begin{array}{cc|c} a & b & e \\ 0 & \frac{ad-cb}{a} & \frac{af-ce}{a} \end{array} \right] \\ R_2 &\leftarrow \frac{a}{ad-cb}R_2 \quad \text{Assuming that } ad-cb \neq 0 \left[\begin{array}{cc|c} a & b & e \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow R_1 - bR_2 \left[\begin{array}{cc|c} a & 0 & e - b\frac{af-ce}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow \frac{R_1}{a} \left[\begin{array}{cc|c} 1 & 0 & \frac{1}{a}(e - b\frac{af-ce}{ad-cb}) \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ &\left[\begin{array}{cc|c} 1 & 0 & \frac{de-bf}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \end{aligned}$$

$$\mathbf{x} = \begin{bmatrix} \frac{de-bf}{ad-cb} \\ \frac{af-ce}{ad-cb} \end{bmatrix} = \frac{1}{ad-cb} \begin{bmatrix} de-bf \\ af-ce \end{bmatrix} \text{ where } ad-cb \neq 0$$

□

- b. If $a = 0$, and $c \neq 0$, is your above computation still valid? How would you modify it? (explain briefly) (**Hint:** recall that you can swap the equations and the result is the same)

Proof. If $a = 0$, and $c \neq 0$, then the above computation will not be valid as we divided by a multiple times when we computed the rref. I would swap the first and second rows so that it would look like

$$\left[\begin{array}{cc|c} c & d & f \\ 0 & b & e \end{array} \right]$$

and compute the rref, assuming that $b \neq 0$. We obtain the rref,

$$\left[\begin{array}{cc|c} 1 & 0 & \frac{bf-de}{bc} \\ 0 & 1 & \frac{e}{b} \end{array} \right]$$

and the solution

$$x = \begin{bmatrix} \frac{bf-de}{bc} \\ \frac{e}{b} \end{bmatrix} \quad \text{where } b \neq 0$$

□

- c. If $a = 0$, $c = 0$, but $b \neq 0$, $d \neq 0$, what are the conditions on e and f such that the system $\mathbf{Ax} = \mathbf{b}$ has a solution? Is the solution unique? (**Hint:** recall that $\mathbf{Ax} = \mathbf{b}$ has a solution if and only if \mathbf{b} can be written as a linear combination of the columns of \mathbf{A})

Proof. If $a = 0$, $c = 0$, $b \neq 0$, $d \neq 0$, we get the augmented matrix

$$\left[\begin{array}{cc|c} 0 & b & e \\ 0 & d & f \end{array} \right]$$

Performing row reduction,

$$\left[\begin{array}{cc|c} 0 & 1 & \frac{e}{b} \\ 0 & 1 & \frac{f}{d} \end{array} \right]$$

Having an infinite amount of solutions is by definition another way of saying that a system that is consistent and that the solutions are not unique.

It is clear that $\det(\mathbf{A}) = 0$ by multiplying the diagonal entries because it is an upper triangular matrix. This means that the solution, if it exists, is not unique by Theorem 3.10 and the corollary to Theorem 4.7. A system is consistent if and only if $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}|\mathbf{b})$ by Theorem 3.11. For this condition to hold,

$$\mathbf{b} \in \text{span}(\mathbf{A}) \Leftrightarrow \begin{pmatrix} \frac{e}{b} \\ \frac{f}{d} \end{pmatrix} = c_1 \begin{pmatrix} 0 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \text{where } c_1, c_2 \in F$$

So the condition of the solution is,

$$\begin{aligned} c_2 &= \frac{e}{b} \\ c_2 &= \frac{f}{d} \\ \frac{e}{b} &= \frac{f}{d} \end{aligned}$$

Thus, there exists a infinite amount of solution.

□

- d. Solve the system

$$\begin{bmatrix} \sqrt{2} & 3\sqrt{2} \\ 2\sqrt{2} & \sqrt{2} \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 5\sqrt{2} \\ 5\sqrt{2} \end{pmatrix}. \quad (2)$$

(**Hint:** You may want to use the formula you just deduced)

Proof.

$$\begin{aligned}
 x_1 &= \frac{de - bf}{ad - cb} \\
 &= \frac{\sqrt{2}(5\sqrt{2}) - 3\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\
 &= \frac{10 - 30}{2 - 12} \\
 &= \frac{-20}{-10} \\
 &= 2
 \end{aligned}$$

$$\begin{aligned}
 x_2 &= \frac{af - ce}{ad - cb} \\
 &= \frac{\sqrt{2}(5\sqrt{2}) - 2\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\
 &= \frac{10 - 20}{-10} \\
 &= 1
 \end{aligned}$$

□

§2

Let

$$\mathbf{A} = \begin{bmatrix} 0 & -\alpha & 2 & 0 \\ 1 & 1 & 0 & 1 \\ 2 & 2 & 2 & 3 \\ -2 & -2 & 4 & 2\alpha \\ 0 & \alpha & -1 & 2\alpha + 1/2 \end{bmatrix}, \quad \text{and } \mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ 4 \\ 2 + \alpha \\ 2\beta + \alpha - 2 \end{bmatrix} \quad (3)$$

What are the conditions on α and β such that the system $\mathbf{Ax} = \mathbf{b}$:

- Has no solution?

Proof. We begin by putting the augmented matrix $(\mathbf{A}|\mathbf{b})$ in its reduced form.

$$\begin{aligned}
 (\mathbf{A}|\mathbf{b}) &= \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 2 & 2 & 2 & 3 & 4 \\ -2 & -2 & 4 & 2\alpha & 2+\alpha \\ 0 & \alpha & -1 & 2\alpha+1/2 & 2\beta+\alpha-2 \end{array} \right] \\
 R_5 \leftarrow R_5 + R_1 & \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 2 & 2 & 2 & 3 & 4 \\ -2 & -2 & 4 & 2\alpha & 2+\alpha \\ 0 & 0 & 1 & 2\alpha+1/2 & 2\beta+\alpha \end{array} \right] \\
 R_3 \leftarrow R_3 - 2R_2 & \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ -2 & -2 & 4 & 2\alpha & 2+\alpha \\ 0 & 0 & 1 & 2\alpha+1/2 & 2\beta+\alpha \end{array} \right] \\
 R_4 \leftarrow R_4 + 2R_2 & \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 4 & 2\alpha+2 & 4+\alpha \\ 0 & 0 & 1 & 2\alpha+1/2 & 2\beta+\alpha \end{array} \right] \\
 R_4 \leftarrow R_4 - 2R_3, R_5 \leftarrow R_5 - \frac{1}{2}R_3 & \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 2\alpha & 2\beta+\alpha+1 \end{array} \right] \\
 R_5 \leftarrow R_5 - R_4 & \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 2\beta-1 \end{array} \right] \\
 R_1 \leftrightarrow R_2 \quad (\mathbf{A}'|\mathbf{b}') &= \left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & -\alpha & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 2\beta-1 \end{array} \right]
 \end{aligned}$$

By Theorem 3.11 and 3.13, a system is consistent if and only if $\text{rank}(\mathbf{A}') = \text{rank}(\mathbf{A}'|\mathbf{b}')$. Thus this system will have no solution if $2\beta - 1 \neq 0$ because then $\mathbf{b}' \notin \text{span}(\mathbf{A}')$.

This is when $\beta \neq \frac{1}{2}$. We observe that there will be no conditions on α . \square

- b. Has an unique solution? Find the solution. (**Hint:** you will need to row reduce the augmented system to echelon form, and then use the theorems seen in class to impose the conditions on α and β).

Proof. Combining Theorem 3.10 and the corollary to Theorem 4.7, we get that a system has a unique solution if and only if $\det(\mathbf{A}) \neq 0$. The determinant of an upper triangular matrix is the product of its diagonal entries by property 4 of determinants in section 4.4. Using this

fact we can compute the condition of α as such that

$$\begin{aligned} 1 * -\alpha * 2 * 2\alpha &\neq 0 \\ -4\alpha^2 &\neq 0 \\ \alpha &\neq 0 \end{aligned}$$

and from (a), $\beta = \frac{1}{2}$. Combining these two conditions we get the following system,

$$\left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & -\alpha & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

By performing back substitution we compute the unique solution

$$\begin{aligned} 2\alpha x_4 &= \alpha \Leftrightarrow x_4 = \frac{1}{2} \\ 2x_3 + x_4 &= 2 \Leftrightarrow x_3 = \frac{3}{4} \\ -\alpha x_2 + 2x_3 &= 2 \Leftrightarrow x_2 = -\frac{1}{2\alpha} \\ x_1 + x_2 + x_4 &= 1 \Leftrightarrow x_1 = \frac{1}{2} + \frac{1}{2\alpha} \end{aligned}$$

$$x = \begin{bmatrix} \frac{1}{2} + \frac{1}{2\alpha} \\ -\frac{1}{2\alpha} \\ \frac{3}{4} \\ \frac{1}{2} \end{bmatrix}$$

as desired. □

- c. Has infinite amount of solutions? Find the solution set in parametric form. (**Hint:** You may have one equations for α and one for β that have to be satisfied simultaneously).

Proof. Having an infinite amount of solutions is by definition another way of saying that a system that is consistent and that the solutions are not unique. A system is consistent if and only if $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}|\mathbf{b})$ by Theorem 3.11. If $\det(\mathbf{A}) = 0$, the solution, if it exists, is not unique by Theorem 3.10 and the corollary to Theorem 4.7.

Using this fact we can compute the condition of α as such that

$$\begin{aligned} 1 * -\alpha * 2 * 2\alpha &= 0 \\ -4\alpha^2 &= 0 \\ \alpha &= 0 \end{aligned}$$

Given that $\alpha = 0$, and that $\beta = \frac{1}{2}$ from part (a) we get the following system,

$$(\mathbf{A}'|\mathbf{b}') = \left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

It is clear that $\text{rank}(\mathbf{A}') = \text{rank}(\mathbf{A}'|\mathbf{b}')$ because \mathbf{b}' is a linear combination of the second and third column from \mathbf{A}' . By Theorem 3.13, we know that the solution set for $\mathbf{A}'\mathbf{x} = \mathbf{b}'$ is equivalent to the solution set for $\mathbf{A}\mathbf{x} = \mathbf{b}$. Hence, both systems are consistent.

We can compute a solution space to $\mathbf{A}\mathbf{x} = \mathbf{b}$ as outlined in Theorem 3.9. We start by first computing the solution set to $\mathbf{A}\mathbf{x} = 0$ denoted by K_H . It is clear that $\text{rank}(\mathbf{A}) = 3$ because the first two columns are the same and the rest of the columns are linearly independent from each other. By Theorem 3.8, $\dim(K_H) = 4 - 3 = 1$. Thus any nonzero solution constitutes a basis for K . For example, since

$$\begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}$$

is a solution to the $\mathbf{A}\mathbf{x} = 0$, it is a basis for K_H by Corollary 2 of Theorem 1.10. So a solution set to K_H would be

$$K_H = \left\{ t \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix} : t \in \mathbb{R} \right\}$$

A solution to $\mathbf{A}\mathbf{x} = \mathbf{b}$ is

$$\begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

Therefore, by Theorem 3.9 we compute solution space when this system has infinite amount of solutions, which is when $\alpha = 0$ and $\beta = \frac{1}{2}$ as

$$K = \left\{ \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix} : t \in \mathbb{R} \right\}$$

□

§3

Let $A \in M_{n \times n}(F)$, for a field F . We want to prove that $\text{rank}(A^2) - \text{rank}(A^3) \leq \text{rank}(A) - \text{rank}(A^2)$. The solution to this exercise requires the notion of quotient spaces. Even though you should already be familiar with quotient spaces we will prove a few properties that will be useful.

Let W be a subspace of a vector space V over a field F . For any $v \in V$ the set $\{v\} + W = \{v + w : w \in W\}$ is called the coset of W containing v . It is customary to denote this coset by $v + W$ rather than $\{v\} + W$. Following this notation we write $V/W = \{v + W : v \in V\}$, which is usually called the quotient space V module W . Addition and scalar multiplication by scalars can be defined in the collection V/W as follows:

$$(v_1 + W) + (v_2 + W) = (v_1 + v_2) + W$$

for all $v_1, v_2 \in V$ and

$$a(v + W) = av + W$$

for all $v \in V$ and $a \in F$

- a. Prove that $v_1 + W = v_2 + W$ if and only if $v_1 - v_2 \in W$. (**Hint:** recall that $v_1 + W$ is a set, thus you need to prove equality between sets)

Proof.

i. $v_1 + W = v_2 + W \Rightarrow v_1 - v_2 \in W$

If $v_1 + W = v_2 + W$, then $\exists w_1, w_2 \in W$ such that $v_1 + w_1 = v_2 + w_2$

$$v_1 - v_2 = w_2 - w_1$$

Since, $w_2 - w_1 \in W$ (closure under addition)

Therefore, $v_1 - v_2 \in W$

ii. $v_1 - v_2 \in W \Rightarrow v_1 + W = v_2 + W$

This means $v_1 - v_2 = w$ where $w \in W$ (*)

Now let $x \in v_1 + W$

By definition, $\exists w_x \in W$ such that $x = v_1 + w_x$

By (*) $v_1 = v_2 + w$

So, $x = v_2 + w + w_x$

Since, $w + w_x \in W$ (closure under addition)

We have $x \in v_2 + W$

So, $v_1 + W \subseteq v_2 + W$

Without loss of generality, we can show $v_2 + W \subseteq v_1 + W$

Therefore, $v_1 + W = v_2 + W$

Therefore, $v_1 + W = v_2 + W$ if and only if $v_1 - v_2 \in W$

□

- b. Show that V/W with the operations defined above is a linear vector space.

VS 1: For all x, y in V/W , $x + y = y + x$ (commutativity of addition)

Proof. Let $x = v_x + W, y = v_y + W$ where $v_x, v_y \in V$.

$$\begin{aligned} x + y &= (v_x + W) + (v_y + W) \\ &= (v_x + v_y) + W \\ &= (v_y + v_x) + W \\ &= (v_y + W) + (v_x + W) \\ &= y + x \end{aligned}$$

□

VS 2: For all x, y, z in V/W , $(x + y) + z = x + (y + z)$ (associativity of addition)

Proof. Let $x = v_x + W, y = v_y + W, z = v_z + W$, where $v_x, v_y, v_z \in V$.

$$\begin{aligned} (x + y) + z &= ((v_x + W) + (v_y + W)) + (v_z + W) \\ &= ((v_x + v_y) + W) + (v_z + W) \\ &= (v_y + v_x + v_z) + W \\ &= (v_x + W) + ((v_y + v_z) + W) \\ &= (v_x + W) + ((v_y + W) + (v_z + W)) \\ &= x + (y + z) \end{aligned}$$

□

VS 3: There exists an element in V/W denoted by $\mathbf{0}$ such that $x + \mathbf{0} = x$ for each x in V/W

Proof. Let $x = v_x + W$ where $v_x \in V$. Let $\mathbf{0}$ in V/W be defined as $0 + W$, where $0 \in V$.

$$\begin{aligned} x + \mathbf{0} &= x + (0 + W) \\ &= (v_x + W) + (0 + W) \\ &= (v_x + 0) + W \\ &= v_x + W \\ &= x \end{aligned}$$

□

VS 4: For each element x in V/W there exists an element y in V/W such that $x + y = \mathbf{0}$

Proof. Let $x = v_x + W$ where $v_x \in V$. Fix y such that $y = -v_x + W$.

$$\begin{aligned} x + y &= (v_x + W) + (-v_x + W) \\ &= (v_x - v_x) + W \\ &= 0 + W \\ &= \mathbf{0} \end{aligned}$$

□

VS 5: For each element x in V/W , $1x = x$

Proof. Let $x = v_x + W$ where $v_x \in V$.

$$\begin{aligned} 1x &= 1(v_x + W) \\ &= (1v_x + W) \\ &= (v_x + W) \\ &= x \end{aligned}$$

□

VS 6: For each pair of elements a, b in \mathbb{F} and each element x in V/W , $(ab)x = a(bx)$

Proof. Let $x = v_x + W$ where $v_x \in V$.

$$\begin{aligned} (ab)x &= abv_x + W \\ &= a(bv_x + W) \\ &= a(bx) \end{aligned}$$

□

VS 7: For each element a in \mathbb{F} and each pair of elements x, y in V/W , $a(x + y) = ax + ay$

Proof. Let $x = v_x + W, y = v_y + W$ where $v_x, v_y \in V$.

$$\begin{aligned} a(x + y) &= a((v_x + v_y) + W) \\ &= ((av_x + av_y) + W) \\ &= (av_x + W) + (av_y + W) \\ &= a(v_x + W) + a(v_y + W) \\ &= ax + ay \end{aligned}$$

□

VS 8: For each pair of elements a, b in \mathbb{F} and each element x in V/W , $(a + b)x = ax + bx$

Proof. Let $x = v_x + W$ where $v_x \in V$.

$$\begin{aligned}
 (a + b)x &= (a + b)(v_x + W) \\
 &= ((a + b)v_x) + W \\
 &= (av_x + bv_x) + W \\
 &= (av_x + W) + (bv_x + W) \\
 &= a(v_x + W) + b(v_x + W) \\
 &= ax + bx
 \end{aligned}$$

□

Therefore, V/W is a vector space because it holds all the properties above.

- c. Prove that if $\dim(V) < \infty$ then $\dim(V/W) = \dim(V) - \dim(W)$. (Hint: Define a linear map $T : V \rightarrow V/W$ such that the range of T is V/W , and then use the rank-nullity theorem)

Proof. We define the linear map $T : V \rightarrow V/W$ by

$$T(v) = v + W$$

We first prove that T is in fact linear, where $v_1, v_2 \in V$ and $c \in F$.

$$\begin{aligned}
 T(cv_1 + v_2) &= (cv_1 + v_2) + W \\
 &= (cv_1 + W) + ((v_2) + W) \\
 &= c(v_1 + W) + ((v_2) + W) \\
 &= cT(v_1) + T(v_2)
 \end{aligned}$$

We claim that $N(T) = W$ and $R(T) = V/W$

1. $R(T) = V/W$

- i. $R(T) \subseteq V/W$
By Theorem 2.1

- ii. $V/W \subseteq R(T)$

Let $y = v_y + W$ where $y \in V/W$ and $v_y \in V$.

For prove that $y \in N(T)$ we need to show that $\exists x$ such that $T(x) = y$.

Notice that $x = v_y$. Therefore, $y \in R(T)$ so $V/W \subseteq R(T)$.

2. $N(T) = W$

- i. $N(T) \subseteq W$

Lemma 3.1

$w_1 + W = w_2 + W$ where $w_1, w_2 \in W$.

Proof.

Let $x_1 \in w_1 + W$ and $x_2 \in w_2 + W$

By definition, $w_1 + W = \{w_1 + w : w \in W\}$.

We know that $w_1 + w \in W$ because W is a subspace and has closure under addition. Hence, $x_1 \in W$ and similarly $x_2 \in W$. Therefore, $w_1 + W \subseteq W$ and $w_2 + W \subseteq W$.

Now consider $x_1 \in W$ and $x_2 \in W$.

$x_1 \in w_1 + W$ because $w_1 + w \in W$ by closure under addition

Similarly, $x_2 \in w_2 + W$.

Hence, $W \subseteq w_1 + W$ and $W \subseteq w_2 + W$.

Therefore, $w_1 + W = W$ and $W = w_2 + W$ so, $w_1 + W = w_2 + W$.

□

Let $x \in N(T)$.

By definition of $N(T)$ and using the lemma above,

$$\begin{aligned} T(x) &= \mathbf{0} \\ &= 0 + W \\ &= w + W \end{aligned}$$

where $0, w \in W$ and $\mathbf{0} \in V/W$.

This must mean that $x = w$ and since $w \in W$, $x \in W$. Therefore, $N(T) \subseteq W$

ii. $W \subseteq N(T)$

Let $w \in W$.

$$\begin{aligned} T(w) &= w + W \\ &= W \\ &= 0 + W \\ &= \mathbf{0} \end{aligned}$$

So $w \in N(T)$. Therefore, $W \subseteq N(T)$.

Notice that $V/W \subseteq V$ so V/W is finite dimensional. Since T is a linear map and that V and V/W are indeed finite dimensional vector spaces (by part b) we can use the rank-nullity theorem (Theorem 2.3).

$$\begin{aligned} \dim(N(T)) + \dim(R(T)) &= \dim(V) \\ \dim(W) + \dim(V/W) &= \dim(V) \\ \dim(V/W) &= \dim(V) - \dim(W) \end{aligned}$$

as desired. □

d. Let $K = F^n$, define $AK = R(L_A)$, and $A^2K = R(L_{A^2})$. Show that AK/A^2K is a vector space of dimension $\text{rank}(A) - \text{rank}(A^2)$.

Proof. We begin by proving that AK/A^2K is a vector space. It is enough to show that A^2K is a subspace of AK to prove that AK/A^2K is a vector space, by part (b). It is clear that AK is a subspace because $AK = R(L_A)$ and Theorem 2.3. Moreover, $A^2K \subseteq AK$ because $R(L_{A^2}) \subseteq R(L_A)$. (Theorem 3.7?) **Not sure if $R(L_{A^2}) \subseteq R(L_A)$ holds**
Because $A^2K = R(L_{A^2})$, A^2K has the properties of a vector space because of Theorem 2.1. It follows from part (c) that

$$\begin{aligned} \dim(AK/A^2K) &= \dim(AK) - \dim(A^2K) \\ &= \dim(R(L_A)) - \dim(R(L_{A^2})) \\ &= \text{rank}(A) - \text{rank}(A^2) \end{aligned}$$

as desired. □

- e. Show that A^2K/A^3K is a vector space of dimension $\text{rank}(A^2) - \text{rank}(A^3)$, where $A^3K = R(L_{A^3})$.

Proof. Similiar to what we did in part (d), we can prove that A^2K/A^3K is a vector space. Then it follows that,

$$\begin{aligned} \dim(A^2K/A^3K) &= \dim(A^2K) - \dim(A^3K) \\ &= \dim(R(L_{A^2})) - \dim(R(L_{A^3})) \\ &= \text{rank}(A^2) - \text{rank}(A^3) \end{aligned}$$

as desired. □

- f. Define $T : AK/A^2K \rightarrow A^2K/A^3K$, by $T(v) = L_A(v)$, i.e, we left multiply each element of v by the matrix A . Show that $R(T) = A^2K/A^3K$.

Proof. To show that $R(T) = A^2K/A^3K$, is another way of saying show that T is onto. We must show that $\forall x \in AK/A^2K [\exists y \in A^2K/A^3K : T(x) = y]$. Let $x = Ak' + A^2K$ where $k' \in \mathbb{F}^n$.

$$\begin{aligned} T(x) &= T(Ak' + A^2K) \\ &= L_A(Ak' + A^2K) \\ &= L_A(Ak') + L_A(A^2K) \\ &= A^2k' + A^3K \in A^2K/A^3K \end{aligned}$$

Therefore, T is onto and $R(T) = A^2K/A^3K$. □

- g. Use the rank-nullity theorem on T to conclude that $\text{rank}(A^2) - \text{rank}(A^3) \leq \text{rank}(A) - \text{rank}(A^2)$.

Proof. We begin by proving that T is linear. We define $x_1 = Ak'_1 + A^2K$ and $x_2 = Ak'_2 + A^2K$ such that $x_1, x_2 \in AK/A^2K$ and $k'_1, k'_2 \in \mathbb{F}^n$ and $c \in F$. We use properties of matrices as

outlined in Theorem 2.12.

$$\begin{aligned}
 T(ck'_1 + k'_2) &= L_A(ck'_1 + k'_2) \\
 &= A(ck'_1 + k'_2) \\
 &= A(c(Ak'_1 + A^2K) + (Ak'_2 + A^2K)) \\
 &= A(c(Ak'_1 + A^2K)) + A(Ak'_2 + A^2K) \\
 &= c(A(Ak'_1 + A^2K)) + A(Ak'_2 + A^2K) \\
 &= cL_A(k'_1) + L_A(k'_2) \\
 &= cT(k'_1) + T(k'_2)
 \end{aligned}$$

Since T is linear, we can use rank nullity theorem. From part (d), (e), (f), and assuming that $\text{nullity}(T) \geq 0$ because the number of elements of a set cannot be negative

$$\begin{aligned}
 \text{rank}(T) + \text{nullity}(T) &= \dim(AK/A^2K) \\
 &= \text{rank}(A) - \text{rank}(A^2) \\
 \text{rank}(T) &\leq \text{rank}(A) - \text{rank}(A^2) \\
 \text{rank}(A^2) - \text{rank}(A^3) &\leq \text{rank}(A) - \text{rank}(A^2)
 \end{aligned}$$

as desired. □

§4

Let V be a finite-dimensional vector space. Let T and P be two linear transformations from V to itself, such that $T^2 = P^2 = 0$, and $T \circ P + P \circ T = I$, where I is the identity in V .

- a. Denote $N_T = N(T)$ and $N_P = N(P)$, the null spaces of T and P , respectively. Show that $N_P = P(N_T)$, and $N_T = T(N_P)$, where $T(N_P) = \{T(v) : v \in N_P\}$ and $P(N_T) = \{P(v) : v \in N_T\}$.

Proof.

- i. Show $N_P = P(N_T)$

1. $N_P \subseteq P(N_T)$

Let $x \in N_P$. By definition, $P(x) = 0$.

$$\begin{aligned}
 P \circ T(x) + T \circ P(x) &= x \\
 P \circ T(x) + T(0) &= x \\
 P \circ T(x) &= x
 \end{aligned}$$

Notice that $T(x) \in N_T$, so $x \in P(N_T)$. Thus, $N_P \subseteq P(N_T)$.

2. $P(N_T) \subseteq N_P$

Let $P(x) \in P(N_T)$ where $x \in N_T$.

$P \circ P(x) = 0$ so $P(x) \in N_P$. Thus, $P(N_T) \subseteq N_P$.

Therefore, $N_P = P(N_T)$.

- ii. Show $N_T = T(N_P)$

1. $N_T \subseteq T(N_P)$

Let $x \in N_T$. By definition, $T(x) = 0$.

$$P \circ T(x) + T \circ P(x) = x$$

$$P(0) + T \circ P(x) = x$$

$$T \circ P(x) = x$$

Notice that $P(x) \in N_P$, so $x \in T(N_P)$. Thus, $N_T \subseteq T(N_P)$.

2. $T(N_P) \subseteq N_T$

Let $T(x) \in T(N_P)$ where $x \in N_P$.

$T \circ T(x) = 0$ so $T(x) \in N_T$. Thus, $T(N_P) \subseteq N_T$.

Therefore, $N_P = P(N_T)$.

□

b. Show that $V = N_T \oplus N_P$.

Proof. Need to prove the following two conditions.

i. $N_T \cap N_P = \{0\}$.

Let $x \in N_T$ and $x \in N_P$. This means that $T(x) = 0$ and $P(x) = 0$.

$$\begin{aligned} x &= P \circ T(x) + T \circ P(x) \\ &= P(0) + T(0) \\ &= 0 \end{aligned}$$

Thus, $N_T \cap N_P = \{0\}$.

ii. $N_T + N_P = V$.

1. $N_T + N_P \subseteq V$

Let $x \in N_T + N_P$. Define $x = n_t + n_p$ where $n_t \in N_T \subseteq V$ and $n_p \in N_P \subseteq V$ by Theorem 2.11. Since V is a vector space $n_t + n_p \in V$ by closure under addition.

Thus, $x \in V$ and it follows that $N_T + N_P \subseteq V$.

2. $V \subseteq N_T + N_P$

Let $x \in V$. We know that $x = P \circ T(x) + T \circ P(x)$.

Notice that $P \circ T(x) \in N_P$ because $P(P \circ T(x)) = 0$.

Similarly, $T \circ P(x) \in N_T$ because $T(T \circ P(x)) = 0$.

Thus, $N_T + N_P \subseteq V$.

Therefore, $N_T + N_P = V$.

Therefore, by definition $V = N_T \oplus N_P$.

□

c. Prove that the dimension of V is even.

Proof.

Lemma 4.1

$$V = W_1 \oplus W_2 \Leftrightarrow \dim(V) = \dim(W_1) + \dim(W_2)$$

Proof. 1.6.29(b) in the textbook **Actually remember to type this in**

□

By the lemma above, $\dim(V) = \dim(N_T) + \dim(N_P)$.

Let's construct a basis β_{N_T} for N_T and β_{N_P} for N_P .

$$\beta_{N_T} = \{u_1, u_2, \dots, u_m\}$$

$$\beta_{N_P} = \{w_1, w_2, \dots, w_n\}$$

From (a), since $N_T = T(N_P)$, this must mean that

$$\text{span}(\beta_{N_T}) = T(\text{span}(\beta_{N_P}))$$

Thus, $\dim(N_P) \leq \dim(N_T)$ because $\dim(T(N_P)) \leq \dim(N_P)$.

Likewise, since $N_P = P(N_T)$, this must mean that

$$\text{span}(\beta_{N_P}) = P(\text{span}(\beta_{N_T}))$$

Thus, $\dim(N_T) \leq \dim(N_P)$ because $\dim(P(N_T)) \leq \dim(N_T)$.

Thus, $\dim(N_P) = \dim(N_T) = n = m$.

$$\begin{aligned} \dim(V) &= \dim(N_T) + \dim(N_P) \\ &= n + m \\ &= n + n = m + m \\ &= 2n = 2m \end{aligned}$$

Therefore, $\dim(V)$ is even. □

d. Suppose that the dimension of V is two. Prove that V has a basis β , such that

$$[T]_{\beta} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad [P]_{\beta} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}. \quad (4)$$

Proof. □